

SEMISTABILITY OF LOGARITHMIC COTANGENT BUNDLE ON SOME PROJECTIVE MANIFOLDS

SESHADRI CHINTAPALLI AND JAYA NN IYER

ABSTRACT. In this paper, we investigate the semistability of logarithmic de Rham sheaves on a smooth projective variety (X, D) , under suitable conditions. This is related to existence of Kähler -Einstein metric on the open variety. We investigate this problem when the Picard number is one. Fix a normal crossing divisor D on X and consider the logarithmic de Rham sheaf $\Omega_X(\log D)$ on X . We prove semistability of this sheaf, when the log canonical sheaf $K_X + D$ is ample or trivial, or when $-K_X - D$ is ample i.e., when X is a log Fano n -fold of dimension $n \leq 6$. We also extend the semistability result for Kawamata coverings, and this gives examples whose Picard number can be greater than one.

CONTENTS

1. Introduction	1
2. Semistability and logarithmic vanishing theorems	3
3. Stability when the sheaf $K_X + \mathcal{O}_X(D)$ is non-negative.	5
4. Stability on Kawamata's finite coverings	6
5. Log Fano manifolds of small dimensions	7
References	14

1. INTRODUCTION

Suppose X is a smooth complex projective variety of dimension n and fix an ample line bundle $\mathcal{O}_X(1)$ on X . In this paper, for any coherent sheaf \mathcal{F} of X , we consider the Mumford-Takemoto stability or μ -stability of \mathcal{F} , see §2.1. .

We would like to investigate the stability properties of some natural bundles on the projective manifold X . In particular, the stability of the cotangent bundle has attracted wide attention since work of Aubin [Au] and Yau [Ya], appeared in the study of existence of a Kähler-Einstein metric on smooth projective varieties, with ample or trivial canonical class. By [Kb], [Lu], it implies the stability of cotangent bundle in these cases. Since

⁰Mathematics Classification Number: 53C55, 53C07, 53C29, 53.50.

⁰Keywords: Logarithmic Fano manifolds, Logarithmic Cotangent bundle, Semistability.

then the stability problem for Fano manifolds has attracted wide attention. Some significant results on stability is done by Steffens [St], Subramanian [Su], Peternell-Wisniewski [Pe-Wi], Hwang [Hw] and the references therein.

Suppose we fix a simple normal crossing divisor D on X . Then it is of interest to investigate semistability of the logarithmic cotangent bundle $\Omega_X(\log D)$. Existence of a Kähler-Einstein metric on the open variety $X - D$ with singularity on the boundary has been investigated by Tian [Ti3]. A recent work on this can be found in [Li-Su] and is related to K-stability. This is perhaps related to the μ -stability of the logarithmic cotangent bundle, although this has not been established yet. In other direction, certain Chern inequalities have been obtained in [Ti3] for log general type pairs (X, D) and Bogomolov-Miyaoka-Yau inequality for log canonical surfaces in [La]. Since these inequalities are related to stability properties (see [Bo]), it is of interest to investigate the stability properties of the logarithmic cotangent sheaves, with respect to a normal crossing divisor.

In this paper we study these sheaves under suitable hypothesis on the divisor components of D and X . We assume that the Picard number is one in the below results.

We begin with *ample log canonical pairs* (X, D) . More precisely:

Theorem 1.1. *Suppose X is a smooth projective variety of dimension n over \mathbb{C} , with the Picard group $\text{Pic}(X) = \mathbb{Z}$. Let $D = \sum_{i=1}^r D_i$ be a simple normal crossing divisor on X and K_X denote the canonical class. If $K_X + \mathcal{O}_X(D)$ is ample or trivial, then $\Omega_X(\log D)$ is semistable.*

The proof is given in §2.

This statement can be extended to Kawamata coverings, see Proposition 4.2. We note that the Picard number of X in the below proposition can be greater than one, which is relevant for other applications.

Proposition 1.2. *Suppose $(Y, \mathcal{O}_Y(1))$ is a smooth projective variety of dimension n and $D = \sum_i D_i$ is a normal crossing divisor on Y . Assume $\text{Pic}(Y) = \mathbb{Z}$ and $K_Y + D$ is ample or trivial. Consider the Kawamata covering $\pi : X \rightarrow Y$, ramified along D (see Proposition 4.1) and $D' = \pi^{-1}(D)_{\text{red}}$. Then the sheaf $\Omega_X(\log D')$ is semistable with respect to the ample class $\pi^*\mathcal{O}_Y(1)$.*

We next investigate log Fano manifolds (X, D) , in small dimensions. In this situation the class $-K_X - D$ is ample. The classification of such pairs (X, D) is due to Maeda [Md] and Fujita [Fu] in small dimensions. We show:

Theorem 1.3. *Suppose (X, D) is a log Fano manifold of dimension n and $\text{Pic}(X) = \mathbb{Z} \cdot \mathcal{O}_X(1)$. Let the canonical class $K_X = \mathcal{O}_X(-s)$ and D is in the linear system $|\mathcal{O}_X(k)|$, for $s, k > 0$.*

Assume one of the following holds:

a) $n = 2$ and $s = 3$,

b) $n = 3$ and $s \leq 4$

c) $n = 4$ and $s \leq 5$

d) $n = 5$ and $s \leq 6$ such that $s = 2, 5, 6$ or $(s, k) = (3, 2), (4, 3)$.

e) $n = 6$ and $s \leq 7$ such that $s \leq 4$, $s = 6, 7$, or $(s, k) = (5, 4), (5, 3)$.

If D is smooth and irreducible then the logarithmic cotangent bundle $\Omega_X(\log D)$ is semistable.

Since the index s of a Fano variety X is at most $\dim X + 1$, we get complete results, when $\dim X \leq 4$. See Proposition 5.2. The proof involves a careful investigation of vanishing theorems in appropriate range using residue sequences, and apply semistability of de Rham sheaves on Fano manifolds in small dimensions. The classification of Maeda yields complete statements for log Del Pezzo surfaces, log Fano threefolds and fourfolds.

When (X, D) is a log Del Pezzo surface with $D = D_1 + D_2$, D_i are lines, then we check that the log cotangent bundle is not semistable. See Lemma 5.3. This suggests that when D is reducible, we may expect different possibilities.

Acknowledgements: We thank Lawrence Ein for making useful remarks on a previous version.

2. SEMISTABILITY AND LOGARITHMIC VANISHING THEOREMS

We recall the notion of stability ([Hu-Le, p.13]), which we will use in this paper.

2.1. Stability of a coherent sheaf. Suppose X is a smooth projective variety. For any coherent sheaf \mathcal{F} on X , we denote $\det(\mathcal{F}) := c_1(\mathcal{F})$. We denote the slope of \mathcal{F} (with respect to $\mathcal{O}_X(1)$):

$$\mu(E) := \frac{c_1(\mathcal{F})}{rk \mathcal{F}} \cdot \mathcal{O}_X(1)^{n-1}.$$

The sheaf \mathcal{F} is called stable in the sense of Mumford-Takemoto if for any coherent subsheaf \mathcal{G} of \mathcal{F} with $0 < rk \mathcal{G} < rk \mathcal{F}$ we have

$$\mu(\mathcal{G}) < \mu(\mathcal{F}).$$

Similarly, \mathcal{F} is semistable if $\mu(\mathcal{G}) \leq \mu(\mathcal{F})$.

2.2. Slope of logarithmic De Rham sheaves. In this section, we discuss the relation between the vanishing of Dolbeault cohomology and stability of the logarithmic de Rham sheaves. Suppose X is a smooth projective manifold of dimension n and $D \subset X$ is a normal crossing divisor, i.e. $D = \sum_{i=1}^r D_i$ such that D_i intersects D_j transversally, for $i \neq j$. Recall the de Rham sheaves Ω_X and its exterior powers Ω_X^a , for $0 \leq a \leq n$. Consider the logarithmic de Rham sheaves $\Omega_X^a(\log D) := \bigwedge^a \Omega_X(\log D)$, whose local sections are meromorphic a -forms having at most a simple pole along D . Then this is a locally free sheaf of rank n . See [Es-Vi, 2.2,p.11] for more details.

In this paper, we assume that the Picard group of X is \mathbb{Z} and is generated by the ample class $\mathcal{O}_X(1)$. Let $D_i \in |\mathcal{O}_X(k_i)|$, for some positive integers k_i , for $1 \leq i \leq r$.

We first compute the slope of the logarithmic cotangent bundle with respect to $\mathcal{O}_X(1)$, and relate it with vanishing of certain cohomology groups. Let the canonical class $K_X = \mathcal{O}_X(-s)$, for some integer s .

Lemma 2.1. *a) The slope $\mu(\Omega_X^a(\log D))$ is given as*

$$\frac{a \cdot (-s + \sum_{i=1}^r k_i)}{n} \cdot \mathcal{O}_X(1)^n.$$

b) The stability of $\Omega_X(\log D)$ is implied by the vanishing

$$H^0(X, \Omega_X^a(\log D)(-t)) = 0$$

for $-t \leq \frac{a \cdot (s - \sum_{i=1}^r k_i)}{n}$ and $1 \leq a < n$. Similar assertion is true for semistability when we have strictly inequality in the slope inequality.

Proof. a) Consider the short exact sequence of sheaves on X

$$0 \rightarrow \mathcal{O}_X(-D) \rightarrow \mathcal{O}_X \rightarrow \mathcal{O}_D \rightarrow 0.$$

Since the first Chern class c_1 is additive over exact sequences, we have the equality:

$$(1) \quad c_1(\mathcal{O}_D) = -c_1(\mathcal{O}_X(-D)) = c_1(\mathcal{O}_X(D)).$$

Consider the usual residue exact sequences [Es-Vi, Properties 2.3.,p.13]:

$$(2) \quad 0 \rightarrow \Omega_X \rightarrow \Omega_X(\log D) \rightarrow \bigoplus_{i=1}^r \mathcal{O}_{D_i} \rightarrow 0,$$

and

$$(3) \quad 0 \rightarrow \Omega_X^a(\log(D - D_1)) \rightarrow \Omega_X^a(\log D) \rightarrow \Omega_{D_1}^{a-1}(\log(D - D_1)|_{D_1}) \rightarrow 0.$$

Using the additivity of c_1 , we have

$$\begin{aligned} c_1(\Omega_X(\log D)) &= c_1(\Omega_X) + c_1\left(\bigoplus_i \mathcal{O}_{D_i}\right) \\ &= c_1(\Omega_X) + \sum_i c_1(\mathcal{O}(D_i)), \text{ using (1), (2)} \\ &= c_1(\Omega_X) + c_1(\mathcal{O}_X(\sum_i k_i)). \end{aligned}$$

The first Chern class modulo the rank, of the sheaf $\Omega_X^a(\log D)$ is

$$\begin{aligned} \frac{c_1(\Omega_X^a(\log D))}{\binom{n}{a}} &= \frac{\binom{n-1}{a-1} \cdot c_1(\Omega_X(\log D))}{\binom{n}{a}} \\ &= \frac{\binom{n}{a} - \binom{n-1}{a}}{\binom{n}{a}} \cdot c_1(\Omega_X(\log D)) \\ &= \frac{a}{n} c_1(\Omega_X(\log D)). \end{aligned}$$

Hence the slope is given as

$$\begin{aligned} \mu(\Omega_X^a(\log D)) &= \frac{a}{n} c_1(\Omega_X(\log D)) \cdot \mathcal{O}_X(1)^{n-1} \\ &= a \cdot \frac{c_1(\Omega_X) + c_1(\mathcal{O}_X(\sum_i k_i))}{n} \cdot \mathcal{O}_X(1)^{n-1} \\ &= \frac{a \cdot (-s + \sum_{i=1}^r k_i)}{n} \cdot \mathcal{O}_X(1)^n. \end{aligned}$$

b) Suppose there is a subsheaf $\mathcal{F} \subset \Omega_X(\log D)$ of rank $a < n$, destabilizing the sheaf. Then taking determinants, we get a nonzero morphism

$$\det(\mathcal{F}) \rightarrow \Omega_X^a(\log D).$$

Let $\det(\mathcal{F}) = \mathcal{O}_X(t)$, for some integer t . Hence the above morphism gives a nonzero section in $H^0(X, \Omega_X^a(\log D))$. The slope condition says that

$$t > \frac{a \cdot (-s + \sum_i k_i)}{n}$$

Hence semistability is implied by the vanishing

$$H^0(X, \Omega_X^a(\log D)(-t)) = 0, \text{ whenever } -t < \frac{a \cdot (s - \sum_i k_i)}{n}.$$

□

3. STABILITY WHEN THE SHEAF $K_X + \mathcal{O}_X(D)$ IS NON-NEGATIVE.

In this subsection, we proceed to investigate the stability under suitable assumptions on the canonical class with respect to the divisor D . More precisely, we show,

Proposition 3.1. *With notations as in the previous subsection, if $K_X + \mathcal{O}_X(\sum_{i=1}^r k_i)$ is ample or trivial, then the sheaf $\Omega_X(\log D)$ is semistable.*

It suffices to prove vanishing of relevant cohomologies as indicated in Lemma 1 b).

We first prove the following vanishing. This is well-known and due to Norimatsu [No]. For the sake of completeness, we provide a simpler proof:

Lemma 3.2. *Suppose $(Y, \mathcal{O}_Y(1))$ is a smooth projective variety of dimension n . Let $D \subset Y$ be a normal crossing divisor and D is written as $\sum_{i=1}^r D_i$. Then for $t < 0$,*

$$H^0(Y, \Omega_Y^a(\log D)(t)) = 0.$$

Proof. We prove this by using induction on the number of components r of the divisor D .

We start with the case $r = 1$.

Consider the residue sequence

$$0 \rightarrow \Omega_Y^a \rightarrow \Omega_Y^a(\log D) \rightarrow \Omega_D^{a-1} \rightarrow 0.$$

Tensor with $\mathcal{O}(t)$, $t < 0$, and take the long exact cohomology sequence:

$$0 \rightarrow H^0(Y, \Omega_Y^a(t)) \rightarrow H^0(Y, \Omega_Y^a(\log D)(t)) \rightarrow H^0(D, \Omega_D^{a-1}(t)) \rightarrow \dots$$

Since $t < 0$, by Kodaira-Akizuki-Nakano theorem [Es-Vi, 1.3,p.4], the first and the third cohomology groups vanish. This implies the middle cohomology $H^0(Y, \Omega_Y^a(\log D)(t))$ also vanishes.

Now assume that the lemma holds for divisors with at most $r - 1$ components. Consider the residue sequence (3) and tensor with $\mathcal{O}(t)$, for $t < 0$. Now take the associated cohomology sequence

$$0 \rightarrow H^0(Y, \Omega_Y^a(\log(D-D_1))(t)) \rightarrow H^0(Y, \Omega_Y^a(\log D)(t)) \rightarrow H^0(D_1, \Omega_{D_1}^{a-1}(\log(D-D_1)|_{D_1})(t)) \rightarrow \dots$$

By induction hypothesis applied to $D - D_1$ on Y and D_1 , we deduce the vanishing of the middle cohomology as required. \square

The proof of Proposition 3.1 is a corollary of above lemma. Indeed, by Lemma 1 b), it suffices to check the vanishing

$$H^0(X, \Omega_X^a(\log D)(t)) = 0$$

for $t < \frac{a \cdot (s - \sum_{i=1}^r k_i)}{n}$ and $1 \leq a < n$. Recall that $K_X = \mathcal{O}_X(-s)$, s is an integer.

The assumption on $K_X + D$ being ample or trivial implies that $s \leq \sum_{i=1}^r k_i$. Hence the slope condition $t < \frac{a \cdot (s - \sum_{i=1}^r k_i)}{n}$ implies $t < 0$. Now by Lemma 3.2 we conclude the proposition.

4. STABILITY ON KAWAMATA'S FINITE COVERINGS

In this subsection, we recall some details concerning branched finite coverings of a complex projective variety, and investigate stability of the logarithmic de Rham sheaves on the covering variety. Note that the Picard group of such coverings can be bigger than \mathbb{Z} (as pointed out by Ein). Hence it is of interest to look at such cases.

We begin by recalling Kawamata's covering construction:

Proposition 4.1. *Let $(Y, \mathcal{O}_Y(1))$ be a nonsingular projective variety of dimension n . Let $D = \sum_{i=1}^r D_i$ be a simple normal crossing divisor on Y and $D_i \in |\mathcal{O}_Y(k_i)|$, for some positive integers k_i . Then there is a smooth variety X together with a finite flat morphism $\pi : X \rightarrow Y$ such that $\pi^* D_i = k_i \cdot D'_i$, for some divisors D'_i on X such that $D' = \sum_{i=1}^r D'_i$ is a normal crossing divisor on X . Furthermore, the canonical class $K_X = \pi^*(K_Y \otimes \mathcal{O}_Y(\sum_{i=1}^r (k_i - 1)))$.*

Proof. See [La, 4.1.6,4.1.12]. \square

Now we investigate the semistability of the logarithmic de Rham sheaves on the covering variety.

Proposition 4.2. *We keep notations as in Proposition 4.1 for the covering variety $\pi : X \rightarrow Y$. Assume that $\text{Pic}(Y) = \mathbb{Z}\mathcal{O}_Y(1)$ and $k := \sum_{i=1}^r k_i$. If $K_Y + \mathcal{O}_Y(k)$ is ample or trivial then $\Omega_X(\log D')$ is semistable.*

Proof. Since $K_Y + \mathcal{O}(k)$ is ample or trivial, by Proposition 3.1, the sheaf $\Omega_Y(\log D)$ is semistable. By the Generalised Hurwitz formula [Es-Vi, Lemma 3.21, p.33], we have

$$\Omega_X(\log D') \simeq \pi^* \Omega_Y(\log D).$$

Now by [Ma, Lemma 1.17, p. 325], we deduce that the pullback sheaf $\Omega_X(\log D')$ is also semistable, with respect to the ample line bundle $\pi^* \mathcal{O}_Y(1)$. \square

Now in the next section, we investigate the situation when the class $K_X + D$ is anti-ample.

5. LOG FANO MANIFOLDS OF SMALL DIMENSIONS

A pair (X, D) is called a log Fano n -fold if the class $-K_X - D$ is ample. Here $D = \sum_i D_i$ is a normal crossing divisor, D_i are smooth irreducible divisors.

Assume that $\text{Pic}(X) = \mathbb{Z}H$ and the anti canonical class is $-K_X = sH$ and $D \in |kH|$, for some $s, k > 0$. Hence the assumption on ampleness of $-K_X - D$ implies that $s > k$. In particular $s \geq 2$.

In this section we would like to discuss stability for possible cases, when n is small.

5.1. $n = 2$. Here (X, D) is a Del Pezzo surface with $\text{Pic}(X) = \mathbb{Z}H$. By Fujita's classification theorem [Md, p.87], the following cases for (X, D) occur:

- a) (\mathbb{P}^2, H) , where H is a line on \mathbb{P}^2 .
- b) $(\mathbb{P}^2, H_1 + H_2)$, where H_1, H_2 are lines on \mathbb{P}^2 .
- c) (\mathbb{P}^2, Q) , where Q is a conic in \mathbb{P}^2 .

5.2. $n = 3$. Here (X, D) is log Fano threefold with $\text{Pic}(X) = \mathbb{Z}H$. By Maeda's classification [Md, §6, p.95] according to the index s , the following cases occur:

- a) $\mathbf{s} = 4$ and $X = \mathbb{P}^3$. Here D is equivalent to $H, 2H$ or $3H$. Hence we have,
 - 1) (\mathbb{P}^3, D) , where D is a smooth cubic surface.
 - 2) (\mathbb{P}^3, D) , where $D = D_1 + D_2$, and D_1 is a smooth quadric surface and D_2 is a plane.
 - 3) (\mathbb{P}^3, D) , where $D = D_1 + D_2 + D_3$, and each D_i is a plane.
 - 4) (\mathbb{P}^3, D) , where D is a smooth quadric surface.
 - 5) (\mathbb{P}^3, D) , where $D = D_1 + D_2$, and each D_i is a plane.
 - 6) (\mathbb{P}^3, D) , where D is a plane.

b) $\mathbf{s} = 3$ and $X = Q$, a smooth quadric threefold in \mathbb{P}^4 . Here D is equivalent to H or $2H$. Hence we have,

- 1) (Q, D) , where D is a smooth quartic surface in \mathbb{P}^4 .
- 2) (Q, D) , where $D = D_1 + D_2$ and each D_i is a smooth quadric surface.
- 3) (Q, D) , where D is a smooth quadric surface.

c) $\mathbf{s} = \mathbf{2}$. There are five different types of Fano 3-folds and D is a smooth irreducible divisor in the linear system $|H|$.

5.3. $n = 4, 5, 6$. Here the possibilities are more and we refer to [Fu].

We can now state the main result of this section.

We will need the following result in the proof.

Lemma 5.1. *Suppose $(Y, \mathcal{O}(1))$ is a smooth projective variety of dimension n , and D is a smooth irreducible divisor in $|\mathcal{O}(k)|$. Fix $q < n - 1$. Then the restriction map*

$$H^0(Y, \Omega_Y^q(c)) \rightarrow H^0(D, \Omega_D^q(c))$$

is surjective, for all $c < k$.

Proof. See proof of [Pe-Wi, Lemma 2.9 a)]. □

Proposition 5.2. *Suppose (X, D) is a log Fano manifold of dimension n and $\text{Pic}(X) = \mathbb{Z}H$. Let $K_X = \mathcal{O}_X(-s)$ and $D \in \mathcal{O}_X(k)$ such that $s, k > 0$.*

Assume one of the following holds:

- a) $n = 2$ and $s = 3$,
- b) $n = 3$ and $s \leq 4$
- c) $n = 4$ and $s \leq 5$
- d) $n = 5$ and $s \leq 6$ such that $s = 2, 5, 6$ or $(s, k) = (3, 2), (4, 3)$.
- e) $n = 6$ and $s \leq 7$ such that $s \leq 4$, $s = 6, 7$, or $(s, k) = (5, 4), (5, 3)$.

If D is smooth and irreducible then the logarithmic cotangent bundle $\Omega_X(\log D)$ is semistable.

Proof. Suppose D is a smooth and irreducible divisor. Note that the ampleness of $-K_X - D$ implies that $s > k$.

From Lemma 2.1 b), the semistability of $\Omega_X(\log D)$ is implied by the vanishing

$$(4) \quad H^0(X, \Omega_X^a(\log D)(t)) = 0,$$

for $t < \frac{a(s-k)}{n}$ and $1 \leq a < n$.

Recall the residue exact sequence;

$$0 \rightarrow \Omega_X^a(t) \rightarrow \Omega_X^a(\log D)(t) \rightarrow \Omega_D^{a-1}(t) \rightarrow 0.$$

Taking the global sections, we have the long exact sequence

$$0 \rightarrow H^0(X, \Omega_X^a(t)) \rightarrow H^0(X, \Omega_X^a(\log D)(t)) \rightarrow H^0(D, \Omega_D^{a-1}(t)) \rightarrow$$

$$H^1(X, \Omega_X^a(t)) \rightarrow H^1(X, \Omega_X^a(\log D)(t)) \rightarrow \dots$$

Then to prove the vanishing (4), it suffices to check that

$$(5) \quad H^0(X, \Omega_X^a(t)) = 0$$

and the map

$$(6) \quad H^0(D, \Omega_D^{a-1}(t)) \rightarrow H^1(X, \Omega_X^a(t))$$

is injective, whenever $t < \frac{a(s-k)}{n}$ and $1 \leq a < n$.

We now look at the cases listed above, according to the dimension n .

a) $\mathbf{n} = 2$ and $\mathbf{s} = 3$.

By §5.1, the only possibility is $(X, D) = (\mathbb{P}^2, D)$, where D is a line or a conic in \mathbb{P}^2 .

But for $X = \mathbb{P}^2$, $H^0(X, \Omega_X(t)) = 0$ for $t \leq 1$. Hence for $\frac{s-k}{2} = \frac{3-k}{2} \leq 1$, $k = 1, 2$, the vanishing (5) holds. When $t < 0$, then clearly $H^0(D, \mathcal{O}_D(t)) = 0$.

When $t = 0$, then by the hard Lefschetz theorem and the cupping map (for instance see [Pe-Wi, Lemma 1.2]) gives the injectivity of (6).

Hence $\Omega_{\mathbb{P}^2}(\log D)$ is semistable.

b) $\mathbf{n} = 3$ and $\mathbf{s} \leq 4$.

Since X is a Fano 3-fold, by [St, 2.4, p.638], we have the stability of Ω_X . Therefore, by Maruyama's result [Ma], Ω_X^a is semistable. Using the slope inequality in Lemma 2.1, we deduce that

$$(7) \quad H^0(X, \Omega_X^a(t)) = 0, \text{ for } t < \frac{a \cdot s}{3},$$

and when $a = 1$, the vanishing holds for $t \leq \frac{s}{3}$.

On the other hand, $\frac{a(s-k)}{3} < \frac{a \cdot s}{3}$ and this verifies (5).

Now we proceed to check (6) below.

Since $-K_D = \mathcal{O}_D(s-k)$ is ample, D is a Del Pezzo surface. But $\text{Pic}(D)$ can be greater than \mathbb{Z} , hence semistability of Ω_D does not always hold. Hence we argue as follows.

Since $0 < k < s \leq 4$ and $1 \leq a < 3$, the possible values for a are 1, 2 and the possible values for k and s are:

if $k = 1$, then $s = 2, 3, 4$.

if $k = 2$, then $s = 3, 4$.

if $k = 3$, then $s = 4$.

If $t < 0$, then the required vanishing of $H^0(D, \Omega_D^a(t))$, follows from Kodaira-Akizuki-Nakano theorem.

Suppose $a = 1$ and we have $0 \leq t < \frac{(s-k)}{3}$.

If we substitute the respective values of k and s in above range then the only possible value is $t = 0$. The required Hodge vanishing holds because both X and D are Fano manifolds.

Suppose $a = 2$ and we have $0 \leq t < \frac{2 \cdot (s - k)}{3}$. In this case the only possible values are $t = 0, 1$ when $(k, s) = (1, 3)$ and $(2, 4)$. We deduce that it is sufficient to prove, when $t = 1$,

$$H^0(D, \Omega_D(1)) = 0$$

and when $t = 0$,

$$H^0(D, \mathcal{O}_D) \longrightarrow H^1(X, \Omega_X)$$

is injective.

Suppose $t = 1$.

First consider the case when $(k, s) = (2, 4)$. Then by §5.2 a) 4), $(X, D) = (\mathbb{P}^3, D)$, where D is a smooth quadric surface. Using Lemma 5.1, we deduce that the restriction map

$$H^0(\mathbb{P}^3, \Omega_{\mathbb{P}^3}(1)) \rightarrow H^0(D, \Omega_D(1))$$

is surjective.

But we noticed in (7) or it also follows from [Bo], that $H^0(\mathbb{P}^3, \Omega_{\mathbb{P}^3}(1)) = 0$. Hence $H^0(D, \Omega_D(1)) = 0$.

When $(k, s) = (1, 3)$, then by §5.2 b)3), $(X, D) = (Q, H)$, where Q is a smooth quadric threefold and H is a hyperplane section. Hence D is again a quadric surface and $H^0(D, \Omega_D(1)) = 0$.

On the other hand for $t = 0$, by [Pe-Wi, Lemma 2.1], the required injectivity follows.

Hence $\Omega_X(\log D)$ is semi-stable.

c) $\mathbf{n} = 4$ and $\mathbf{s} \leq 5$.

Since X is a Fano 4-fold with $Pic(X) = \mathbb{Z}$, by [Pe-Wi, 2.10,p.15], Ω_X is stable. Therefore, by Maruyama's result the exterior powers are semistable and by Lemma 2.1, we have

$$(8) \quad H^0(X, \Omega_X^a(t)) = 0, \text{ for } t < \frac{a \cdot s}{4}.$$

On the other hand, $\frac{a \cdot (s - k)}{4} < \frac{a \cdot s}{4}$ and we have $H^0(X, \Omega_X^a(t)) = 0$ for $t < \frac{a \cdot (s - k)}{4}$.

Since $-K_D = \mathcal{O}_D(s - k)$ is ample, D is a Fano 3-fold with $Pic(D) = \mathbb{Z} \cdot H|_D$ (by Lefschetz hyperplane section theorem). Hence Ω_D is stable [St, 2.4,p.638].

Therefore, again by Maruyama's result we have the semistability of its exterior powers and by Lemma 2.1,

$$H^0(D, \Omega_D^{a-1}(t)) = 0, \text{ for } t < \frac{(a-1) \cdot (s-k)}{3}.$$

Since $\frac{(a-1).(s-k)}{3} < \frac{a.(s-k)}{4}$, we only have to discuss the situation

$$\frac{(a-1).(s-k)}{3} \leq t < \frac{a.(s-k)}{4}.$$

Since $0 < k < s \leq 3$ and $1 \leq a < 4$, the possible values for a are 1, 2, 3 and the possible values for k and s are:

if $k = 1$, then $s = 2, 3$,

if $k = 2$, then $s = 3$.

If we substitute the respective values of k , s and a in $\frac{(a-1).(s-k)}{3} \leq t < \frac{a.(s-k)}{4}$ then the only possible value which remains is $t = 0$ and when $a = 1$.

Therefore, as before injectivity of (6) follows from [Pe-Wi, Lemma 1.2].

Suppose $\mathbf{s} = \mathbf{4}$, then we note that we need vanishing of only $H^0(D, \Omega_D^{a-1}(t))$, if $a = 2$ and $k = 1$. In this case X is a smooth quadric 4-fold and D is a smooth quadric threefold in \mathbb{P}^3 . Hence we can apply Lemma 5.1, to get the desired vanishing.

Suppose $\mathbf{s} = \mathbf{5}$, then $X = \mathbb{P}^4$. We note that we need to check vanishing of $H^0(D, \Omega_D^{a-1}(t))$ only when $a = 2, k = 2, t = 1$ and when $a = 3, k = 2, t = 2$. In this case, D is a smooth quadric threefold. Both these vanishings follow from [Sn, Theorem (1), p.174].

Hence $\Omega_X(\log D)$ is semi-stable.

d) $\mathbf{n} = \mathbf{5}$ and $\mathbf{s} \leq \mathbf{6}$.

Since X is a Fano 5-fold, Ω_X is stable [Hw, Theorem 2, p.605]. Therefore, by Maruyama's result,

$$H^0(X, \Omega_X^a(t)) = 0, \text{ for } t < \frac{a.s}{5}.$$

Note that D is a Fano fourfold with $Pic(D) = \mathbb{Z}.H|_D$.

If $t \leq 0$ and except when $a = 1, t = 0$, then by Kodaira-Nakano vanishing theorem and by rational connectedness of D ,

$$H^0(D, \Omega_D^{a-1}(t)) = 0.$$

So when $t \leq 0$, we have the desired vanishing $H^0(X, \Omega_X^a(\log D)(t)) = 0$.

Suppose $t = 0$ and $a = 1$ then by [Pe-Wi, Lemma 2.1] the injectivity of (6) follows.

As in the previous case, we need to look at the case:

$$\frac{(a-1)(s-k)}{4} \leq t < \frac{a.(s-k)}{5}$$

to obtain vanishing of $H^0(D, \Omega_D^{a-1}(t))$.

We note that we need to check the following cases only:

1) $(s = 3, k = 1, a = 3, t = 1)$,

2) $(s = 4, k = 1, a = 2, t = 1)$,

- 3) $(s = 4, k = 2, a = 3, t = 1)$
- 4) $(s = 5, k = 1, a = 2, t = 1)$
- 5) $(s = 5, k = 1, a = 3, t = 2)$
- 6) $(s = 5, k = 1, a = 4, t = 3)$
- 7) $(s = 5, k = 2, a = 2, t = 1)$
- 8) $(s = 5, k = 3, a = 3, t = 1)$
- 9) $(s = 6, k = 2, a, t = a - 1)$
- 10) $(s = 6, k = 3, a = 2, t = 1)$
- 11) $(s = 6, k = 4, a = 3, t = 1)$.

We check that in 4),5),6), D is a smooth quadric hypersurface in \mathbb{P}^5 . Hence the vanishing $H^0(D, \Omega_D^{a-1}(a-1)) = 0$ holds by [Sn, Theorem 1],p. 174]. Similarly, 9) also hold because D is a smooth quadric hypersurface in \mathbb{P}^6 . We again use Snow's theorem and apply Lemma 5.1 to get the required vanishing on D , in case of 8), 10), and 11).

The remaining cases are not known to us.

Hence $\Omega_X(\log D)$ is semi-stable in the cases claimed.

d) $\mathbf{n} = \mathbf{6}$ and $\mathbf{s} \leq \mathbf{7}$.

Since X is a Fano 6-fold, by [Hw, Theorem 3,p.605], Ω_X is semi-stable. Therefore, by Maruyama's result

$$H^0(X, \Omega_X^a(t)) = 0 \text{ for } t < \frac{a \cdot s}{6}.$$

On the other hand, $\frac{a \cdot (s - k)}{6} < \frac{a \cdot s}{6}$ we have

$$H^0(X, \Omega_X^a(t)) = 0, \text{ for } t < \frac{a \cdot (s - k)}{6}.$$

Since $-K_D = \mathcal{O}_D(s - k)$ is ample, D is a Fano 5-fold with $Pic(D) = \mathbb{Z} \cdot H|_D$ and hence Ω_D is stable [Hw, Theorem 2,p.605]. Therefore, again by Maruyama's result we have

$$H^0(D, \Omega_D^{a-1}(t)) = 0, \text{ for } t < \frac{(a-1) \cdot (s-k)}{5}.$$

Since $\frac{(a-1) \cdot (s-k)}{5} < \frac{a \cdot (s-k)}{6}$, so we have only to discuss the situation

$$\frac{(a-1) \cdot (s-k)}{5} \leq t < \frac{a \cdot (s-k)}{6}.$$

Since $0 < k < s \leq 4$ and $1 \leq a < 6$, the possible values for a are 1, 2, 3, 4, 5 and the possible values for k and s are:

if $k = 1$, then $s = 2, 3, 4$,

if $k = 2$, then $s = 3, 4$,

if $k = 3$, then $s = 4$.

Suppose $a = 1$ we have $0 \leq t < \frac{(s-k)}{6}$.

If we substitute the respective values of k and s in above range then the only possible value is $t = 0$. In this case it is enough to show $H^0(D, \mathcal{O}_D) \rightarrow H^1(X, \Omega_X^1)$ is injective. But this follows from the cupping map [Pe-Wi, Lemma 2.1].

Suppose $a = 2$ then we have $\frac{(s-k)}{5} \leq t < \frac{2 \cdot (s-k)}{6}$. In this case t does not exist.

Suppose $a = 3, 4$, and if we substitute the respective values of k , s and a in

$$\frac{(a-1) \cdot (s-k)}{5} \leq t < \frac{a \cdot (s-k)}{6}.$$

Then the possible value is $t = 1$ when

- $a = 3$ and $(k, s) = (1, 4)$,
- $a = 4$ and $(k, s) = (1, 3), (2, 4)$.

In this case it is enough to show

$$H^0(D, \Omega_D^2(1)) = 0 \text{ when } (k, s) = (1, 4)$$

and

$$H^0(D, \Omega_D^3(1)) = 0 \text{ when } (k, s) = (1, 3), (2, 4).$$

But both the claims follows from stability of Ω_D .

Suppose $a = 5$ then we have $\frac{4 \cdot (s-k)}{5} \leq t < \frac{5 \cdot (s-k)}{6}$. If we substitute the respective values of k and s in above range then the only possible value is $t = 2$ when $(k, s) = (1, 4)$.

In this case it is enough to show

$$H^0(D, \Omega_D^4(2)) = 0 \text{ when } (k, s) = (1, 4).$$

But this follows from stability of Ω_D .

The following cases need only to be discussed:

s = 6:

- 1) $(s = 6, k = 1, a, t = a - 1)$
- 2) $(s = 6, k = 2, a = 2, t = 1)$
- 3) $(s = 6, k = 3, a = 2, t = 1)$
- 4) $(s = 6, k = 4, a = 3, t = 1)$.

s = 7:

- 5) $(s = 7, k = 2, a, t = a - 1)$
- 6) $(s = 7, k = 3, a = 2, t = 1)$.

In 1) and 5), we note that D is a smooth quadric hypersurface in \mathbb{P}^6 and by [Sn, Theorem 1,p.174], the vanishing $H^0(D, \Omega_D^{a-1}(a-1)) = 0$ holds. Again by Snow's theorem, and applying Lemma 5.1 we deduce the required vanishing in case 1)-4), 6).

Hence $\Omega_X(\log D)$ is semi-stable in the cases claimed. □

5.4. Counterexample when D is reducible. We now investigate the situation when D is reducible and when (X, D) is Del Pezzo surface.

Lemma 5.3. *Suppose $(X, D) = (\mathbb{P}^2, D_1 + D_2)$, where D_1, D_2 are lines on \mathbb{P}^2 . Then $\Omega_{\mathbb{P}^2}(\log D)$ is not semistable.*

Proof. The semistability of $\Omega_{\mathbb{P}^2}(\log D)$ is equivalent to the vanishing (see (4)):

$$H^0(\mathbb{P}^2, \Omega_{\mathbb{P}^2}(\log D)(t)) = 0$$

for $t < \frac{(3-2)}{2}$, i.e. when $t \leq 0$.

When $t < 0$, this follows from Lemma 3.2.

When $t = 0$, we note that the injectivity of map

$$\bigoplus_{i=1,2} H^0(D_i, \mathcal{O}_{D_i}) \rightarrow H^1(\mathbb{P}^2, \Omega_{\mathbb{P}^2})$$

(see (2)) fails. Indeed, here $\bigoplus_{i=1,2} H^0(D_i, \mathcal{O}_{D_i})$ is of rank two and $H^1(\mathbb{P}^2, \Omega_{\mathbb{P}^2})$ is of rank one. □

Remark 5.4. *We suspect that in higher dimensional cases with several divisor components, the semistability may fail.*

REFERENCES

- [Ak-Na] Y. Akizuki, S. Nakano : *Note on Kodaira-Spencer's proof of Lefschetz's theorem.* Proc. Jap. Acad., Ser A **30** (1954), 266 - 272.
- [Au] T. Aubin : *Nonlinear analysis on manifolds.* Berlin Heidelberg New York: Springer 1982.
- [Bo] F. Bogomolov : *Unstable vector bundles and curves on surfaces.* Proceedings of the International Congress of Mathematicians (Helsinki, 1978), pp. 517524, Acad. Sci. Fennica, Helsinki, 1980.
- [Bo] R. Bott : *Homogeneous vector bundles* Ann. of Math.,**66**,203-248, 1957.
- [De] P. Deligne : *Théorie de Hodge. II.* (French) Inst. Hautes Études Sci. Publ. Math. No. **40** (1971), 557.
- [Do] S. K. Donaldson : *Infinite determinants, stable bundles and curvature.* Duke Math. J. 54, (1987), no. **1**, 231-247.
- [Ha] R. Hartshorne : *Algebraic Geometry*, Grad. Texts in Math. **52**, Springer-Verlag, New York, 1977.
- [Es-Vi] H. Esnault, E. Viehweg : *Lectures on vanishing theorems.* DMV Seminar, **20**. Birkhäuser Verlag, Basel, 1992. vi+164 pp.
- [Fu] K. Fujita : *Simple normal crossing Fano varieties and log Fano manifolds.*, arXiv.math 1206.1994.
- [Hu-Le] D. Huybrechts, M. Lehn : *The geometry of moduli spaces of sheaves.* Second edition. Cambridge Mathematical Library. Cambridge University Press, Cambridge, 2010. xviii+325 pp.

- [Hw] J-M. Hwang : *Stability of tangent bundles of low-dimensional Fano manifolds with Picard number 1*. Math. Ann. 312 (1998), no. 4, 599-606.
- [La] A. Langer : *The Bogomolov-Miyaoka-Yau inequality for log canonical surfaces*. J. London Math. Soc. (2) 64 (2001), no. 2, 327-343.
- [Li-Su] Chi Li, Song Sun : *Conical Kähler-Einstein metric revisited*, arXiv.math 1207.5011.
- [Lu] M. Lübke : *Stability of Einstein-Hermitian vector bundles*. Manuscr. Math. **42**, 245-257 (1983).
- [Kb] S. Kobayashi : *Differential geometry of holomorphic vector bundles*. Berkeley lecture notes, 1986.
- [Ko] J. Kollar : *Rational curves on algebraic varieties*. Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge., **32**. Springer-Verlag, Berlin, 1996. viii+320.
- [La] R. Lazarsfeld : *Positivity in Algebraic Geometry. 1*, Results in Mathematics and Related Areas. 3rd Series. A Series of A Modern Surveys in Mathematics, **48**. Springer-Verlag, Berlin, 2004.
- [Ma] M. Maruyama : *The theorem of Grauert-Mulich-Spindler*, Math. Ann. 255, (1981), no. 3, 317-333.
- [Md] H. Maeda : *Classification of logarithmic Fano threefolds*. Compositio Math. 57 (1986), no. 1, 81-125.
- [No] Y. Norimatsu : *Kodaira vanishing theorem and Chern classes for δ -manifolds*, Proc. Japan Acad Ser. A Math Sci, 54, (1978), no. 4, 107-108.
- [Pe-Wi] T. Peternell, J. A. Wisniewski : *On stability of tangent bundles of Fano manifolds with $b_2 = 1$* J. Algebraic Geom. 4 (1995), no. 2, 363-384.
- [Sn] D. Snow : *Cohomology of twisted holomorphic forms on Grassmann manifolds and Quadric hypersurfaces*, Math. Annalen, **276**, 159-176, 1986.
- [St] A. Steffens : *On the stability of the tangent bundle of Fano manifolds*. Math. Ann. 304 (1996), no. 4, 635-643.
- [Su] S. Subramanian : *Stability of the tangent bundle and existence of a Kähler-Einstein metric*. Math. Ann. 291 (1991), no. 4, 573-577.
- [Si] Y-T. Siu : *Lectures on Hermitian-Einstein metrics and Kähler-Einstein metrics*. Basel Boston: Birkhauser 1987.
- [Ti] G. Tian : *On Kähler-Einstein metrics on certain Kähler manifolds with $c_1(M) > 0$* . Invent. Math. **89**, 225-246 (1987).
- [Ti2] G. Tian : *On stability of the tangent bundles of Fano varieties*, International Journal of math. **3**, 401-413 (1992).
- [Ti3] G. Tian : *Kähler-Einstein metrics on algebraic manifolds*. Transcendental methods in algebraic geometry (Centraro, 1994), 143-185, Lecture Notes in Math., **1646**, Springer, Berlin, 1996.
- [Ts] H. Tsuji : *Stability of tangent bundles of minimal algebraic varieties*. Topology **27**, 429-442, (1988).
- [Ya] S-T. Yau : *On the Ricci curvature of a compact Kähler manifold and the complex Monge-Ampère equation. I*. Comm. Pure Appl. Math. 31 (1978), no. 3, 339-411.

THE INSTITUTE OF MATHEMATICAL SCIENCES, CIT CAMPUS, TARAMANI, CHENNAI 600113, INDIA

E-mail address: seshadrich@imsc.res.in

E-mail address: jniyer@imsc.res.in