



## Mitochondrial DNA Part B Resources

ISSN: (Print) 2380-2359 (Online) Journal homepage: <http://www.tandfonline.com/loi/tmdn20>

# Complete mitochondrial genome sequence of Asiatic lion (*Panthera leo persica*)

Wajeeda Tabasum, Sreenivas Ara, Niraj Rai, Kumarasamy Thangaraj & Ajay Gaur

To cite this article: Wajeeda Tabasum, Sreenivas Ara, Niraj Rai, Kumarasamy Thangaraj & Ajay Gaur (2016) Complete mitochondrial genome sequence of Asiatic lion (*Panthera leo persica*), *Mitochondrial DNA Part B*, 1:1, 619-620, DOI: [10.1080/23802359.2016.1214541](https://doi.org/10.1080/23802359.2016.1214541)

To link to this article: <http://dx.doi.org/10.1080/23802359.2016.1214541>



© 2016 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group.



Published online: 05 Sep 2016.



Submit your article to this journal [↗](#)



Article views: 127



View related articles [↗](#)



View Crossmark data [↗](#)

Full Terms & Conditions of access and use can be found at  
<http://www.tandfonline.com/action/journalInformation?journalCode=tmdn20>

## Complete mitochondrial genome sequence of Asiatic lion (*Panthera leo persica*)

Wajeeda Tabasum<sup>a</sup>, Sreenivas Ara<sup>a</sup>, Niraj Rai<sup>b</sup>, Kumarasamy Thangaraj<sup>b</sup> and Ajay Gaur<sup>a</sup>

<sup>a</sup>Conservation Genetics Lab, LaCONES, CSIR-Centre for Cellular and Molecular Biology, Hyderabad, India; <sup>b</sup>Evolutionary Biology and Medical Genetics Lab, CSIR-Centre for Cellular and Molecular Biology, Hyderabad, India

### ABSTRACT

The complete mitochondrial genome sequence 17,059 bp of Asiatic lion (*Panthera leo persica*) has been sequenced with the use of next generation sequencing technology using Ion Torrent<sup>PGM</sup> platform. The complete mitochondrial genome sequence of Asiatic lion consists of 13 protein-coding, 22 tRNA, and two rRNA genes, and 1 control region (CR). The mitochondrial genome is relatively similar to other felid mitochondrial genomes with respect to gene arrangement, composition, tRNA structures and skews of AT/GC bases to be typical of those reported for other mammals. The nucleotide composition of Asiatic lion mitogenome shows that there is more A-T% than G-C% on the positive strand as revealed by positive AT and CG skews. The overall base composition is 31.9% of A, 27.2% of C, 14.5% of G, and 26.2% of T. Most of the genes have ATA start codons, except ND1, COX2, ATP8, ATP6, ND4, and ND5 have ATG start codons.

### ARTICLE HISTORY

Received 15 December 2015  
Revised 15 July 2016  
Accepted 15 July 2016

### KEYWORDS

Asiatic lion; complete mitochondrial genome; ngs; conservation

The Asiatic lion (*Panthera leo persica*) is classified as a member of Felidae family consisting of two major sub families Pantherinae and Felinae. Pantherinae includes six big cats i.e. Tiger, Lion, Jaguar, Leopard, Snow leopard, and Clouded leopard (Johnson et al. 1996, 2006). Asiatic lions are the top carnivores and possess a prominent position atop the food chain in their only remaining natural habitat in Gir Forest of Gujarat in India, thus the reduction in their population size would lead to ecological imbalance (Donald & Kenneth 2010).

The blood sample used for DNA extraction and analysis was sampled from a male Asiatic lion named Siddharath with a studbook No. 370 from Nehru Zoological Park, Hyderabad, India (N: 17°21'04", E: 78°26'59").


In this study, the complete mitochondrial genome of Asiatic lion was sequenced and characterized. Two set of primers were designed based on highly conserved sequences of an alignment with full-length mitochondrial genomes from the available public database (unpublished) and used for PCR, with an average amplicon size of 8500 bp each with an overlap 614 bp were used for sequencing. The overall characteristics of mitogenome of Asiatic lion were identical to the typical vertebrate mitochondrial genome. The complete mitochondrial genome of Asiatic lion is 17,059 bp in length (GeneBank accession number KU234271), made up of 37 genes, which includes 13 protein-coding genes, 2 rRNA genes, and 22 tRNA genes, as well as a control region, is highly conserved among most vertebrates mitochondrial genome (Boore 1999).

Gene order and origin of reading frame of all protein-coding genes were identical to other members of Carnivora

except for ND6 and nine tRNA genes. The total length of the 13 protein-coding genes was 11,364 bp, which corresponds to 66.62% of the mitochondrial genome sequence length. The longest gene was ND5 (1805 bp) and the shortest gene was ATP8 (195 bp). The start and stop codons appeared universal among species, except for ND2, COX1, COX3, ND3, ND4L, ND6, and CYTB (ATA start codon), the remaining protein-coding genes start with ATG. The length of the 22 tRNA genes ranges from 58 to 74 bp in Asiatic lion. In Asiatic lion, origin of L-strand replication (OL) was within a cluster of five tRNA genes i.e. tRNA<sup>Trp</sup>, tRNA<sup>Ala</sup>, tRNA<sup>Asn</sup>, tRNA<sup>Cys</sup>, and tRNA<sup>Tyr</sup> and could also be folded into a stable stem-loop secondary structure. The noncoding region i.e. CR, was located between the tRNA-Pro and tRNA-Phe genes and is 1624 bps long.

The phylogenetic position of Asiatic lion was estimated using maximum-likelihood (ML) and the Bayesian inference (BI) (Figure 1) containing concatenated 13 protein-coding genes of 14 species derived from different families in sub-order feliformia. The Bayesian analysis was performed using MRBAYES v 3.1.2 (Huelsenbeck et al. 2001) with four chains of  $1.1 \times 10^5$  generations and sampling the trees every 100 generations. ML tree was similar to the BI tree. The monophyly of genus *panthera* was clearly depicted and were statistically supported by high bootstrap values.

The characterization of Asiatic lion mitogenome will contribute to more refined phylogeny of big cats. Our study will also facilitate concentrated efforts towards the conservation and proper management of the only remaining population of Asiatic lions in India.



**Figure 1.** The Bayesian majority-rule consensus tree is inferred from the combined data set of 13 mitochondrial protein-coding genes. The node value represents the Bayesian posterior probabilities. The phylogenetic tree was rooted using *Cuon alpinus* (NC\_013445). Our sample sequence was *Panthera leo persica* (KU\_234271). The analysed species and the corresponding NCBI accession numbers are as follows: *Panthera leo leo* (KF776494), *Panthera pardus* (EF\_551002), *Uncia uncia* (NC\_010638), *Panthera onca* (KM\_236783), *Panthera tigris* (EF\_551003), *Neofelis nebulosa* (NC\_008450), *Acinonyx jubatus* (NC\_005212), *Prionailurus bengalensis* (KP\_246843), *Nandinia binotata* (NC\_024567), *Viverricula indica* (NC\_025296), *Genetta servalina* (NC\_024568), *Paguma larvata* (NC\_029403) and *Prionodon pardicolor* (NC\_024569).

### Nucleotide sequence accession number

The complete genome sequence of Asiatic lion (*Panthera leo persica*) has been assigned and Genbank accession number KU234271.

### Acknowledgements

The authors would like to thank Director, CCMB and CSIR for all the support.

### Disclosure statement

The authors report no conflicts of interest. The authors alone are responsible for the content and writing of the paper.

### References

- Boore JL. 1999. Animal mitochondrial genomes. *Nucleic Acids Res.* 27:1767–1780.
- Donald RS, Kennen TF. 2010. Human involvement in food webs. *Annu Rev Environ Resour.* 35:1–23.
- Huelsenbeck JP, Ronquist F, Nielsen R, Bollback JP. 2001. Bayesian inference of phylogeny and its impact on evolutionary biology. *Science.* 294:2310–2314.
- Johnson WE, Dratch PA, Martenson JS. 1996. Resolution of recent radiations within three evolutionary lineages of Felidae using mitochondrial restriction fragment length polymorphism variation. *J Mamm Evol.* 3:97–120.
- Johnson WE, Eizirik E, Pecon-Slatery J, Murphy WJ, Antunes A, Teeling E, O'Brien SJ. 2006. The late Miocene radiation of modern Felidae: a genetic assessment. *Science.* 311:73–77.