

Mycobacterium tuberculosis directs T helper 2 cell differentiation by inducing interleukin-1 β production in dendritic cells

Ved Prakash Dwivedi¹, Debapriya Bhattacharya¹, Samit Chatterjee¹, Debprasad Chattopadhyay², Luc Van Kaer³, William R. Bishai⁴ and Gobardhan Das^{1,*}

¹Immunology Group, International Centre for Genetic Engineering and Biotechnology, Aruna Asaf Ali Marg, New Delhi 110067, India.

²ICMR Virus Unit, Calcutta, ID & BG Hospital, GB 4, Beliaghata, Kolkata, India

³Department of Pathology, Microbiology and Immunology, Vanderbilt University School of Medicine, Nashville, TN 37232-2363.

⁴Kwazulu-Natal Research Institute for TB and HIV, Durban, 4001, South Africa.

***Address for correspondence:**

Gobardhan Das, Ph.D.
Staff Research Scientist,
International Centre for Genetic Engineering and Biotechnology,
New Delhi, India
E-mail: dasgo@icgeb.res.in

Or
Prof. William R Bishai, Director
Kwazulu-Natal Research Institute for TB and HIV
Suite Z310 Doris Duke Building
719 Umbilo Road
Durban, 4001, South Africa

Running title: IL-1 β directs Th2 cell differentiation in *M. tb* infection.

Background: *Mycobacterium tuberculosis* (*M. tb*) survives within the host by modulating host immune responses.

Results: RD-1/ESAT-6 from *M. tb* induces IL-1 β in dendritic cells to direct Th2-differentiation, which facilitates disease progression by inhibiting host protective Th1-responses.

Conclusion: Cytokines produced by *M. tb*-infected cells play a role in promoting Th2-responses to subvert host immunity.

Significance: These findings contribute to understanding the immune-evasion mechanisms of *M. tb*.

Abstract:

Mycobacterium tuberculosis (*M. tb*), the causative agent of tuberculosis (TB), resides and replicates within phagocytes and persists in susceptible hosts by modulating protective innate immune responses. Furthermore, *M. tb* promotes T helper 2 (Th2) immune responses by altering the balance of T cell polarising cytokines in infected cells. However, cytokines that regulate Th2 cell differentiation during TB infection remain unknown. Here we show that IL-1 β , produced by phagocytes infected by virulent *M. tb* strain H37Rv, directs Th2 cell differentiation. In sharp contrast, the vaccine strain BCG as well as RD-1 and ESAT-6 mutants of H37Rv failed to induce IL-1 β and promote Th2 cell differentiation. Furthermore, ESAT-6 induced IL-1 β production in dendritic cells (DCs), and CD4⁺ T cells co-cultured with infected DCs differentiated into Th2 cells. Taken together, our findings indicate that IL-1 β induced by RD-1/ESAT-6 plays an important role in the differentiation of Th2 cells, which in turn facilitates progression of TB by inhibiting host protective Th1 responses.

Introduction:

Tuberculosis (TB) is a global health problem that causes over nine million new cases and claims two million lives each year (1). According to the world health organization (WHO), nearly 2 billion people or nearly one third of the global population is latently infected with *Mycobacterium tuberculosis* (*M. tb*), the etiologica agent of TB (2). Current therapy for TB is lengthy and consists of multiple antibiotics, which have the risk of generating

drug-resistant variants of *M. tb* (3-5). In fact, almost all countries, irrespective of their socio-economic status, are now under threat from multiple drug resistant (MDR) and extensively drug resistant (XDR) stains of *M. tb* (3, 5, 6). Unfortunately, the rate by which new drug-resistant variants of this deadly organism are generated is much faster than the rate by which new generation antibiotics are being discovered, which has already resulted in the appearance of totally drug resistant (TDR) TB strains (7, 8). Therefore, alternative treatment methods that avoid the generation of MDR strains are urgently needed. Immunotherapy is one possible option that might avoid such risks.

It has been well-established that T helper 1 (Th1) cells producing interferon (IFN)- γ play a central role in host resistance to *M. tb* infection (9, 10). Consequently, interleukin (IL)-12-, IL-12 receptor (IL-12R)-, signal transducer of activation and transcription (STAT)-4-, and T-bet-deficient animals, which are defective in the generation of T helper 1 (Th1) responses, are highly susceptible to *M. tb* infection (9, 11). Similarly, individuals with mutations in IL-12 or its signalling components are highly susceptible to *M. tb* infection (12, 13). It has been well-documented that *M. tb* inhibits Th1 cells by inhibiting IL-12 production in infected cells (14). On the other hand, transfroming growth factor (TGF)- β produced by the infected cells directs the differentiation of induced regulatory T cells (iTregs) (15-17). In fact, it has been shown that iTregs are induced during the progression of TB, and this is associated with inhibition of protective T cell responses in the host (18, 19). *M. tb* infected cells also produce IL-6, which, in combination with TGF- β , provides favourable conditions for the differentiation of Th17 cells (20, 21). All Th cell subsets, including Th1 cells, Th2 cells, Th17 cells, and Tregs are in a dynamic balance, and it has been therefore assumed that impaired Th1 responses during the progression of TB are associated with increased Th2 responses. However, the cytokines produced by the infected cells that play a role in the development of Th2 responses during progression of TB have not been identified.

It has previously been shown that *M. tb* infected macrophages produce copious amounts of IL-1 β (22, 23). IL-1 β plays an important role in the differentiation of Th2 cells (24). Thus, we examined whether IL-1 β produced by infected

phagocytes is responsible for Th2 cell differentiation during progression of TB. We observed that dendritic cells (DCs) infected with the virulent *M. tb* strain H37Rv induced copious amounts of IL-1 β , whereas the vaccine strain Baccille Calmette Guérin (BCG) and *M. tb* strains carrying mutations in the region of difference (RD)-1 (H37Rv Δ RD-1) or early secretory antigenic target (ESAT)-6 (H37Rv Δ ESAT-6) failed to induce IL-1 β . Importantly, complementation of H37Rv Δ ESAT-6 with ESAT-6 restored IL-1 β production in DCs. On the other hand, H37Rv, BCG and all H37Rv mutants induced comparable amounts of IL-12. H37Rv infected DCs induced both IFN- γ - and IL-4-producing CD4⁺ T cells, whereas BCG, H37Rv Δ RD-1, and H37Rv Δ ESAT-6 selectively induced IFN- γ -producing cells. We confirmed these findings by adoptive transfer experiments with naive CD4⁺ T cells isolated from IL-1 β R knockout mice. Taken together, our findings indicate that IL-1 β produced by infected phagocytes plays an important role in Th2 cell differentiation during TB.

Experimental Procedures:

Ethics Statement- Animal experiments were performed according to the guidelines approved by the Institutional Animal Ethics Committee meeting held on 22nd November 2007 at ICGEB (approval ID; ICGEB/IAEC/IMM-13/2007), New Delhi, India and Department of Biotechnology (DBT) guidelines, Government of India. All mice used for experiments were ethically sacrificed by asphyxiation in carbon dioxide according to institutional and DBT regulations.

Mice- C57BL/6 and OT-II T cell receptor (TCR) transgenic mice (6–8 wks of age) were initially purchased from The Jackson Laboratory, USA. IL-1 β R^{-/-} mice were purchased from the Jackson Laboratory, USA. All animals were subsequently bred and maintained in the animal facility of the International Centre for Genetic Engineering and Biotechnology (ICGEB), New Delhi, India.

Bacteria- *Mycobacterium tuberculosis* strain H37Rv was a kind gift from the Colorado State University repository. BCG, H37Rv Δ RD-1, H37Rv Δ ESAT-6 and ESAT-6-complemented H37Rv Δ ESAT-6 were a kind gift from Prof. David Sherman (SBRI, Seattle, USA). All

mycobacterial strains were grown in 7H9 (Middlebrook, Difco™, USA) medium supplemented with 10% ADC (albumin, dextrose, and catalase; Difco™, USA) and with 0.05% Tween 80 and 0.2% glycerol, and cultures were grown to mid-log phase. Aliquots of the cultures in 20% glycerol were preserved at -80°C and these cryo-preserved stocks were used for infections.

Reagents- Luminex kits were purchased from Bio-Rad. GM-CSF and IL-4 were obtained from R&D Biosystems, USA. Purified anti-mouse IL-1 β antibody (clone B122) was purchased from e-Biosciences, USA.

Generation of dendritic cells (DCs) - Mice were euthanized and the femurs were isolated. Bone marrow (BM) was flushed out with RPMI-1640 medium using a 2.0 ml syringe (26.5G). The cells were washed twice with PBS and then cultured in complete RPMI-1640 (Gibco, UK) medium supplemented with granulocyte monocyte-colony stimulating factor (GM-CSF, 40 ng/ml) and IL-4 (10 ng/ml) on 24-well plates (1 million cells/ml). On the third day, 75% of the medium was replaced with fresh DC culture medium. On day 5, the suspended cells were removed and the loosely adherent cells were collected as immature DCs (CD11c-positive cells were >90%). For mature DCs, immature DCs were stimulated with lipopolysaccharide (LPS, 1 μ g/ml) for 24 hours. FACS analysis using anti-CD11c, -CD80, -CD86, and -MHC class II antibodies suggested that >95% of the cells were conventional DCs.

Bacterial infections and co-culture of DCs with CD4⁺ T cells- *In vitro:* BM cells were isolated from mice (C57BL/6), differentiated into immature DCs as described above and cultured in 24-well plates (1x10⁶ cells per well). Cells were infected with H37Rv, H37Rv Δ RD-1, BCG, H37Rv Δ ESAT-6, or ESAT-6-complemented H37Rv Δ ESAT-6 (MOI of 1:10). Supernatants from cells were collected at 24, 48 and 72 hrs for cytokine profiling. For T cell differentiation, CD4⁺ T cells (1x10⁶) were purified by magnetic activated cell sorting (MACS) method (CD4⁺ T cell isolation beads kit; Miltenyi Biotech, Germany) from OT-II TCR transgenic mice and co-cultured with immature DCs (1x10⁶) infected with H37Rv, H37Rv Δ RD-1, BCG, H37Rv Δ ESAT-6, or ESAT-6-complemented H37Rv Δ ESAT-6 in the presence of ovalbumin (OVA) (10 μ g/ml) peptide (Thermo Scientific, USA) for 72 hrs.

Then, CD4⁺ T cells were harvested and subjected to intracellular staining for IL-4 and IFN- γ expression.

In vivo: Mice (C57BL/6) used for these experiments were infected with H37Rv, H37Rv Δ RD-1, H37Rv Δ ESAT-6, BCG or ESAT-6-complemented H37Rv Δ ESAT-6 through the aerosol route.

Detection of cytokines- Cytokines in the culture supernatant of DCs were assayed by a Luminex microbead-based multiplexed assay using commercially available kits according to the manufacturer's protocol (BioPlex, Bio-Rad, USA).

Antibody treatment- For depletion of IL-1 β , we cultured cells in the presence of anti-IL-1 β mAb (10 μ g/ml) and collected the supernatant for cytokine profiling after different time points.

T cell adoptive transfer- For adoptive transfer experiments, Thy1.1⁺ mice were gamma-irradiated (8 rads/sec for 100 seconds) and rested for a day. CD4⁺ T cells, isolated from lymph nodes of IL-1R^{-/-} mice (Thy1.2⁺ background) were then adoptively transferred into the irradiated recipient mice (2-4 X 10⁶ cells per mouse). After 15 days recipient mice were challenged with H37Rv through the aerosol route.

Intracellular cytokine staining- For intracellular cytokine staining, cells were treated with 50 ng/ml phorbol myristate acetate (PMA) and 500 ng/ml ionomycin in the presence of 10 μ g/ml brefeldin A (Sigma-Aldrich or eBiosciences, USA) added during the last 6 h of culture. Cells were washed twice with PBS and resuspended in a permeabilization buffer (Cytotfix/Cytoperm kit; BD), and stained with the following fluorescently conjugated monoclonal antibodies: anti-CD4 (clone GK1.5)-allophycocyanin (APC), anti-IFN- γ (clone XMG1.2)-fluorescein isothiocyanate (FITC), anti-IFN- γ (clone XMG1.2)-APC, anti-IL-4 (clone GK1.5)-phycoerythrin (PE) (eBiosciences, USA). Fluorescence intensity was measured by flow cytometry (FACS Calibur or FACS CantoII; BD) and data were analysed with FlowJo (Tree star, USA).

Western Blot analysis- DCs derived from C57BL/6 mice were infected with H37Rv or H37Rv Δ ESAT-6. For T cell differentiation, CD4⁺ T cells (1x10⁶) were purified by MACS (CD4⁺ T cell isolation beads kit; Miltenyi Biotech, Germany) from OT-II TCR transgenic mice and co-cultured with DCs

(1x10⁶) infected with H37Rv or H37Rv Δ ESAT-6 in the presence of OVA-peptide (5 μ g/ml). Whole cell lysate was prepared by using lysis buffer (50 mM Tris HCL pH 7.4, 5 mM EDTA, 120 mM NaCl, 0.5% NP40, 0.5 mM NaF, 1 mM dithiothreitol, 0.5 mM phenylmethylsulfonyl fluoride) along with HALTTM phosphatase inhibitor cocktail (78420 Thermoscientific, USA) and protease inhibitor cocktail (78410 Thermoscientific, USA) for 1 hr. Samples were electrophoresed on a 10% SDS-polyacrylamide gel and electroblotted onto polyvinylidene difluoride (PVDF) membranes. Blots were blocked for 1 hr in 5% BSA in PBST (PBS with 0.1% Tween-20). STAT6, pSTAT6, STAT4 and pSTAT4 proteins were detected with anti-STAT6 (Sc-621), -pSTAT6 (Sc-11762-R), -STAT4 (Sc-486) and -pSTAT4 (Sc-16317-R) polyclonal antibodies, respectively, at a dilution of 1:250 and as recommended by the manufacturer (SantaCruz Biotechnology, USA). Goat anti-rabbit-immunoglobulin G-conjugated horseradish peroxidase (Sc-2004) (diluted 1:2500) was used as a secondary antibody (SantaCruz Biotechnology, USA). Immunoblotting for GAPDH was carried out to confirm equal loading.

Statistical analysis- All data were derived from at least three independent experiments. Statistical analyses were conducted using SPSS10 software and values were presented as mean \pm SD. Significant differences between the groups were determined by ANOVA followed by Tukey's multiple comparison test (SPSS software). A value of p<0.05 was accepted as an indication of statistical significance.

Results:

Virulent M. tb H37Rv but not BCG induces IL-1 β and Th2 cells. It is now well established that Th1 cell responses play a protective role (10) whereas Th2 and iTreg cells facilitate progression of TB (11, 18). Previous studies have provided evidence that BCG induces Th1 cell responses (25), whereas virulent H37Rv induces both Th1 and Th2 cell responses (26). Despite adequate Th1 cell responses, *M. tb* continues to progress towards disease in susceptible hosts suggesting that Th1 immunity is not sufficient for optimal host protection. Differentiation of Th1 and Th2 subsets requires IL-12 and IL-4, respectively (14, 27). While IL-

IL-12 produced by professional antigen presenting cells (APCs) drives Th1 cell differentiation (13), professional APCs do not produce IL-4. However, it has been shown that under some circumstances, IL-1 β produced by APCs plays an important role in Th2 cell differentiation (28, 29). Therefore, we determined the cytokines produced by *M. tb* or BCG infected DCs. We generated bone marrow-derived DCs from C57BL/6 mice and infected these cells with H37Rv or BCG. We determined the T cell polarizing cytokines IL-12 and IL-1 β . We found that both H37Rv and BCG induced IL-12 whereas only H37Rv induced IL-1 β (Fig. 1A). We next determined whether these infected DCs differentially facilitate the differentiation of Th cells. To test this hypothesis, we co-cultured CD4⁺ T cells isolated from OT-II TCR Tg mice in the presence of OVA peptide. We observed that H37Rv infected DCs induced large amounts of both IFN- γ and IL-4, whereas BCG infected DCs induced only IFN- γ in T cells (Fig. 1B). This observation suggests that *M. tb* induces either Th0 cells, which produce both IFN- γ and IL-4, or induce distinct populations of Th1 (IFN- γ producers) and Th2 (IL-4 producers) cells. Therefore, to distinguish between these possibilities, we performed intracellular staining of CD4⁺ T cells for IFN- γ and IL-4 production. We found that H37Rv induced few if any CD4⁺ T cells producing both IFN- γ and IL-4 simultaneously. In sharp contrast, BCG induced IFN- γ -producing cells only (Fig. 1C).

RD-1/ESAT-6 induces IL-1 β production and Th2 cell differentiation. From the preceding section it was clear that H37Rv induces both Th1 and Th2 responses, whereas BCG selectively induces Th1 responses. Compared with H37Rv, BCG lacks several genomic segments that are called regions of difference (RD), and a total of 22 such RD regions have been characterized. Among them, RD-1 has been shown to play an important role in the pathogenicity of *M. tb* and its capacity to subvert the immune system (30, 31). Therefore, we tested whether RD-1 has a role in inducing IL-1 β in DCs and to promote Th2 cell responses. ESAT-6 and CFP-10 are two important antigenic proteins encoded within the RD-1 region that are secreted as a complex. Thus, mutation in either protein impairs formation and secretion of the complex (30, 32, 33). Therefore, to understand the role of RD-1 we performed experiments with deletion

mutants of H37Rv in RD-1 (H37Rv Δ RD-1) or ESAT-6 (H37Rv Δ ESAT-6). Interestingly, neither the H37Rv Δ RD-1 mutant nor the H37Rv Δ ESAT-6 mutant was able to induce IL-1 β , whereas both mutants induced levels of IL-12 comparable to H37Rv and BCG (Fig. 2A). Taken together, these observations indicated that the RD-1 region of H37Rv plays an important role in the production of IL-1 β by infected DCs, but has little if any role in the production of IL-12 (Fig. 2A). Next we tested whether cytokines (and possibly IL-1 β) produced by the H37Rv-infected DCs assist in the differentiation of Th2 cells. For this purpose, we co-cultured naïve CD4⁺ T cells from OT-II TCR Tg mice with infected DCs pulsed with OVA peptide. We determined cytokines produced in the culture supernatant as well as cytokines produced by the cells intracellularly. We found that DCs infected with H37Rv induced both IL-4- and IFN- γ -producing CD4⁺ T cells whereas BCG-infected DCs selectively induced IFN- γ -producing CD4⁺ T cells (Fig. 2B). These observations clearly indicate that the RD-1 region of virulent strain H37Rv plays a dominant role in inducing Th2 responses.

Neutralization of IL-1 β abrogates Th2 cell differentiation in T cell co-cultures with infected DCs. To further investigate the role of IL-1 β in Th2 cell differentiation induced by H37Rv, we performed T cell co-culture experiments with infected DCs in the presence of anti-IL-1 β antibodies. Neutralization of IL-1 β with antibody dramatically abrogated IL-4 production by CD4⁺ T cells (Fig. 3A). Interestingly, DCs infected with BCG, H37Rv Δ RD-1 or H37Rv Δ ESAT-6 mutants selectively induced IFN- γ production and this was not affected by anti-IL-1 β antibodies, which thus served as a negative control (Fig. 3B). We also neutralized IL-1 β in co-cultures of T cells with DCs infected with RD-1-complemented BCG. Similar to the results obtained for H37Rv, we found that neutralization of IL-1 β dramatically abrogated Th2 cell differentiation induced by RD-1-complemented BCG. ***To confirm these findings, we investigated the molecular mechanism of Th cell differentiation in infected DCs co-cultured with T cells. For this purpose, we performed Western blot analysis of STAT6, pSTAT6, STAT4 and pSTAT4 proteins in OT-II CD4⁺ T cells from cultures with infected DCs. We observed***

phosphorylation of STAT-6 in CD4⁺ T cells co-cultured with H37Rv infected DCs, and this was dramatically reduced when IL-1 β was neutralized by antibodies (Fig. 3C). In contrast, CD4⁺ T cells co-cultured with H37Rv Δ ESAT-6 were unable to induce STAT-6 phosphorylation. We also examined STAT-4 phosphorylation and found an opposite pattern (Fig. 3C). These data suggest that IL-1 β plays an important role in Th2 cell polarization during *M. tb* infection, and that ESAT-6 or factors controlled by ESAT-6 are responsible for induction of IL-1 β .

Cytokine profiles in lungs of infected animals. Next, we complemented our *in vitro* finding showing that the RD-1 region of virulent strain H37Rv induces IL-1 β to promote Th2 cell differentiation, with *in vivo* experiments. We also performed cytokine profiling studies of lung homogenates obtained from C57BL/6 mice infected with H37Rv, BCG or H37Rv Δ ESAT-6. Consistent with our *in vitro* data, we observed that H37Rv infected animals produced both IL-4 and IFN- γ , whereas BCG infected animals induced only IFN- γ at all time points analysed (Fig. 4A). To obtain further information about the relevant T cell differentiating cytokines, we determined the levels of IL-12 and IL-1 β , as in our *in vitro* experiments. We detected both IL-1 β and IL-12 in lung homogenates from H37Rv-infected mice, whereas BCG infected animals selectively produced IL-12, at levels similar to H37Rv infected animals (Fig. 4A). We also determined cytokines in the lung homogenates of animals infected with H37Rv Δ RD-1 or H37Rv Δ ESAT-6. As expected, both groups of infected animals produced IL-12 and IFN- γ but not IL-1 β in their lungs (Fig 4B). In contrast, RD-1 complemented BCG induced IL-12 as well as IL-1 β , and both IFN- γ and IL-4 were present (Fig. 4B). Collectively these data indicate that the RD-1 region plays an important role in the induction of IL-1 β by H37Rv-infected DCs, which in turn drives Th2 cell differentiation.

*IL-1 β directly acts on CD4⁺ T cells to promote Th2 cell differentiation during *M. tb* infection.* To provide further evidence that IL-1 β plays an important role in the differentiation of Th2 cells in *M. tb* infected animals, we isolated CD4⁺ T cells from congenic wild type Thy1.2 or IL-1 β R^{-/-} Thy1.2 mice and adoptively transferred these cells into irradiated Thy1.1 congenic

animals followed by infection with H37Rv. Thirty days after infection, spleen cells from these animals were isolated and challenged with H37Rv complete soluble antigen (CSA) *ex vivo*, and we stained T cells for the intracellular cytokines IL-4 and IFN- γ . Cytokines produced by adoptively transferred CD4⁺ T cells were determined by gating on Thy1.2. We observed that CD4⁺ T cells from wild type animals had adopted a phenotype characterized by IFN- γ or IL-4 production. In contrast, CD4⁺ T cells from IL-1 β R^{-/-} mice predominantly differentiated into IFN- γ -producing cells (Fig. 5). Therefore, we conclude that IL-1 β produced during the progression of TB disease is responsible for the generation of Th2 cell responses.

Discussion:

M. tb survives within susceptible hosts by altering host protective Th cell responses. IFN- γ -induced macrophage activation that results in the production of nitric oxide radicals is the key mechanism for elimination of the harboured pathogens. Thus, animals that are defective in the production of IL-12, inducible nitric oxide synthase (iNOS), IFN- γ , or proteins involved in their signalling pathways exhibit increased susceptibility to *M. tb* infection (34). However, *M. tb* successfully inhibits IL-12 production in susceptible hosts and thus inhibits the development of Th1 responses (35). In addition, it has clearly been shown that *M. tb* not only prevents Th1 responses, but also facilitates Th2 responses (36), which counter regulate host protective Th1 responses. Phagocytes, the natural home for *M. tb*, are the predominant source of IL-12 that drives Th1 cell differentiation. In contrast, IL-4, the key cytokine required for the differentiation of Th2 cells, is not produced by APCs. Thus, it is likely that another cytokine is involved in promoting Th2 cell differentiation during the progression of TB. Interestingly, previous studies have shown that IL-1 β can act as a co-stimulatory factor for Th2 cells (29). Indeed, we found that IL-1 β is induced by *M. tb* infected DCs. We further showed that IL-1 β is directly involved in Th2 cell differentiation during *M. tb* infection. Therefore, it appears that *M. tb* ensures survival not only by directly inhibiting Th1 responses, but also indirectly by mounting Th2 responses. In fact, it has been reported that inhibition of Th2 responses or genetic deficiency in the

generation of Th2 responses confers partial resistance to *M. tb* infection (37).

Previously, it has been shown that IL-1 β -mediated signalling plays an important role in host resistance to *M. tb* infection (22, 38). In fact, IL-1 β - and IL-1 β R-deficient animals have increased susceptibility to *M. tb* infection (39). Similarly, mice deficient in the IL-1 β signalling adaptor molecule MyD88 are highly susceptible to *M. tb* infection (40). However, the precise nature of the host protective immune responses induced by IL-1 β during *M. tb* infection is unclear. Our results demonstrate that IL-1 β induces Th2 cell differentiation, which has been shown to promote progression of TB by inhibiting host protective Th1 responses (11). Therefore, IL-1 β might originate two divergent immune responses: (i) IL-1 β might induce the MyD88 signalling pathway to promote protective innate host immune responses; and (ii) IL-1 β might have a direct role in the differentiation of Th2 cells, which counter-regulates protective host adaptive immune responses. Therefore, it will be of future interest to delineate the conditions and mechanisms by which IL-1 β elicits these divergent responses.

Previously, we have shown that BCG, *M. tb* H37Rv, and mutant strains H37Rv Δ RD-1 and H37Rv Δ ESAT-6 induce IL-12 in DCs in a toll-like receptor (TLR)-2-independent but MyD88-dependent manner (41), suggesting that production of IL-12 is independent of RD-1. In this report we have shown that IL-1 β production is dependent on RD-1/ESAT-6. Thus, *M. tb* has at least two different sets of factors that control

IL-12 and IL-1 β production to modulate Th cell differentiation during the progression of TB. Furthermore, it is known that Tregs also assist in the progression of TB (19), whereas the role of Th17 cells remains unclear and might differ during primary and secondary infection (41). Taken together, these data indicate that Th1 cells are protective whereas Th2 cells and Treg cells promote disease. These Th cell subsets are in a dynamic balance during *M. tb* infection. In this context, it has been recently shown that mycobacterial strains that can mount Th1 responses but are defective in the induction of Th2 cells or Tregs exhibit dramatically increased vaccine efficacy (36, 42). Consistent with these observations, our recent data indicate that animals that are unable to mount Th2 cells and Tregs (Stat-6^{-/-}TGF- β RIIDN mice) are completely resistant to *M. tb* infection (Bhattacharya et al., under review). Therefore, in future studies it will be interesting to identify the components that are responsible for the induction of Th2 cells and Tregs, which will enable us to design improved vaccines and therapies.

In conclusion, we have demonstrated that the RD-1 region of *M. tb* is responsible for the capacity of this organism to promote Th2 cell differentiation. One of our future goals will be to identify the bacterial component(s) that induce(s) Tregs during TB disease. Our findings suggest that *M. tb* mutants that are defective in the induction of both Th2 cells and Tregs may serve as superior vaccines.

References:

1. WHO. (2007). Global Tuberculosis Control Surveillance, Planning, Financing.
2. WHO. (2004). "BCG Vaccine". *Weekly Epidemiological Record* 79 (4): 27–38. .
3. Almeida Da Silva PE, Palomino JC. (2011). Molecular basis and mechanisms of drug resistance in *Mycobacterium tuberculosis*: classical and new drugs. *J Antimicrob Chemother* 66: 1417-30
4. Davies J. (1996). Origins and evolution of antibiotic resistance. *Microbiologia* 12: 9-16
5. Gillespie SH. (2002). Evolution of drug resistance in *Mycobacterium tuberculosis*: clinical and molecular perspective. *Antimicrob Agents Chemother* 46: 267-74
6. Shah NS, Wright A, Bai GH, Barrera L, Boulahbal F, Martin-Casabona N, Drobniewski F, Gilpin C, Havelkova M, Lepe R, Lumb R, Metchock B, Portaels F, Rodrigues MF, Rusch-Gerdes S, Van Deun A, Vincent V, Laserson K, Wells C, Cegielski JP. (2007). Worldwide emergence of extensively drug-resistant tuberculosis. *Emerg Infect Dis* 13: 380-7
7. Velayati AA, Farnia P, Masjedi MR, Ibrahim TA, Tabarsi P, Haroun RZ, Kuan HO, Ghanavi J, Farnia P, Varahram M. (2009). Totally drug-resistant tuberculosis strains: evidence of adaptation at the cellular level. *Eur Respir J* 34: 1202-3
8. Velayati AA, Masjedi MR, Farnia P, Tabarsi P, Ghanavi J, Ziazarifi AH, Hoffner SE. (2009). Emergence of new forms of totally drug-resistant tuberculosis bacilli: super extensively drug-resistant tuberculosis or totally drug-resistant strains in iran. *Chest* 136: 420-5
9. Flynn JL, Chan J. (2001). Immunology of tuberculosis. *Annu Rev Immunol* 19: 93-129
10. Sweeney KA, Dao DN, Goldberg MF, Hsu T, Venkataswamy MM, Henao-Tamayo M, Ordway D, Sellers RS, Jain P, Chen B, Chen M, Kim J, Lukose R, Chan J, Orme IM, Porcelli SA, Jacobs WR, Jr. (2011). A recombinant *Mycobacterium smegmatis* induces potent bactericidal immunity against *Mycobacterium tuberculosis*. *Nat Med* 17: 1261-8
11. Lienhardt C, Azzurri A, Amedei A, Fielding K, Sillah J, Sow OY, Bah B, Benagiano M, Diallo A, Manetti R, Manneh K, Gustafson P, Bennett S, D'Elis MM, McAdam K, Del Prete G. (2002). Active tuberculosis in Africa is associated with reduced Th1 and increased Th2 activity in vivo. *Eur J Immunol* 32: 1605-13
12. Cooper AM, Solache A, Khader SA. (2007). Interleukin-12 and tuberculosis: an old story revisited. *Curr Opin Immunol* 19: 441-7
13. Trinchieri G. (2003). Interleukin-12 and the regulation of innate resistance and adaptive immunity. *Nat Rev Immunol* 3: 133-46
14. Hickman SP, Chan J, Salgame P. (2002). *Mycobacterium tuberculosis* induces differential cytokine production from dendritic cells and macrophages with divergent effects on naive T cell polarization. *J Immunol* 168: 4636-42
15. Chen W, Konkel JE. (2010). TGF-beta and 'adaptive' Foxp3(+) regulatory T cells. *J Mol Cell Biol* 2: 30-6
16. Scott-Browne JP, Shafiani S, Tucker-Heard G, Ishida-Tsubota K, Fontenot JD, Rudensky AY, Bevan MJ, Urdahl KB. (2007). Expansion and function of Foxp3-expressing T regulatory cells during tuberculosis. *J Exp Med* 204: 2159-69
17. Yoshimura A, Wakabayashi Y, Mori T. (2010). Cellular and molecular basis for the regulation of inflammation by TGF-beta. *J Biochem* 147: 781-92
18. Kursar M, Koch M, Mittrucker HW, Nouailles G, Bonhagen K, Kamradt T, Kaufmann SH. (2007). Cutting Edge: Regulatory T cells prevent efficient clearance of *Mycobacterium tuberculosis*. *J Immunol* 178: 2661-5

19. Shafiani S, Tucker-Heard G, Kariyone A, Takatsu K, Urdahl KB. (2010). Pathogen-specific regulatory T cells delay the arrival of effector T cells in the lung during early tuberculosis. *J Exp Med* 207: 1409-20
20. Bettelli E, Carrier Y, Gao W, Korn T, Strom TB, Oukka M, Weiner HL, Kuchroo VK. (2006). Reciprocal developmental pathways for the generation of pathogenic effector TH17 and regulatory T cells. *Nature* 441: 235-8
21. Mangan PR, Harrington LE, O'Quinn DB, Helms WS, Bullard DC, Elson CO, Hatton RD, Wahl SM, Schoeb TR, Weaver CT. (2006). Transforming growth factor-beta induces development of the T(H)17 lineage. *Nature* 441: 231-4
22. Juffermans NP, Florquin S, Camoglio L, Verbon A, Kolk AH, Speelman P, van Deventer SJ, van Der Poll T. (2000). Interleukin-1 signaling is essential for host defense during murine pulmonary tuberculosis. *J Infect Dis* 182: 902-8
23. Mayer-Barber KD, Barber DL, Shenderov K, White SD, Wilson MS, Cheever A, Kugler D, Hieny S, Caspar P, Nunez G, Schlueter D, Flavell RA, Sutterwala FS, Sher A. (2010). Caspase-1 independent IL-1beta production is critical for host resistance to mycobacterium tuberculosis and does not require TLR signaling in vivo. *J Immunol* 184: 3326-30
24. Ben-Sasson SZ, Hu-Li J, Quiel J, Cauchetaux S, Ratner M, Shapira I, Dinarello CA, Paul WE. (2009). IL-1 acts directly on CD4 T cells to enhance their antigen-driven expansion and differentiation. *Proc Natl Acad Sci U S A* 106: 7119-24
25. Fine PE. (1995). Variation in protection by BCG: implications of and for heterologous immunity. *Lancet* 346: 1339-45
26. Bhattacharyya S, Singla R, Dey AB, Prasad HK. (1999). Dichotomy of cytokine profiles in patients and high-risk healthy subjects exposed to tuberculosis. *Infect Immun* 67: 5597-603
27. Noben-Trauth N, Hu-Li J, Paul WE. (2002). IL-4 secreted from individual naive CD4+ T cells acts in an autocrine manner to induce Th2 differentiation. *Eur J Immunol* 32: 1428-33
28. Hebel K, Rudolph M, Kosak B, Chang HD, Butzmann J, Brunner-Weinzierl MC (2011). IL-1beta and TGF-beta act antagonistically in induction and differentially in propagation of human proinflammatory precursor CD4+ T cells. *J Immunol* 187: 5627-35
29. Mejri N, Brossard M. (2007). Splenic dendritic cells pulsed with Ixodes ricinus tick saliva prime naive CD4+T to induce Th2 cell differentiation in vitro and in vivo. *Int Immunol* 19: 535-43
30. Guinn KM, Hickey MJ, Mathur SK, Zakel KL, Grotzke JE, Lewinsohn DM, Smith S, Sherman DR. (2004). Individual RD1-region genes are required for export of ESAT-6/CFP-10 and for virulence of Mycobacterium tuberculosis. *Mol Microbiol* 51: 359-70
31. Sherman DR, Guinn KM, Hickey MJ, Mathur SK, Zakel KL, Smith S. (2004). Mycobacterium tuberculosis H37Rv: Delta RD1 is more virulent than M. bovis bacille Calmette-Guerin in long-term murine infection. *J Infect Dis* 190: 123-6
32. McLaughlin B, Chon JS, MacGurn JA, Carlsson F, Cheng TL, Cox JS, Brown EJ. (2007). A mycobacterium ESX-1-secreted virulence factor with unique requirements for export. *PLoS Pathog* 3: e105
33. Majlessi L, Brodin P, Brosch R, Rojas MJ, Khun H, Huerre M, Cole ST, Leclerc C. (2005). Influence of ESAT-6 secretion system 1 (RD1) of Mycobacterium tuberculosis on the interaction between mycobacteria and the host immune system. *J Immunol* 174: 3570-9

34. Flynn JL, Chan J, Triebold KJ, Dalton DK, Stewart TA, Bloom BR. (1993). An essential role for interferon gamma in resistance to *Mycobacterium tuberculosis* infection. *J Exp Med* 178: 2249-54
35. Pathak SK, Basu S, Bhattacharyya A, Pathak S, Kundu M, Basu J. (2005). *Mycobacterium tuberculosis* lipoarabinomannan-mediated IRAK-M induction negatively regulates Toll-like receptor-dependent interleukin-12 p40 production in macrophages. *J Biol Chem* 280: 42794-800
36. Rook GA. (2007). Th2 cytokines in susceptibility to tuberculosis. *Curr Mol Med* 7: 327-37
37. Potian JA, Rafi W, Bhatt K, McBride A, Gause WC, Salgame P. (2011). Preexisting helminth infection induces inhibition of innate pulmonary anti-tuberculosis defense by engaging the IL-4 receptor pathway. *J Exp Med* 208: 1863-74
38. Wilkinson RJ, Patel P, Llewelyn M, Hirsch CS, Pasvol G, Snounou G, Davidson RN, Toossi Z. (1999). Influence of polymorphism in the genes for the interleukin (IL)-1 receptor antagonist and IL-1beta on tuberculosis. *J Exp Med* 189: 1863-74
39. Yamada H, Mizumo S, Horai R, Iwakura Y, Sugawara I. (2000). Protective role of interleukin-1 in mycobacterial infection in IL-1 alpha/beta double-knockout mice. *Lab Invest* 80: 759-67
40. Fremont CM, Togbe D, Doz E, Rose S, Vasseur V, Maillet I, Jacobs M, Ryffel B, Quesniaux VF. (2007). IL-1 receptor-mediated signal is an essential component of MyD88-dependent innate response to *Mycobacterium tuberculosis* infection. *J Immunol* 179: 1178-89
41. Chatterjee S, Dwivedi VP, Singh Y, Siddiqui I, Sharma P, Van Kaer L, Chattopadhyay D, Das G. (2011). Early secreted antigen ESAT-6 of *Mycobacterium tuberculosis* promotes protective T helper 17 cell responses in a toll-like receptor-2-dependent manner. *PLoS Pathog* 7: e1002378
42. Koch MA, Tucker-Heard G, Perdue NR, Killebrew JR, Urdahl KB, Campbell DJ. (2009). The transcription factor T-bet controls regulatory T cell homeostasis and function during type 1 inflammation. *Nat Immunol* 10: 595-602

Legends to Figures:

Figure 1. Capacity of *M. tb* strains to induce IL-1 β secretion in DCs co-cultured with OT-II CD4⁺ T cells and to promote Th2 cell differentiation. (A) IL-12p40 and IL-1 β levels after infection of DCs with H37Rv or BCG. IL-1 β levels were significantly increased ($p < 0.01$) in H37Rv infected DCs as compared with DCs infected with BCG. (B) IFN- γ and IL-4 levels after infection of DCs with H37Rv or BCG. IL-4 levels were significantly higher ($p < 0.01$) in H37Rv infected DCs than BCG infected DCs. (C) Intracellular IFN- γ and IL-4 levels produced by CD4⁺ T cells from OT-II TCR transgenic mice co-cultured with DCs infected with H37Rv or BCG. The results shown are representative of at least 3-4 independent experiments.

Figure 2. Cytokine profiling from DCs infected with H37Rv or mutant strains. (A) Measurements of levels of IL-12p40 and IL-1 β from supernatants of DCs infected with H37Rv or mutant strains (H37Rv Δ RD1, H37Rv Δ ESAT-6, or ESAT-6-complemented H37Rv Δ ESAT-6) of H37Rv at different points. (B) Levels of IFN- γ and IL-4 from the culture supernatants of DCs infected with H37Rv or mutants (H37Rv Δ RD1, H37Rv Δ ESAT-6, or ESAT-6-complemented H37Rv Δ ESAT-6). H37Rv and ESAT-6-complemented H37Rv Δ ESAT-6 (Comp-ESAT-6) direct differentiation of both Th1 (IL-12p40 and IFN- γ) and Th2 (IL-4) cytokine responses as well as IL-1 β production, whereas H37Rv Δ RD1 and H37Rv Δ ESAT-6 induce only Th1 cytokines in infected DCs co-cultured with naïve OT-II CD4⁺ T cells. IL-4 and IL-1 β levels were significantly increased ($p < 0.01$) in H37Rv and Comp-ESAT-6 infected DCs as compared with DCs infected with the other strains. The results shown are representative of at least three independent experiments.

Figure 3. ESAT-6 induces Th2 cell differentiation via IL-1 β production. (A) IL-4 production in DCs co-cultured with naïve OT-II CD4⁺ T cells after infection with H37Rv or BCG in the presence or absence of anti-IL-1 β . (B) IL-4 production in DCs co-cultured with naïve OT-II CD4⁺ T cells after infection with H37Rv or mutant strains (H37Rv Δ RD1, H37Rv Δ ESAT-6 and Comp-ESAT-6) in the presence or absence of anti-IL-1 β . IL-4 levels were higher in cultures with H37Rv or Comp-ESAT-6 as compared with cultures with BCG or other mutant strains ($p < 0.003$). (C) *Western blot analysis of STAT6, pSTAT6, STAT4 and pSTAT4 proteins from DCs infected with H37Rv or H37Rv Δ ESAT-6 and co-cultured with OT-II CD4⁺ T cells, to show modulated Th1 and Th2 responses in the presence or absence of anti-IL-1 β .* The results shown are representative of at least four independent experiments.

Figure 4. H37Rv induces significantly higher Th2 responses and the proinflammatory cytokine IL-1 β as compared with BCG or H37Rv Δ ESAT-6. (A) Cytokine levels in lung homogenates of mice at different time points after infection with H37Rv or BCG. (B) Cytokine levels in lung homogenates of mice at different time points after infection with H37Rv or mutant strain H37Rv Δ ESAT-6. Statistical significance between different strains was determined by one-way ANOVA: * $p < 0.01$; ** $p < 0.017$, *** $p < 0.001$. The results shown are representative of at least three independent experiments with six mice within each group.

Figure 5. IL-1 β plays an important role in Th2 cell differentiation after infection with H37Rv without affecting Th1 cell differentiation. CD4⁺ T cells were isolated from congenic wild type Thy1.2 or IL-1 β R^{-/-} Thy1.2 mice and adoptively transferred into irradiated Thy1.1 congenic animals followed by infection with H37Rv. Thirty days after infection, spleen cells from these animals were isolated and challenged with H37Rv complete soluble antigen (CSA) *ex vivo*. T cells were then stained for the intracellular cytokines IL-4 and IFN- γ . The results shown are representative of at least three independent experiments with three mice within each group.






Fig.1. Dwivedi et al

A



B






Fig.2. Dwivedi et al

A



B



C





Fig.3. Dwivedi et al

A



B




Fig.4. Dwivedi et al




Fig.5. Dwivedi et al

Mycobacterium tuberculosis directs T helper 2 cell differentiation by inducing interleukin-1 β production in dendritic cells

Ved Prakash Dwivedi, Debapriya Bhattacharya, Samit Chatterjee, Debprasad Chattopadhyay, Luc Van Kaer, William R. Bishai and Gobardhan Das

J. Biol. Chem. published online July 18, 2012

Access the most updated version of this article at doi: [10.1074/jbc.M112.375154](https://doi.org/10.1074/jbc.M112.375154)

Alerts:

- [When this article is cited](#)
- [When a correction for this article is posted](#)

[Click here](#) to choose from all of JBC's e-mail alerts