

## Positive solutions of the semilinear Dirichlet problem with critical growth in the unit disc in $\mathbb{R}^2$

ADIMURTHI

TIFR Centre, P.B. No. 1234, Bangalore 560 012, India

MS received 28 June 1988

**Abstract.** We prove the existence of a positive solution of the following problem

$$-\Delta u = f(r, u) \quad \text{in } D$$

$$u > 0$$

$$u = 0, \quad \text{on } \partial D$$

where  $D$  is the unit disc in  $\mathbb{R}^2$  and  $f$  is a superlinear function with critical growth.

**Keywords.** Sub-critical growth, critical growth, super critical growth; Laplacian; Palais-Smale condition; Semilinear Dirichlet problem; unit disc.

### Introduction

$D$  be the unit disc in  $\mathbb{R}^2$ . We are looking for positive radial solutions of the following problem: Find  $u$  in  $C^2(D) \cap C^0(\bar{D})$  such that

$$\begin{aligned} -\Delta u &= f(r, u) \quad \text{in } D \\ u &> 0 \\ u &= 0, \quad \text{on } \partial D \end{aligned} \tag{1.1}$$

where  $f$  is superlinear,  $f(r, 0) = 0$ ,  $(\partial f / \partial t)(r, 0) < \lambda_1$  with  $\lambda_1$  being the first eigenvalue of the Dirichlet problem. For  $n \geq 3$  and  $f$  of critical growth, Brezis–Nirenberg [4] studied the existence and non-existence of solutions of problem (1.1). For  $n = 2$ , the critical growth is of exponential type whereas in the case of  $n \geq 3$ , it is of polynomial type and the method adopted for  $n \geq 3$  fails in the case of  $n = 2$ .

Carleson–Chang [5] obtained a positive solution for  $f(u) = \lambda u \exp(\lambda u^2)$  with  $\lambda < \lambda_1$  via a variational method. For growths of type  $f(u) = u^m \exp(bu^2)$ , Atkinson–Gutierrez [3] used the shooting argument to obtain a solution of (1.1). They assumed that  $\log f$  is strictly convex for large  $u$ .

In this paper we relax the conditions on  $f$  and use a variational method to obtain a solution of (1.1). Since we are interested in radial solutions, (1.1) is equivalent to

finding an  $u$  in  $C^2(D) \cap C^0(\bar{D})$  with  $u$  radial and satisfying

$$\begin{aligned} L_1 u &\equiv -(ru')' = f(r, u)r \quad \text{in } [0, 1) \\ u &> 0 \quad \text{in } [0, 1) \\ u'(0) &= u(1) = 0. \end{aligned} \tag{1.2}$$

where  $u' = du/dr$ .

The idea of the method is to approximate the energy functional by functionals satisfying Palais–Smale conditions. Then obtain the critical points of these approximate functionals by a constrained minimization problem similar to that of Zeev–Nehari [8] and then pass to the limit. The method of the proof is in the spirit of Brezis–Nirenberg [4]. Here, we also get a constant “ $a$ ” which is strictly less than the best possible constant and thereby the existence of solutions of (1.2) is guaranteed.

In [1] we also prove the existence of infinitely many solutions of (1.1) when  $f$  is odd and of critical growth. Also in [2] we prove the existence of solutions of (1.1) if  $D$  is replaced by an arbitrary smooth domain.

## 2. Statements

Let  $E = \{u \in C^1[0, 1]; u(1) = 0\}$ . For  $0 \leq \alpha \leq 1$  and  $u$  in  $E$  define

$$\begin{aligned} |u|_\alpha^2 &= \int_0^1 u^2(r) r^\alpha dr \\ \|u\|_\alpha^2 &= \int_0^1 u'(r)^2 r^\alpha dr. \end{aligned}$$

Let  $H_\alpha^*$  be the completion of  $E$  with respect to  $\|\cdot\|_\alpha$ . Define the operator  $L_\alpha$  by

$$L_\alpha = -\frac{1}{r^\alpha} \frac{d}{dr} \left( r^\alpha \frac{d}{dr} \right). \tag{2.1}$$

Let  $(\lambda_\alpha, \phi_\alpha)$  be the first eigenvalue and the corresponding first eigenvector with  $\phi_\alpha(0) = 1$  of the following eigenvalue problem.

$$\begin{aligned} L_\alpha \phi &= \lambda \phi \quad \text{in } [0, 1] \\ \phi'(0) &= \phi(1) = 0. \end{aligned} \tag{2.2}$$

### DEFINITION 2.1

Let  $f: [0, 1] \times [0, \infty) \rightarrow [0, \infty)$  be a  $C^1$ -function. We say  $f$  is of class **A** if

- (i)  $f(r, 0) = 0$ .
- (ii) There exists a  $\delta_0 > 0$  and for  $(r, t) \in Q_{\delta_0} \equiv [0, \delta_0] \times [0, \infty)$   $(\partial f / \partial r)(r, t) \geq 0$ .
- (iii) There exists a  $t_0 > 0$  such that  $f(r, t) < \lambda_1 t$  for all  $(r, t) \in [0, 1] \times [0, t_0]$ .
- (iv) There exist constants  $t_1 > 0$ ,  $\beta > 2$  such that  $\beta F(r, t) \leq f(r, t)t$  for all  $(r, t) \in [0, 1] \times [t_1, \infty)$  where  $F(r, t) = \int_0^t f(r, s) ds$ .

Let

$$A' = \left\{ f \in A; \frac{\partial f}{\partial t} > \frac{f}{t} \text{ in } [0, 1] \times (0, \infty) \right\}.$$

We consider the following three types of functions in our discussions.

*Sub-critical:*  $f$  in  $A$  is said to be sub-critical if there exists a  $\delta > 0$  and for every  $\varepsilon > 0$

$$\sup_{(r,t) \in [0,\delta] \times [0,\infty)} f(r,t) \exp(-\varepsilon t^2) < \infty \quad (2.3)$$

*Critical:*  $f$  in  $A'$  is said to be critical if there exists  $\delta_1 > 0$  such that

- (i)  $f(r,t) = h(r,t) \exp(h(r)t^2) \quad \forall (r,t) \in Q_{\delta_1} \equiv [0,\delta_1] \times [0,\infty)$   
 (ii)  $\forall \varepsilon > 0,$

$$\sup_{(r,t) \in Q_{\delta_1}} h(r,t) \exp(-\varepsilon t^2) < \infty. \quad (2.4)$$

- (iii) For every  $c > 0$ ,  $h(0,t) \exp(ct^2) \rightarrow \infty$  as  $t \rightarrow \infty$ .

*Super critical:*  $f \in A'$  is said to be super critical if for every  $c > 0$

$$\sup_{w \rightarrow 1^-} \int_0^1 f(r,cw) w r dr = \infty. \quad (2.5)$$

For  $f \in A$ ,  $0 \leq x \leq 1$ , let  $\Sigma_x$  be the set of  $C^2$ -solutions of the following problem

$$\begin{aligned} L_x u &= f(r,u) \quad \text{in } [0,1] \\ u &> 0 \\ u'(0) &= u(1) = 0. \end{aligned} \quad (2.6)$$

## DEFINITION 2.2

$u$  in  $H_0^1(D)$  is said to be a weak solution of (1.2) if

- (i)  $u > 0$  in  $[0,1]$   
 (ii)  $\int_0^1 f(r,u) u r dr < \infty$   
 (iii)  $\forall \phi \in C^2[0,1]$  with  $\phi(1) = 0$

$$\int_0^1 u(L_1 \phi) r dr = \int_0^1 f(r,u) \phi r dr.$$

Since we are interested in only positive solutions of (1.2) and hence extending  $f$  for  $t \leq 0$  is irrelevant. Therefore we make the following conventions.

- 1) Whenever we say  $f$  is in  $A$ , then we extend  $f$  by  $f(r,t) = 0$  for  $t \leq 0$  and  $r \in [0,1]$ .  
 2) Whenever we say  $f$  is in  $A'$ , then we extend  $f$  by  $f(r,t) = -f(r,-t)$  for  $t \leq 0$ . (2.8)

For  $u$  in  $H_0^\alpha$ , define

$$\bar{I}_\alpha(u) = \frac{1}{2} \|u\|_\alpha^2 - \int_0^1 F(r, u) r^\alpha dr. \quad (2.9)$$

$$l_\alpha = \inf_{\Sigma_\alpha} \bar{I}_\alpha.$$

Then we have

**Theorem 2.1.** *Let  $f$  be in  $A$ . Then there exists an  $\alpha_0 < 1$  such that for every  $\alpha_0 \leq \alpha < 1$ ,  $\Sigma_\alpha$  is non-empty and  $\{l_\alpha\}$  is bounded. Let  $l = \lim_{\alpha \rightarrow 1} l_\alpha$ . Suppose there exists  $b > 0$ ,  $M > 0$  such that*

- (i)  $f(r, t) \leq M \exp(bt^2)$  for all  $(r, t) \in [0, \delta] \times [0, \infty)$
  - (ii)  $bl < 1$ .
- (2.10)

Then there exists a solution  $u$  of (1.2).

#### COROLLARY 2.1

If  $f$  is sub-critical, then there exists a solution.

*Proof.* If  $f$  is sub-critical, we can take  $b$  as small as we want and satisfying (i) and (ii) of Theorem (2.1). Hence the solution exists.

*Criterion to satisfy (2.10).* Let  $f$  be in  $A$  satisfying (i) of Theorem (2.1). Suppose there exists an  $m > 0$  such that

$$\int_0^{1/2} F\left(r, \frac{m}{2}\right) r dr \geq 2m^2. \quad (2.11)$$

$$2m^2 b < 1$$

Then  $f$  satisfies (ii) of Theorem (2.1).

For  $f$  in  $A^1$  and for  $0 \leq \alpha < 1$ , define

$$\begin{aligned} B_\alpha &= \left\{ u \in H_0^\alpha \setminus \{0\}; \|u\|_\alpha^2 \leq \int_0^1 f(r, u) u r^\alpha dr \right\} \\ \partial B_\alpha &= \left\{ u \in B_\alpha; u \geq 0; \|u\|_\alpha^2 = \int_0^1 f(r, u) u r^\alpha dr \right\} \\ B_1 &= \left\{ u \in H_0^1 \cap L^\infty \setminus \{0\}; \|u\|_1^2 \leq \int_0^1 f(r, u) u r dr \right\} \\ B_1^* &= \left\{ u \in H_0^1 \setminus \{0\}; u \text{ is non-increasing, } \|u\|_1^2 \leq \int_0^1 f(r, u) u r dr \right\} \\ \partial(B_1 \cup B_1^*) &= \left\{ u \in B_1 \cup B_1^*; u \geq 0; \|u\|_1^2 = \int_0^1 f(r, u) u r dr \right\} \\ B_{01} &= \{u \in B_1; u \text{ is constant in a nhd of zero}\}. \\ \partial B_{01} &= \left\{ u \in B_{01}; u \geq 0; \|u\|_1^2 = \int_0^1 f(r, u) u r dr \right\} \end{aligned}$$

For  $0 \leq \alpha \leq 1$ ,  $f \in A'$ ,  $u$  in  $H_0^\alpha$ , define

$$I_\alpha(u) = \frac{1}{2} \int_0^1 f(r, u) u r^\alpha dr - \int_0^1 F(r, u) r^\alpha dr \quad (2.13)$$

since  $f \in A'$ ;  $f(r, t)t - 2F(r, t) \geq 0$  for all  $(r, t) \in [0, 1] \times \mathbb{R}$ , hence  $I_\alpha(u) \geq 0$ . Define  $a_\alpha$  by

$$\frac{a_\alpha^2}{2} = \inf_{\Sigma_\alpha} I_\alpha. \quad (2.14)$$

**Theorem 2.2.** Let  $f$  be in  $A'$ . Then there exists an  $\alpha_0 < 1$  such that for  $\alpha_0 \leq \alpha < 1$ ,  $\Sigma_\alpha$  is non-empty and  $\{a_\alpha\}$  is bounded and satisfying

$$\frac{a_\alpha^2}{2} = \inf_{B_\alpha} I_\alpha(u) = \inf_{\partial B_\alpha} I_\alpha(u). \quad (2.15)$$

Case 1. If  $f$  is super critical then  $\lim_{\alpha \rightarrow 1} a_\alpha = 0$ .

Case 2. If  $f$  is critical and suppose there exists a  $t_2 > 0$  such that

$$\begin{aligned} t_2 h\left(0, \left(\frac{2}{b(0)}\right)^{1/2} t_2\right) &> 2\left(\frac{2}{b(0)}\right)^{1/2} \\ \exp(-t_2) &< \delta_1 \quad [\text{see (2.4)}] \end{aligned} \quad (2.16)$$

then  $\lim_{\alpha \rightarrow 1} a_\alpha = a$  exists and is non-zero. Moreover there exists  $u$  satisfying (1.2) such that

$$I_1(u) = \frac{a^2}{2} = \inf_{B_1 \cup B_1^*} I_1 = \inf_{B_{01}} I_1 = \inf_{\partial B_{01}} I_1 \quad (2.17)$$

**Remark 2.1.** Suppose there exists a sequence  $t_n \rightarrow \infty$  such that  $h(0, t_n)t_n \rightarrow \infty$ , then (2.16) is satisfied.

### Examples

1. *Carleson-Chang.* Let  $f_\lambda(t) = \lambda t \exp(\lambda t^2)$  for  $0 < \lambda < \lambda_1$ . Then  $f_\lambda$  is in  $A'$  and satisfies (2.16). Hence (1.1) has a solution.

2. *Atkinson-Peletier.*  $f(t) = t^m \exp(bt^2)$ ,  $m > 1$ ,  $b > 0$ . Then  $f$  is in  $A'$  satisfying (2.16). Hence (1.1) has a solution.

3.  $f(t) = \lambda t^m \exp(bt^2 + \sin t^2)$ ,  $b \geq 1$

$$m = 1, \quad 0 < \lambda < \lambda_1,$$

$$m > 1, \quad \lambda > 0.$$

Then  $f$  is in  $A'$  and satisfying (2.16). Hence (1.1) has a solution. Here  $\log f$  is not convex for large  $t$ .

4. Let  $b(r)$  be a  $C^1$ -function on  $[0, 1]$  such that  $0 \leq b(r) \leq 1$ ,  $b(r) \equiv 1$  in a neighbourhood of zero. Let  $f(r, t) = t^m \exp(b(r)t^2 + (1 - b(r))\exp(t))$ . Then  $f$  is in  $A'$  satisfying (2.16). Hence (1.1) has a solution.

### 3. Proofs of theorems (2.1) and (2.2)

*Lemma 3.1.* For  $0 \leq \alpha < 1$ , we have

- (i)  $H_0^\alpha$  is compactly embedded in  $C[0, 1]$ .
- (ii)  $\lambda_\alpha < \lambda_1$  and  $\lambda_\alpha \rightarrow \lambda_1$  as  $\alpha \rightarrow 1$
- (iii)  $u$  in  $H_0^1$ ,  $r_1 < r_2$ ,

$$|u(r_1) - u(r_2)|^2 \leq \|u\|_1^2 \log \frac{r_2}{r_1}.$$

*Proof.* Let  $r_1 \leq r_2$  and  $u$  is in  $H_0^\alpha$ . Then by integration by parts

$$\begin{aligned} |u(r_2) - u(r_1)|^2 &= \left( \int_{r_1}^{r_2} u'(r) dr \right)^2 \\ &\leq \|u\|_\alpha^2 \int_{r_1}^{r_2} r^{-\alpha} dr \\ &= \|u\|_\alpha^2 \frac{r_2^{1-\alpha} - r_1^{1-\alpha}}{1-\alpha}. \end{aligned} \quad (3.1)$$

Hence (i) follows from (3.1) and Arzela–Ascoli's theorem. Let  $u$  is in  $H_0^1$ , then

$$\begin{aligned} |u(r_2) - u(r_1)|^2 &= \left( \int_{r_1}^{r_2} u'(r) dr \right)^2 \\ &\leq \|u\|_1^2 \left( \int_{r_1}^{r_2} r^{-1} dr \right) \\ &= \|u\|_1^2 \log \frac{r_2}{r_1}. \end{aligned} \quad (3.2)$$

This proves (iii).

We have

$$\begin{aligned} -(r\phi'_\alpha)' &= \lambda_\alpha \phi_\alpha r - (1-\alpha)\phi'_\alpha \\ -(r\phi'_1)' &= \lambda_1 \phi_1 r. \end{aligned}$$

Hence

$$\begin{aligned} \lambda_1 \int_0^1 \phi_1 \phi_\alpha r dr &= - \int_0^1 (r\phi'_1)' \phi_\alpha dr \\ &= - \int_0^1 (r\phi'_1)' \phi_\alpha dr \\ &= \lambda_\alpha \int_0^1 \phi_\alpha \phi_1 r dr - (1-\alpha) \int_0^1 \phi'_\alpha \phi_1 dr. \end{aligned}$$

i.e.

$$(\lambda_1 - \lambda_\alpha) \int_0^1 \phi_1 \phi_\alpha r dr = -(1-\alpha) \int_0^1 \phi'_\alpha \phi_1 dr.$$

Since  $\phi'_\alpha \leq 0$  and hence  $\lambda_\alpha \leq \lambda_1$  and  $\lambda_\alpha \rightarrow \lambda_1$  as  $\alpha \rightarrow 1$ . This proves (ii).

**Lemma 3.2.** *Let  $f$  be in  $A$ , then there exists an  $\alpha_0 < 1$  such that for  $\alpha_0 \leq \alpha < 1$ ,*

i)  $\bar{I}_\alpha$  satisfies the Palais-Smale condition.

ii) Let  $m > 0$  be such that

$$\int_0^{1/2} F\left(r, \frac{m}{2}\right) r \, dr \geq 2m^2 \quad (3.3)$$

[Such a  $m$  exists because of the condition (iv) of definition (2.1)].

Then there exists a  $u_\alpha$  in  $C^2[0, 1]$  satisfying

$$\begin{aligned} L_\alpha u_\alpha &= f(r, u_\alpha) \quad \text{in } [0, 1] \\ u_\alpha &> 0 \end{aligned} \quad (3.4)$$

$$u'_\alpha(0) = u_\alpha(1) = 0.$$

and

$$\bar{I}_\alpha(u_\alpha) \leq 2m^2.$$

*Proof.* Proof of this lemma is standard (see [7]). For the sake of completeness we will prove it.

**Step 1.** Let  $u_n$  in  $H_0^\alpha$  be a sequence such that

$$|\bar{I}_\alpha(u_n)| \leq M \quad (3.5)$$

$$\bar{I}'_\alpha(u_n) \rightarrow 0 \quad \text{as } n \rightarrow \infty.$$

$$\begin{aligned} \beta \bar{I}_\alpha(u_n) - \langle \bar{I}'_\alpha(u_n), u_n \rangle &= \left(\frac{\beta}{2} - 1\right) \int_0^1 u'_n(r)^2 r^\alpha \, dr - \int_0^1 [\beta F(r, u_n) - f(r, u_n)u_n] r^\alpha \, dr \\ &\geq \left(\frac{\beta}{2} - 1\right) \int_0^1 u'_n(r)^2 r^\alpha \, dr - \int_{|u_n| \leq t_1} [\beta F(r, u_n) - f(r, u_n)u_n] r^\alpha \, dr \\ &\geq \left(\frac{\beta}{2} - 1\right) \int_0^1 u'_n(r)^2 r^\alpha \, dr + C, \end{aligned} \quad (3.6)$$

where  $C$  is a constant depending only on  $F$ . Since  $\beta > 2$ , (3.5) and (3.6) imply  $\{\|u_n\|_\alpha\}$  is bounded. Let  $u_n$  converge to  $u$  weakly in  $H_0^\alpha$  and strongly in  $C[0, 1]$ .

$$\begin{aligned} \langle \bar{I}'_\alpha(u_n), u_n - u \rangle &= \int_0^1 u'_n(r)^2 r^\alpha \, dr - \int_0^1 u'_n(r)u'(r) r^\alpha \, dr \\ &\quad - \int_0^1 f(r, u_n)(u_n - u) r^\alpha \, dr \end{aligned} \quad (3.7)$$

(3.5) and (3.7) imply

$$\int_0^1 u'_n(r)^2 r^\alpha \, dr \rightarrow \int_0^1 u'(r)^2 r^\alpha \, dr.$$

Hence  $u_n$  converges strongly to  $u$  and this proves (i).

*Step 2.* From (ii) of Lemma (3.1) and (iii) of Definition (2.1) there exists an  $\alpha_0 < 1$  and a  $\lambda > 0$  such that

$$F(r, t) \leq \frac{\lambda t^2}{2} < \frac{\lambda_{\alpha} t^2}{2} \quad \text{for all } r \in [0, 1], \quad 0 < |t| < t_0. \quad (3.8)$$

Let  $u$  in  $H_0^{\alpha}$  be such that

$$\|u\|_{\alpha}^2 \leq \frac{(1-\alpha)}{2} t_0^2. \quad (3.9)$$

From (3.1) and (3.9) we have

$$|u(r)|^2 \leq t_0^2. \quad (3.10)$$

Hence (3.8) and (3.10) give

$$F(r, u(r)) \leq \frac{\lambda u(r)^2}{2} \quad (3.11)$$

$$\begin{aligned} \bar{I}_{\alpha}(u) &= \frac{1}{2} \|u\|_{\alpha}^2 - \int_0^1 F(r, u) r^{\alpha} dr \\ &\geq \frac{1}{2} \|u\|_{\alpha}^2 - \frac{\lambda}{2} \int_0^1 u(r)^2 r^{\alpha} dr \\ &\geq \frac{1}{2} \left[ \|u\|_{\alpha}^2 - \frac{\lambda}{\lambda_{\alpha}} \|u\|_{\alpha}^2 \right] \\ &= \frac{1}{2} \left( 1 - \frac{\lambda}{\lambda_{\alpha}} \right) \|u\|_{\alpha}^2. \end{aligned} \quad (3.12)$$

Hence zero is a local minima.

*Step 3.* Define  $u_0$  in  $H_0^0$  by

$$u_0(r) = \begin{cases} \frac{m}{2} & 0 \leq r < \frac{1}{2} \\ m(1-r) & \frac{1}{2} \leq r \leq 1 \end{cases} \quad (3.13)$$

Then

$$\begin{aligned} \bar{I}_{\alpha}(u_0) &= \frac{1}{2} \int_{1/2}^1 m^2 r^{\alpha} dr - \int_0^1 F(r, u_0) r^{\alpha} dr \\ &\leq \frac{m^2}{2(1+\alpha)} \left( 1 - \frac{1}{2^{1+\alpha}} \right) - \int_0^{1/2} F(r, u_0) r^{\alpha} dr \\ &\leq \frac{m^2}{2(1+\alpha)} \left( 1 - \frac{1}{2^{1+\alpha}} \right) - \int_0^{1/2} F\left(r, \frac{m}{2}\right) r^{\alpha} dr \\ &\leq \frac{m^2}{2(1+\alpha)} \left( 1 - \frac{1}{2^{1+\alpha}} \right) - 2m^2 < 0 \end{aligned} \quad (3.14)$$



and for  $0 \leq t \leq 1$ ,

$$\begin{aligned} \bar{I}_\alpha(tu_0) &\leq \frac{t^2}{2} \|u_0\|_\alpha^2 \\ &\leq \frac{m^2}{2(1+\alpha)} \left(1 - \frac{1}{2^{1+\alpha}}\right) \leq 2m^2 \end{aligned} \quad (3.15)$$

Hence  $\bar{I}_\alpha$  satisfies all the hypotheses of Mountain pass theorem and hence there exists a critical point  $u_\alpha$  of  $\bar{I}_\alpha$  such that

$$\bar{I}_\alpha(u_\alpha) \leq \sup_{t \in [0,1]} \bar{I}_\alpha(tu_0).$$

Now from (3.15) it follows that

$$\bar{I}_\alpha(u_\alpha) \leq 2m^2$$

and  $u_\alpha$  satisfies (3.4).

*Lemma 3.3.* Let  $f$  be in  $A'$ , then there exists  $\alpha_0 < 1$  such that for all  $\alpha_0 \leq \alpha < 1$ ,  $\Sigma_\alpha$  is non-empty and an  $u_\alpha \in \Sigma_\alpha$  satisfying

$$\frac{a_\alpha^2}{2} = I_\alpha(u_\alpha) = \inf_{u \in \partial B_\alpha} I_\alpha(u) = \inf_{u \in B_\alpha} I_\alpha(u) \quad (3.16)$$

and for all  $w$  in  $H_0^\alpha$ ,  $\|w\|_\alpha = 1$ ,

$$\int_0^1 f(r, a_\alpha w) w r^\alpha dr \leq a_\alpha. \quad (3.17)$$

*Proof.* Let  $u$  be in  $B_\alpha$ . Define  $\gamma \leq 1$  such that

$$\|u\|_\alpha^2 = \frac{1}{\gamma} \int_0^1 f(r, \gamma u) u r^\alpha dr. \quad (3.18)$$

Such a  $\gamma$  exists because  $f(r, t)/t$  is an increasing function and  $u$  is in  $B_\alpha$  and  $|f(r, t)| < \lambda_\alpha |t|$  for  $|t| < t_0$ ;  $\alpha_0 \leq \alpha < 1$ .

Define  $v = \gamma u$ , then

$$\begin{aligned} \|v\|_\alpha^2 &= \gamma^2 \|u\|_\alpha^2 = \int_0^1 f(r, \gamma u) (\gamma u) r^\alpha dr \\ &= \int_0^1 f(r, v) v r^\alpha dr. \end{aligned} \quad (3.19)$$

Hence  $v$  is in  $\partial B_\alpha$  and since  $\gamma \leq 1$ , and  $f \in A'$ , we have

$$I_\alpha(v) = I_\alpha(\gamma u) \leq I_\alpha(u).$$

this together with  $\partial B_\alpha \subset B_\alpha$  imply that

$$d_\alpha = \inf_{\partial B_\alpha} I_\alpha = \inf_{B_\alpha} I_\alpha. \quad (3.20)$$

Let  $u_n$  in  $\partial B_\alpha$  be a sequence such that  $u_n \geq 0$  and  $I_\alpha(u_n) \rightarrow d_\alpha$ . Such a sequence exists because for  $u$  in  $\partial B_\alpha$  implies  $|u|$  is in  $\partial B_\alpha$  and  $I_\alpha(u) = I_\alpha(|u|)$ .

We claim that  $\{\|u_n\|_\alpha\}$  is bounded. Let  $N$  be such that for all  $n \geq N$ ,

$$d_\alpha \leq I_\alpha(u_n) \leq d_\alpha + 1 \quad (3.21)$$

$$\begin{aligned} d_\alpha + 1 &\geq I_\alpha(u_n) = \frac{1}{2} \int_0^1 [f(r, u_n)u_n - 2F(r, u_n)]r^\alpha dr \\ &= \frac{1}{2} \int_0^1 [f(r, u_n)u_n - \beta F(r, u_n)]r^\alpha dr \\ &\quad + \left(\frac{\beta}{2} - 1\right) \int_0^1 F(r, u_n)r^\alpha dr. \end{aligned} \quad (3.22)$$

From (iv) of Definition (2.1), there exists a constant  $C$  depending only on  $f$  such that for all  $v$  in  $H_0^\alpha$ ,

$$\int_0^1 [f(r, v)v - \beta F(r, v)]r^\alpha dr \geq C. \quad (3.23)$$

From (3.22) and (3.23) there exists a constant  $C_1$  independent of  $n$  such that

$$\int_0^1 F(r, u_n)r^\alpha dr \leq C_1. \quad (3.24)$$

From (3.21) and (3.24) we have

$$\begin{aligned} \|u\|_\alpha^2 &= 2I_\alpha(u_n) + 2 \int_0^1 F(r, u_n)r^\alpha dr \\ &\leq 2(d_\alpha + 1) + 2C_1 \end{aligned}$$

and this proves the claim.

Let  $u_\alpha = \text{weak limit of } u_n$  and  $\alpha_0$  be as in Lemma (3.2). We claim that for  $\alpha_0 \leq \alpha < 1$ ,  $u_\alpha \in \Sigma_\alpha$  satisfying (3.16).

First we will show that  $u_\alpha$  is non-zero. Suppose  $u_\alpha \equiv 0$ , then from Lemma (3.1),  $u_n$  converges to 0 in  $C[0, 1]$ . Let  $N$  be an integer such that

$$u_n(r) < t_0 \quad \text{for all } n \geq N, r \in [0, 1]. \quad (3.25)$$

Then from (iii) of Definition (2.1) and the choice of  $\alpha_0$ ,

$$f(r, u_n(r)) < \lambda_\alpha u_n(r). \quad (3.26)$$

Since  $u_n \in \partial B_\alpha$ , we have from (3.26)

$$\begin{aligned} \|u_n\|_\alpha^2 &= \int_0^1 f(r, u_n)u_n r^\alpha dr \\ &< \lambda_\alpha \int_0^1 u_n(r)^2 r^\alpha dr \leq \|u_n\|_\alpha^2 \end{aligned}$$

which is a contradiction and hence  $u_\alpha \not\equiv 0$  and

$$\begin{aligned} I_\alpha(u_\alpha) &= \lim_{n \rightarrow \infty} I_\alpha(u_n) = d_\alpha \\ \|u_\alpha\|_\alpha^2 &\leq \lim_{n \rightarrow \infty} \|u_n\|_\alpha^2 = \int_0^1 f(r, u_\alpha) u_\alpha r^\alpha dr, \end{aligned} \quad (3.27)$$

$u_\alpha$  is in  $\partial B_\alpha$ . If not, then by (3.27) we can choose a  $\gamma < 1$  such that

$$\|u_\alpha\|^2 = \frac{1}{\gamma} \int_0^1 f(r, \gamma u_\alpha) u_\alpha r^\alpha dr.$$

Then  $\gamma u_\alpha$  is in  $\partial B_\alpha$  and

$$d_\alpha \leq I(\gamma u_\alpha) < I(u_\alpha) = d_\alpha.$$

This proves that  $u_\alpha$  is in  $\partial B_\alpha$ . Since  $u_\alpha$  is a minimizer and hence there exists a real number  $\rho$  such that for all  $\phi$  in  $H_0^\alpha$ ,

$$\begin{aligned} &\int_0^1 u'_\alpha(r) \phi'(r) r^\alpha dr - \int_0^1 f(r, u_\alpha) \phi r^\alpha dr \\ &= \rho \left\{ 2 \int_0^1 u'_\alpha(r) \phi'(r) r^\alpha dr - \int_0^1 f(r, u_\alpha) \phi r^\alpha dr - \int_0^1 \frac{\partial f}{\partial t}(r, u_\alpha) u_\alpha \phi r^\alpha dr \right\}. \end{aligned} \quad (3.28)$$

Putting  $\phi = u_\alpha$  in (3.28) and using the fact that  $u_\alpha \in \partial B_\alpha$ , we have

$$\rho \left\{ 2 \int_0^1 u'_\alpha(r)^2 r^\alpha dr - \int_0^1 f(r, u_\alpha) u_\alpha r^\alpha dr - \int_0^1 \frac{\partial f}{\partial t}(r, u_\alpha) u_\alpha(r)^2 r^\alpha dr \right\} = 0.$$

Since  $u_\alpha$  is in  $\partial B_\alpha$ , we have

$$\rho \int_0^1 \left[ \frac{f(r, u_\alpha)}{u_\alpha} - \frac{\partial f}{\partial t}(r, u_\alpha) \right] u_\alpha(r)^2 r^\alpha dr = 0.$$

Since  $f$  is in  $A'$ , and  $u$  is not zero, it implies that  $\rho = 0$ . Hence from (3.28) and by regularity of elliptic operator, it follows that  $u_\alpha$  is in  $\Sigma_\alpha$  and  $I_\alpha(u_\alpha) = d_\alpha$ . Since  $\Sigma_\alpha \subset \partial B_\alpha$ , we have  $a_\alpha^2/2 = \inf_{\Sigma_\alpha} I_\alpha = I_\alpha(u_\alpha) = d_\alpha$  and this proves (3.16). Let  $\|w\|_\alpha = 1$ . Choose  $\gamma > 0$  such that

$$1 = \frac{1}{\gamma} \int_0^1 f(r, \gamma w) w r^\alpha dr. \quad (3.29)$$

Then  $\gamma w$  is in  $\partial B_\alpha$ . Hence

$$\frac{a_\alpha^2}{2} \leq I_\alpha(\gamma w) \leq \frac{\gamma^2}{2} \|w\|_\alpha^2 = \frac{\gamma^2}{2}$$

implies  $a_\alpha \leq \gamma$ . Since  $f$  is in  $A'$ , we have

$$\frac{1}{a_\alpha} \int_0^1 f(r, a_\alpha w) w r^\alpha dr \leq \frac{1}{\gamma} \int_0^1 f(r, \gamma w) w r^\alpha dr = 1$$

i.e.

$$\int_0^1 f(r, a_\alpha w) w r^\alpha dr \leq a_\alpha$$

proving (3.17).

**Lemma 3.4.** *Let  $f$  be in  $A'$  and  $\alpha_0$  is as in Lemma (3.3). Then  $\{a_\alpha\}$  is bounded on  $[\alpha_0, 1)$ . Let  $a = \lim_{\alpha \rightarrow 1} a_\alpha$ . Then for all  $w \in H_0^1$  with  $\|w\|_1 = 1$ , we have*

$$\int_0^1 f(r, aw) w r dr \leq a. \quad (3.30)$$

*Proof.* From Lemma (3.2) and (3.3) we have  $l_\alpha = a_\alpha^2/2$  and  $l_\alpha \leq 2m^2$ . Hence  $\{a_\alpha\}$  is bounded on  $[\alpha_0, 1)$ . Let  $\alpha_n$  be a sequence such that  $a_{\alpha_n} \rightarrow a$  as  $\alpha_n \rightarrow 1$  and  $w$  be in  $E$  with  $\|w\|_1 = 1$ . Let  $v_n = w/\|w\|_{\alpha_n}$ . Then from (3.17) we have

$$\int_0^1 f(r, a_{\alpha_n} v_n) v_n r^\alpha dr \leq a_{\alpha_n}.$$

Letting  $\alpha_n \rightarrow 1$ ,  $v_n \rightarrow w$ ,  $a_{\alpha_n} \rightarrow a$ , we get

$$\int_0^1 f(r, aw) w r dr \leq a. \quad (3.31)$$

Since  $f$  is odd, and hence by Fatou's (3.31) holds for all  $w$  in  $H_0^1$ .

**Lemma 3.5.** *Let  $f$  be in  $A$ ,  $0 \leq \alpha < 1$ ,  $0 \leq \varepsilon \leq 1$ , and  $u$  in  $\Sigma_\alpha$ . Then we have*

$$u(r) = \frac{1-r^{1-\alpha}}{1-\alpha} \int_0^r f(t, u(t)) t^\alpha dt + \int_r^1 t^\alpha \left( \frac{1-t^{1-\alpha}}{1-\alpha} \right) f(t, u(t)) dt \quad (3.32)$$

$$\begin{aligned} \frac{1}{2} \varepsilon^{1+\alpha} u'(\varepsilon)^2 &= (1+\alpha) \int_0^\varepsilon F(r, u) r^\alpha dr + \int_0^\varepsilon \frac{\partial F}{\partial r}(r, u) r^{1+\alpha} dr \\ &\quad + \frac{1-\alpha}{2} \int_0^\varepsilon u'(r)^2 r^\alpha dr - \varepsilon^{1+\alpha} F(\varepsilon, u(\varepsilon)). \end{aligned} \quad (3.33)$$

*Proof.* If  $v(r)$  is the right hand side of (3.32), then by differentiating twice,  $v$  satisfies

$$\begin{aligned} L_\alpha v &= f(r, u) \\ v'(0) &= v(1) = 0. \end{aligned} \quad (3.34)$$

Hence by uniqueness,  $v = u$ . This proves (3.32).  $u$  is in  $\Sigma_\alpha$ , hence

$$(r^\alpha u')' = -f(r, u(r)) r^\alpha. \quad (3.35)$$

multiply (3.35) by  $ru'(r)$  and integrate from 0 to  $\varepsilon$  we get

$$\int_0^\varepsilon (r^\alpha u'(r))' u'(r) r dr = - \int_0^\varepsilon f(r, u) u' r^{1+\alpha} dr. \quad (3.36)$$

Since  $(dF/dr)(r, u(r)) = (\partial F/\partial r)(r, u(r)) + f(r, u(r))u'(r)$ , we have

$$\begin{aligned} \frac{1}{2} \varepsilon^{1+\alpha} u'(\varepsilon)^2 - \frac{(1-\alpha)}{2} \int_0^\varepsilon u'(r)^2 r^\alpha dr &= - \int_0^\varepsilon \frac{dF}{dr} r^{1+\alpha} dr + \int_0^\varepsilon \frac{\partial F}{\partial r} r^{1+\alpha} dr \\ &= -F(\varepsilon, u(\varepsilon)) \varepsilon^{1+\alpha} + (1+\alpha) \int_0^\varepsilon F(r, u) r^\alpha dr \\ &\quad + \int_0^\varepsilon \frac{\partial F}{\partial r} r^{1+\alpha} dr. \end{aligned}$$

Hence

$$\begin{aligned} \frac{1}{2} \varepsilon^{1+\alpha} u'(\varepsilon)^2 &= (1+\alpha) \int_0^\varepsilon F(r, u) r^\alpha dr + \int_0^\varepsilon \frac{\partial F}{\partial r} r^{1+\alpha} dr + \frac{1-\alpha}{2} \int_0^\varepsilon u'(r)^2 r^\alpha dr \\ &\quad - F(\varepsilon, u(\varepsilon)) \varepsilon^{1+\alpha}. \end{aligned}$$

This proves (3.33).

**Lemma 3.6.** *Let  $f$  be in  $A$ ,  $\alpha_n \rightarrow 1$ ,  $u_n$  is in  $\Sigma_{\alpha_n}$  and a constant  $M$  independent of  $n$  such that*

$$\begin{aligned} \text{(i)} \quad &\|u_n\|_{\alpha_n} \leq M \\ \text{(ii)} \quad &\lim_{n \rightarrow \infty} u'_n(1) = \eta \neq 0. \end{aligned} \quad (3.37)$$

*Then there exists a subsequence (still denoted by  $\alpha_n$ ) such that the weak limit  $u$  of  $u_n$  in  $H_0^1$  is a weak solution of (1.2). Furthermore*

$$\lim_{n \rightarrow \infty} \int_0^1 F(r, u_n) r^{\alpha_n} dr = \int_0^1 F(r, u) r dr. \quad (3.38)$$

*Proof.*  $\|u_n\|_1 \leq \|u_n\|_{\alpha_n} \leq M$ , hence by going to a subsequence the weak limit  $u$  of  $u_n$  in  $H_0^1$  exists. From (iii) of Lemma (3.1),  $u_n$  converges to  $u$  uniformly on compact subsets of  $(0, 1]$ . We claim  $u$  is not identically zero. For, if  $u \equiv 0$ , then, since  $u_n$  in  $\Sigma_{\alpha_n}$ , we have for  $0 < r \leq 1$ ,

$$r^{\alpha_n} u'_n(r) = u'_n(1) + \int_r^1 f(r, u_n) r^{\alpha_n} dr. \quad (3.39)$$

From (ii) of (3.37) and (3.39) and using  $u_n \rightarrow 0$  on  $[r, 1]$  uniformly

$$r \lim_{n \rightarrow \infty} u'_n(r) = \eta. \quad (3.40)$$

Hence by Fatou's lemma, and (3.40)

$$\infty = \eta^2 \int_0^1 \frac{r dr}{r^2} < \int_0^1 \lim_{n \rightarrow \infty} u'_n(r)^2 r^{\alpha_n} dr \leq \lim_{n \rightarrow \infty} \|u_n\|_{\alpha_n}^2 \leq M$$

which is a contradiction. Hence  $u \not\equiv 0$  and  $u$  satisfies

$$\begin{aligned} -(ru')' &= f(r, u)r \quad \text{in } (0, 1] \\ u(1) &= 0. \end{aligned} \quad (3.41)$$

Now by Fatous, we have

$$\int_0^1 f(r, u)ur \, dr \leq \liminf \int_0^1 f(r, u_n)u_n r^2 \, dr \leq M. \quad (3.42)$$

Hence

$$\int_0^1 f(r, u)r \, dr \leq \int_{u \leq 1} f(r, u)r \, dr + \int_{u > 1} f(r, u)ur \, dr < \infty. \quad (3.43)$$

For any  $0 < r \leq 1$ , integrating (3.41) from  $r$  to 1, we get

$$ru'(r) = u'(1) + \int_r^1 f(t, u)t \, dt. \quad (3.44)$$

(3.44) gives  $ru'(r)$  is monotone and hence limit  $r \rightarrow 0$  exists. We claim that

$$\lim_{r \rightarrow 0} ru'(r) = 0. \quad (3.45)$$

For, if  $\lim_{r \rightarrow 0} ru'(r) = C < 0$ , then there exists  $\varepsilon > 0$  such that  $-u'(r) \geq C/r$  for  $0 < r \leq \varepsilon$ . Hence

$$\infty = C^2 \int_0^\varepsilon \frac{r \, dr}{r^2} \leq \int_0^\varepsilon ru'(r)^2 \, dr < \infty.$$

Hence (3.45) is true. Using (3.44) and (3.45) we get

$$u'(1) = - \int_0^1 f(t, u)t \, dt. \quad (3.46)$$

Let  $\phi$  be in  $C^2[0, 1]$  with  $\phi(1) = 0$ . Multiply  $\phi'$  to (3.44) and integrate from 0 to 1, and using (3.46) we have

$$\begin{aligned} \int_0^1 u'(r)\phi'(r)r \, dr &= u'(1)(\phi(1) - \phi(0)) + \int_0^1 \phi'(r) \int_r^1 f(t, u)t \, dt \, dr \\ &= u'(1)(\phi(1) - \phi(0)) + \int_0^1 f(t, u)\phi(t)t \, dt \\ &\quad - \phi(0) \int_0^1 f(t, u)t \, dt \\ &= \int_0^1 f(t, u)\phi(t)t \, dt \end{aligned}$$

and hence  $u$  is a weak solution of (1.2).

From (3.33) and (3.37) we have

$$\lim_{n \rightarrow \infty} \left\{ (1 + \alpha_n) \int_0^1 F(r, u_n) r^{\alpha_n} dr + \int_0^1 \frac{\partial F}{\partial r} r^{1+\alpha_n} dr \right\} = \frac{1}{2} \eta^2 \quad (3.47)$$

Now multiply  $ru'(r)$  to (3.41) and integrate from  $r$  to 1, we have

$$\begin{aligned} -\frac{1}{2} r^2 u'(r)^2 + \frac{1}{2} u'(1)^2 &= - \int_r^1 \frac{dF}{dt} t^2 dt + \int_{r_1}^1 \frac{\partial F}{\partial t} t^2 dt \\ &= F(r, u(r)) r^2 + 2 \int_r^1 F(t, u) t + \int_r^1 \frac{\partial F}{\partial t} t^2 dt. \end{aligned} \quad (3.48)$$

Since  $ru'(r) \rightarrow 0$ ,  $\int_0^1 F(t, u) t dt < \infty$ ,  $\partial F / \partial r > 0$  in  $[0, \delta_0]$  and  $\int_{\delta_0}^1 (\partial F / \partial t) t^2 dt < \infty$ , we conclude that  $\lim_{r \rightarrow 0} F(r, u(r)) r^2$  exists and claim that

$$\lim_{r \rightarrow 0} F(r, u(r)) r^2 = 0. \quad (3.49)$$

If not, there exists a constant  $C > 0$  and  $\varepsilon > 0$  such that

$$F(r, u(r)) r^2 \geq C \quad \text{for all } 0 < r < \varepsilon.$$

Hence

$$\infty = \int_0^\varepsilon \frac{C}{r} dr \leq \int_0^\varepsilon F(r, u(r)) r dr < \infty$$

which is a contradiction.

Now using (3.49), (3.48) becomes

$$\frac{1}{2} u'(1)^2 = 2 \int_0^1 F(r, u) r dr + \int_0^1 \frac{\partial F}{\partial r} (r, u) r^2 dr. \quad (3.50)$$

Since  $u'(1) = \lim_{n \rightarrow \infty} u'_n(1)$ , and hence from (3.47) and (3.50) we have

$$\begin{aligned} &2 \int_0^1 F(r, u) r dr + \int_0^1 \frac{\partial F}{\partial r} (r, u) r^2 dr \\ &= \lim_{n \rightarrow \infty} \left\{ (1 + \alpha_n) \int_0^1 F(r, u_n) r^{\alpha_n} dr + \int_0^1 \frac{\partial F}{\partial r} (r, u_n) r^{1+\alpha_n} dr \right\}. \end{aligned} \quad (3.51)$$

By Fatou's and using (ii) of Definition (2.1) we have

$$\begin{aligned} &2 \int_0^1 F(r, u) r dr \leq \liminf_{n \rightarrow \infty} (1 + \alpha_n) \int_0^1 F(r, u_n) r^{\alpha_n} dr \\ &\int_0^1 \frac{\partial F}{\partial r} (r, u) r^2 dr \leq \liminf_{n \rightarrow \infty} \int_0^1 \frac{\partial F}{\partial r} (r, u_n) r^{\alpha_n+1} dr. \end{aligned} \quad (3.52)$$

By going to a subsequence, we conclude from (3.51) and (3.52) that

$$\lim_{n \rightarrow \infty} (1 + \alpha_n) \int_0^1 F(r, u_n) r^{\alpha_n} dr = 2 \int_0^1 F(r, u) r dr$$

and

$$\lim_{n \rightarrow \infty} \int_0^1 \frac{\partial F}{\partial r}(r, u_n) r^{\alpha_n+1} dr = \int_0^1 \frac{\partial F}{\partial r}(r, u) r^2 dr.$$

*Lemma 3.7.* Let  $f$  in  $A'$  be critical. Then

$$\frac{2}{b(0)} = \sup \left\{ c^2; \sup_{\|w\|_1 \leq 1} \int_0^1 f(r, cw) wr dr < \infty \right\} \quad (3.53)$$

*Proof.*  $f = h(r, t) \exp [b(r)t^2]$  for  $(r, t)$  in  $Q_{\delta_1}$ .

Let

$$C_0^2 = \sup \left\{ c^2; \sup_{\|w\|_1 \leq 1} \int_0^1 f(r, cw) wr dr < \infty \right\}$$

*Step 1.*  $C_0^2 \geq 2/b(0)$ .

If not, then choose  $\varepsilon > 0$ ,  $c > 0$  and a  $\delta < (\delta_1, \delta_0)$  such that

$$\frac{2}{b(0)} < c^2 < (c + \varepsilon)^2 < C_0^2. \quad (3.54)$$

For  $r_0 \in [0, \delta_1]$ , define

$$\begin{aligned} W_{r_0}(r) &= \frac{\log \frac{1}{r}}{\left( \log \frac{1}{r_0} \right)^{1/2}} \quad \text{for } r_0 \leq r \leq 1 \\ W_{r_0}(r) &= \left( \log \frac{1}{r_0} \right)^{1/2} \quad \text{for } 0 \leq r \leq r_0. \end{aligned} \quad (3.55)$$

Then  $\|w_{r_0}\|_1 = 1$ . Since  $(\partial f / \partial r)(r, t) \geq 0$  in  $Q_{\delta_0}$ , we have

$$h(0, t) \exp [b(0)t^2] \leq h(r, t) \exp [b(r)t^2] \quad \text{in } Q_{\delta_0}.$$

$(c + \varepsilon)^2 < C_0^2$  implies that there exists an absolute constant  $M$  depending only on  $b$  and  $f$  such that

$$\begin{aligned} M &\geq \int_0^1 f(r, (c + \varepsilon)w_{r_0}) w_{r_0} r dr \geq \int_0^\delta f(r, (c + \varepsilon)w_{r_0}) w_{r_0} r dr \\ &\geq \int_0^{r_0} f \left( 0, (c + \varepsilon) \left( \log \frac{1}{r_0} \right)^{1/2} \right) \left( \log \frac{1}{r_0} \right)^{1/2} r dr \\ &= \frac{1}{2} \left( \log \frac{1}{r_0} \right)^{1/2} h \left( 0, (c + \varepsilon) \left( \log \frac{1}{r_0} \right)^{1/2} \right) \exp \left[ b(0)(c + \varepsilon)^2 \log \frac{1}{r_0} \right] r_0^2 \\ &\geq \frac{\frac{1}{2} \left( \log \frac{1}{r_0} \right)^{1/2} h \left( 0, (c + \varepsilon) \left( \log \frac{1}{r_0} \right)^{1/2} \right) \exp \left[ \varepsilon^2 \left( \log \frac{1}{r_0} \right) \right]}{r_0^{(c^2 b(0) - 2)}} \rightarrow \infty \end{aligned}$$

as  $r_0 \rightarrow 0$ .



Hence  $C_0^2 \leq 2/b(0)$ .

Step 2.  $C_0^2 = 2/b(0)$ .

Suppose not, then choose  $\varepsilon > 0$ ,  $\delta > 0$  such that  $\delta \leq \min(\delta_1, \delta_0)$  and for all  $r$  in  $[0, \delta]$ ,

$$C_0^2 < (C_0 + \varepsilon)^2 < \frac{2 - \varepsilon}{b(r)}.$$

Let  $\|w\|_1 \leq 1$ , then

$$\int_0^1 f(r, (C_0 + \varepsilon)w)wr \, dr = \int_0^\delta + \int_\delta^1. \quad (3.56)$$

Since  $\|w\|_1 = 1$  implies from Lemma (3.1)

$$|w(r)| \leq \log \frac{1}{r},$$

hence there exists a constant  $M_1$  such that

$$\sup_{\|w\|_1 \leq 1} \int_\delta^1 f(r, (C_0 + \varepsilon)w)wr \, dr \leq M_1 \quad (3.57)$$

and

$$\begin{aligned} \int_0^\delta f(r, (C_0 + \varepsilon)w)wr \, dr &\leq \int_0^\delta h(r, (C_0 + \varepsilon)w) [\exp(C_0 + \varepsilon)^2 b(r) w^2] wr \, dr \\ &\leq \int_0^\delta h(r, (C_0 + \varepsilon)w) [\exp(2 - \varepsilon) w^2] wr \, dr \\ &\leq M_2 \int_0^\delta [\exp(2 - \varepsilon/2) w^2] r \, dr \\ &\leq M_2 \int_0^\delta r^{s/2-1} \, dr \leq M_3 \end{aligned} \quad (3.58)$$

where

$$M_2 = \sup_{(r,t) \in Q_\delta} h(r,t) t \exp -\frac{\varepsilon}{2} t^2.$$

This implies  $C_0 > (C_0 + \varepsilon)$  which is a contradiction. Hence  $C_0^2 = 2/b(0)$ .

*Lemma 3.8.* Let  $f$  in  $A'$  be critical and suppose there exists a  $t_0 > 0$  satisfying

$$\begin{aligned} \exp -t_0^2 &< \delta_1 \\ h\left(0, \left(\frac{2}{b(0)}\right)^{1/2}\right) t_0 &> 2\left(\frac{2}{b(0)}\right)^{1/2} \end{aligned} \quad (3.59)$$

Let  $a \geq 0$  such that

$$\sup_{\|w\|_1 \leq 1} \int_0^1 f(r, aw)wr \, dr \leq a \quad (3.60)$$

then  $a^2 < 2/b(0)$ .

*Proof.* From Lemma (3.7),  $a^2 \leq 2/b(0)$ . Suppose  $a^2 = 2/b(0)$ , then take  $r_0 = \exp -t_0^2$ ,  $w_{r_0}$  as in (3.55) and from (3.60) we have

$$\begin{aligned} \left(\frac{2}{b(0)}\right)^{1/2} &= a \geq \int_0^{r_0} f(r, aw_{r_0})w_{r_0}r \, dr \\ &\geq \int_0^{r_0} f(0, aw_{r_0})w_{r_0} \, dr \\ &= f(0, at_0)t_0 \frac{r_0^2}{2} \\ &= t_0 h(0, at_0) \exp 2 \left( \log \frac{1}{r_0} \right) \frac{r_0^2}{2} \\ &= \frac{1}{2} t_0 h \left( 0, \left( \frac{2}{b(0)} \right)^{1/2} t_0 \right) > \left( \frac{2}{b(0)} \right)^{1/2} \end{aligned}$$

which is a contradiction. Hence the result.

*Lemma 3.9.* For any  $\varepsilon > 0$ ,  $0 \leq \alpha < 1$ ,

$$\sup_{0 \leq r \leq 1} r^\varepsilon \left( \frac{1 - r^{1-\alpha}}{1 - \alpha} \right) \leq \frac{1}{\varepsilon}. \quad (3.61)$$

*Proof.* Let  $g(r) = r^\varepsilon(1 - r^{1-\alpha}/1 - \alpha)$ . Then  $g(0) = g(1) = 0$ . Let  $0 < r_0 < 1$  such that

$$g(r_0) = \sup_{0 \leq r \leq 1} g(r)$$

then

$$0 = g'(r_0) = \varepsilon r_0^{\varepsilon-1} \left( \frac{1 - r_0^{1-\alpha}}{1 - \alpha} \right) - r_0^{\varepsilon-\alpha}.$$

Hence

$$\frac{1 - r_0^{1-\alpha}}{1 - \alpha} = \frac{r_0^{1-\alpha}}{\varepsilon}.$$

Therefore

$$g(r) \leq g(r_0) \leq \frac{r_0^{1-\alpha+\varepsilon}}{\varepsilon} \leq \frac{1}{\varepsilon}.$$

*Lemma 3.10.* Let  $f$  in  $A'$  be critical, then

$$\inf_{B_1 \cup B_1^*} I_1 = \inf_{\partial(B_1 \cup B_1^*)} I_1 = \inf_{B_{01}} I_1 \quad (3.62)$$

*Proof.*  $u$  is in  $B_1 \cup B_1^*$  implies  $|u|$  also in  $B_1 \cup B_1^*$  and  $I_1(u) = I_1(|u|)$ . Let  $u \in B_1 \cup B_1^*$ ; choose a  $\gamma < 1$  such that

$$\|u\|_1^2 = \frac{1}{\gamma} \int_0^1 f(r, \gamma u) u r \, dr.$$

Then  $\gamma u$  is in  $\partial(B_1 \cup B_1^*)$  and  $I_1(\gamma u) \leq I_1(u)$ . Hence

$$\inf_{B_1 \cup B_1^*} I_1 = \inf_{\partial(B_1 \cup B_1^*)} I_1.$$

Now let  $u \geq 0$  is in  $\partial(B_1 \cup B_1^*)$ . Since  $f$  is critical, we have for any  $s > 1$

$$\int_0^1 f(r, su) u r \, dr < \infty.$$

Let  $v = su$ , then

$$\begin{aligned} \|v\|_1^2 &= s^2 \|u\|_1^2 = s^2 \int_0^1 f(r, u) u r \, dr \\ &= s \int_0^1 f\left(r, \frac{v}{s}\right) v r \, dr < \int_0^1 f(r, v) v r \, dr \end{aligned} \quad (3.63)$$

because  $s > 1$  and  $f(r, t)/t$  is increasing.

Choose an  $\varepsilon_0 > 0$  such that for any  $0 < \varepsilon < \varepsilon_0$

$$\|v\|_1^2 < \int_\varepsilon^1 f(r, v) v r \, dr \leq \int_0^1 f(r, v) v r \, dr \quad (3.64)$$

and define

$$v_\varepsilon = \begin{cases} v(\varepsilon) & \text{if } 0 \leq r \leq \varepsilon \\ v(r) & \text{if } \varepsilon \leq r \leq 1. \end{cases} \quad (3.65)$$

Then from (3.64)  $v_\varepsilon$  is in  $B_{01}$ .

Now we claim that  $I_1(v_\varepsilon) \rightarrow I_1(v)$  as  $\varepsilon \rightarrow 0$ .

*Case 1.* If  $v$  is in  $B_1$ , then  $\|v_\varepsilon\|_\infty \leq \|v\|_\infty$  and hence by dominated convergence theorem  $I_1(v_\varepsilon) \rightarrow I_1(v)$ .

*Case 2.* If  $v$  is in  $B_1^*$ , then  $v_\varepsilon \uparrow v$  and hence by Monotone convergence theorem,  $I_1(v_\varepsilon) \rightarrow I_1(v)$ . Hence

$$\inf_{B_{01}} I_1 \leq I_1(v_\varepsilon) \rightarrow I_1(v) \quad \text{as } \varepsilon \rightarrow 0. \quad (3.66)$$

$f$  is critical and is in  $A'$ , we have for  $1 \leq s \leq 2$

$$f(r, su)su - 2F(r, su) \leq 2f(r, 2u)u - 2F(r, 2u)$$

and is in  $L^1$ . Hence by dominated convergence theorem,

$$I_1(v) \rightarrow I_1(u) \quad \text{as } s \rightarrow 1. \quad (3.67)$$

Combining (3.66) and (3.67) we have

$$\inf_{B_{01}} I_1 \leq \inf_{\partial(B_1 \cup B_1^*)} I_1 \leq \inf_{B_{01}} I_1$$

and hence the result.

*Proof of theorem (2.1).* From Lemma (3.2), there exists  $\alpha_0 < 1$  such that  $\Sigma_\alpha$  is non-empty for  $\alpha_0 \leq \alpha < 1$  and  $\{l_\alpha\}$  is bounded by  $2m^2$  where  $m$  is given by (3.3). Let  $l = \lim_{\alpha \rightarrow 1} l_\alpha$ .

Let  $f$  satisfies (2.10). Let  $\eta > 0$ ,  $\gamma > 0$ ,  $\alpha_n \rightarrow 1$ ,  $u_n$  in  $\Sigma_{\alpha_n}$  such that

$$\begin{aligned} \text{(i)} \quad & l_{\alpha_n} \rightarrow l \quad \text{as } \alpha_n \rightarrow 1 \\ \text{(ii)} \quad & l_{\alpha_n} \leq \bar{I}_{\alpha_n}(u_n) < \left( l_{\alpha_n} + \frac{\eta}{2} \right). \\ \text{(iii)} \quad & (l_{\alpha_n} + \eta)b \leq \gamma < 1. \end{aligned} \quad (3.68)$$

We claim that

$$\lim_{\alpha_n \rightarrow 1} u'_n(1) \neq 0. \quad (3.69)$$

If not, then  $u'_n(1) \rightarrow 0$ . Since  $u_n \in \Sigma_{\alpha_n}$ , we have

$$u'_n(1) = - \int_0^1 f(r, u_n) r^{\alpha_n} dr \rightarrow 0 \quad \text{as } \alpha_n \rightarrow 1.$$

Since for any  $0 \leq r \leq 1$  we have

$$r^\alpha u'_n(r) = u'_n(1) + \int_r^1 f(t, u_n) t^{\alpha_n} dt$$

we have

$$\sup_{r \in [0,1]} |r^\alpha u'_n(r)| \rightarrow 0 \quad \text{as } \alpha_n \rightarrow 1.$$

This shows for any  $0 < r_0 \leq 1$ ,

$$\sup_{r_0 \leq r \leq 1} |u'_n(r)| \rightarrow 0 \quad \text{as } \alpha_n \rightarrow 1. \quad (3.70)$$

This in turn implies

$$\sup_{r_0 \leq r \leq 1} |u_n(r)| \leq \int_{r_0}^1 |u'_n(t)| dt \rightarrow 0 \quad \text{as } \alpha_n \rightarrow 1. \quad (3.71)$$

From (ii) of definition (2.1) and (3.33) we have

$$\begin{aligned} \frac{1}{2} \delta_0^2 u_n'(\delta_0)^2 &= (1 + \alpha_n) \int_0^{\delta_0} F(r, u_n) r^{\alpha_n} dr + \int_0^{\delta_0} \frac{\partial F}{\partial r}(r, u_n) r^{1 + \alpha_n} dr \\ &\quad + \frac{(1 - \alpha)}{2} \int_0^{\delta_0} u_n'(r)^2 r^{\alpha_n} dr - \delta_0^{1 + \alpha_n} F(\delta_0, u_n(\delta_0)) \\ &\geq (1 + \alpha) \int_0^{\delta_0} F(r, u_n) r^{\alpha_n} dr - \delta_0^{1 + \alpha_n} F(\delta_0, u_n(\delta_0)) \end{aligned} \quad (3.72)$$

Hence by (3.70) and (3.72) we have

$$\int_0^{\delta_0} F(r, u_n) r^{\alpha_n} dr \rightarrow 0 \quad \text{as } \alpha_n \rightarrow 1. \quad (3.73)$$

From (3.71) and by dominated convergence theorem

$$\int_{\delta_0}^1 F(r, u_n) r^{\alpha_n} dr \rightarrow 0 \quad \text{as } \alpha_n \rightarrow 1. \quad (3.74)$$

Combining (3.73) and (3.74) we have

$$\int_0^1 F(r, u_n) r^{\alpha_n} dr \rightarrow 0 \quad \text{as } \alpha_n \rightarrow 1. \quad (3.75)$$

Let  $N_0$  be such that for all  $n \geq N_0$ ,

$$\int_0^1 F(r, u_n) r^{\alpha_n} dr < \frac{\eta}{2}. \quad (3.76)$$

From (ii) and (iii) of (3.68) and (3.76)

$$\begin{aligned} \frac{1}{2} \|u_n\|_{\alpha_n}^2 &= \bar{I}_{\alpha_n}(u_n) + \int_0^1 F(r, u_n) r^{\alpha_n} dr \\ &< \left( l_{\alpha_n} + \frac{\eta}{2} \right) + \frac{\eta}{2} = (l_{\alpha_n} + \eta) \\ &\leq \frac{\gamma}{b}. \end{aligned}$$

Hence

$$\begin{aligned} |u_n(r)|^2 &\leq \|u\|_1^2 \log \frac{1}{r} \\ &< 2(l_{\alpha_n} + \eta) \log \frac{1}{r} \\ &\leq \frac{2\gamma}{b} \log \frac{1}{r}. \end{aligned} \quad (3.77)$$

From (3.32), (3.70) and (3.77) we have

$$\begin{aligned}
 u_n(0) &= \int_0^1 t^{\alpha_n} \left( \frac{1-t^{1-\alpha_n}}{1-\alpha_n} \right) f(t, u_n) dt \\
 &= \int_0^{\delta_1} t^{\alpha_n} \left( \frac{1-t^{1-\alpha_n}}{1-\alpha_n} \right) f(t, u_n) dt + \int_{\delta_1}^1 t^{\alpha_n} \left( \frac{1-t^{1-\alpha_n}}{1-\alpha_n} \right) f(t, u_n) dt \\
 &\leq M \int_0^{\delta_1} t^{\alpha_n} \left( \frac{1-t^{1-\alpha_n}}{1-\alpha_n} \right) \exp(bu_n^2) dt + M_1 \\
 &\leq M \int_0^{\delta_1} t^{\alpha_n} \left( \frac{1-t^{1-\alpha_n}}{1-\alpha_n} \right) \exp\left(2\gamma \log \frac{1}{t}\right) dt + M_1 \\
 &\leq M \int_0^{\delta_1} t^{\alpha_n-2\gamma} \left( \frac{1-t^{1-\alpha_n}}{1-\alpha_n} \right) dt + M_1
 \end{aligned} \tag{3.78}$$

Now choose  $\varepsilon > 0$  such that

$$\alpha_n > 2\gamma - 1 + \varepsilon \quad \text{for all } n, \text{ large.}$$

Then from (3.61) and (3.78) we have

$$\begin{aligned}
 u_n(0) &\leq M \int_0^{\delta_1} t^{\alpha_n-2\gamma-\varepsilon/2} t^{\varepsilon/2} \left( \frac{1-t^{1-\alpha_n}}{1-\alpha_n} \right) dt + M_1 \\
 &\leq \frac{2M}{\varepsilon} \frac{1}{\left( \alpha_n - 2\gamma + 1 - \frac{\varepsilon}{2} \right)} + M_2 \leq \frac{4M}{\varepsilon^2} + M_1.
 \end{aligned} \tag{3.79}$$

Hence

$$\|u_n\|_{\infty} = u_n(0) \leq \frac{4M}{\varepsilon^2} + M_1.$$

Since  $u_n$  is in  $\Sigma_{\alpha_n}$  and  $\{\|u_n\|_{\infty}\}$  is bounded and hence  $u_n$  converges strongly in  $C[0, 1]$  and in  $H_0^1$  to a function  $u$ . From (3.71)  $u_n(r) \rightarrow 0$  as  $\alpha_n \rightarrow \infty$  for every  $r \neq 0$ , we have  $u \equiv 0$  and hence  $u_n(0) \rightarrow 0$ . Now choose  $N$  large such that  $\|u_n\|_{\infty} \leq t_0$  for all  $n \geq N$ . From (iii) of Definition (2.1) we have

$$\begin{aligned}
 \lambda_{\alpha_n} \int_0^1 u_n \phi_{\alpha_n} r^{\alpha_n} dr &= - \int_0^1 (r^{\alpha_n} \phi'_{\alpha_n}) u_n dr \\
 &= - \int_0^1 (r^{\alpha_n} u'_n)' \phi_{\alpha_n} dr \\
 \lambda_{\alpha_n} \int_0^1 u_n \phi_{\alpha_n} r^{\alpha_n} dr &= - \int_0^1 (r^{\alpha_n} u'_n)' \phi_{\alpha_n} dr \\
 &= \int_0^1 f(r, u_n) \phi_{\alpha_n} r^{\alpha_n} dr \\
 &< \lambda_{\alpha_n} \int_0^1 u_n \phi_{\alpha_n} r^{\alpha_n} dr.
 \end{aligned}$$

and hence a contradiction. This proves the claim. Hence by going to a subsequence, we assume that

$$\begin{aligned} \lim_{\alpha_n \rightarrow 1} u'_n(1) &\neq 0 \\ l_{\alpha_n} &\leq \bar{I}_{\alpha_n}(u_n) < 2l_{\alpha_n} \leq 4m^2. \end{aligned} \quad (3.80)$$

Now

$$\begin{aligned} 4m^2 &\geq \frac{1}{2} \int_0^1 [f(r, u_n)u_n - 2F(r, u_n)]r^{\alpha_n} dr \\ &= \frac{1}{2} \int_0^1 [f(r, u_n)u_n - \beta F(r, u_n)]r^{\alpha_n} dr + \frac{\beta-2}{2} \int_0^1 F(r, u_n)r^{\alpha_n} dr \\ &\geq M_1 + \left(\frac{\beta-2}{2}\right) \int_0^1 F(r, u_n)r^{\alpha_n} dr \end{aligned}$$

where  $M_1$  is constant independent of  $n$ . Hence  $\exists M_2 > 0$  such that

$$\begin{aligned} \int_0^1 F(r, u_n)r^{\alpha_n} dr &\leq M_2 \\ \frac{1}{2} \|u_n\|_{\alpha_n}^2 &= \bar{I}_{\alpha_n}(u_n) + \int_0^1 F(r, u_n)r^{\alpha_n} dr \leq 4m^2 + M_2. \end{aligned} \quad (3.81)$$

Hence  $\{\|u_n\|_{\alpha_n}\}$  is uniformly bounded. Hence from (3.80) and Lemma (3.6),  $u_n$  converges weakly to a non-zero solution  $u$  of (1.2).

From condition (i) of Theorem (2.1), we have for every  $1 \leq p < \infty$ ,  $f(u) \in L^p(D)$  (see Moser [6]). Hence by regularity of elliptic operators,  $u \in W^{2,p}(D)$  and hence by Sobolev imbedding  $u$  is in  $C^1(\bar{D})'$  and hence in  $C^2(\bar{D})$ . This proves the result.

*Remark 3.1.* From the proof of Theorem (2.1) it follows that if  $m > 0$  is satisfying (2.11), then from Lemma (3.2)  $l_{\alpha} \leq 2m^2$  and hence  $l \leq 2m^2$ . Therefore if  $2m^2b < 1$  implies  $lb < 1$ . This proves the criterion (2.10).

*Proof of Theorem 2.2.* From Lemma (3.2) there exists  $\alpha_0 < 1$  such that  $\Sigma_{\alpha}$  is non-empty and  $\{a_{\alpha}\}$  is bounded for  $\alpha_0 \leq \alpha < 1$ . Lemma (3.3) gives (2.15).

*Case (1).* Let  $f$  be super critical and  $\overline{\lim_{\alpha \rightarrow 1} a_{\alpha}} = a \neq 0$ . Then from Lemma (3.4) we have

$$\sup_{\|w\|_1 \leq 1} \int_0^1 f(r, aw)wr dr \leq a.$$

contradicting the fact that  $f$  is super critical. Hence  $a = 0$ .

*Case 2.* If  $f$  is critical, let  $a = \overline{\lim_{\alpha \rightarrow 1} a_{\alpha}}$ . Then from (3.30) it follows that

$$\sup_{\|w\|_1 = 1} \int_0^1 f(r, aw)wr dr \leq a$$

and from Lemma (3.8),

$$\frac{a^2}{2} b(0) < 1. \quad (3.82)$$

Now choose an  $\varepsilon$  and  $\delta$  positive such that

$$\begin{aligned} \text{(i)} \quad & f(r, t) \leq M \exp[(b(0) + \varepsilon)t^2] \quad \text{for all } (r, t) \in Q_\delta. \\ \text{(ii)} \quad & \frac{a^2}{2}(b(0) + \varepsilon) < 1. \end{aligned} \quad (3.83)$$

Such a choice is possible because of (3.82) and the condition that  $f$  is critical.

Since  $a_\alpha^2/2 = l_\alpha$ , and hence  $f$  satisfies (2.10) of Theorem (2.1) with  $b$  replaced by  $(b(0) + \varepsilon)$  and hence there exists a sequence  $u_n$  in  $\Sigma_{\alpha_n}$  and a weak solution  $u$  of (1.2) such that

$$\begin{aligned} \text{(iii)} \quad & I_{\alpha_n}(u_n) \rightarrow \frac{a^2}{2} \quad \text{as } \alpha_n \rightarrow 1 \\ \text{(iv)} \quad & u_n \rightarrow u \quad \text{in } H_0^1. \\ \text{(v)} \quad & \lim_{n \rightarrow \infty} \int_0^1 F(r, u_n) r^\alpha dr = \int_0^1 F(r, u) r dr. \end{aligned} \quad (3.84)$$

In fact (iii) follows from Lemma (3.6). From weak lower semicontinuity of the norm we have

$$\|u\|_1^2 \leq \liminf_{\alpha_n \rightarrow 1} \|u_n\|_{\alpha_n}.$$

and hence from (iii) we have

$$I_1(u) \leq \liminf_{\alpha_n \rightarrow 1} I_{\alpha_n}(u_n) = \frac{a^2}{2}. \quad (3.85)$$

Let  $w$  be in  $B_{01}$ . Choose  $\gamma_\alpha$  such that

$$\|w\|_\alpha^2 = \frac{1}{\gamma_\alpha} \int_0^1 f(r, \gamma_\alpha w) w r^\alpha dr.$$

Such a  $\gamma_\alpha$  exists and  $\lim_{\alpha \rightarrow 1} \gamma_\alpha = \gamma_1$  exists and is  $\leq 1$  because  $w$  is in  $B_{01}$  and  $\gamma_\alpha w$  is in  $B_\alpha$ . Hence

$$\frac{a_\alpha^2}{2} \leq I(\gamma_\alpha w).$$

Taking the  $\overline{\lim}$  as  $\alpha \rightarrow 1$ , we get

$$\frac{a^2}{2} \leq I_1(\gamma w) \leq I_1(w)$$



This implies

$$\frac{a^2}{2} \leq \inf_{B_{01}} I_1. \quad (3.86)$$

From Lemma (3.10), (3.85) and (3.86) and using the fact that  $u$  is in  $B_1^*$ , we get

$$I_1(u) = \frac{a^2}{2} = \inf_{B_{01}} I_1$$

and  $a \neq 0$  because  $u \neq 0$ . This proves Theorem (2.2).

*Remark 3.2.* Suppose  $f(r, t) \leq 0$  for  $r \in [0, 1]$  and  $0 \leq t \leq t_0$  and satisfying all other hypothesis on  $f$ , then also the Theorems (2.1) and (2.2) are valid.

### Acknowledgements

I am extremely thankful to Dr P N Srikanth for having many helpful discussions and also providing me the useful references. I also thank Professor Mancini for many helpful discussions.

### References

- [1] Adimurthi, Infinite number of solutions of semilinear Dirichlet problem with critical growths in the unit disc in  $\mathbb{R}^2$  (in Preparation)
- [2] Adimurthi, Existence of positive solutions of the semilinear Dirichlet problem with critical growth for the  $n$ -Laplacian (preprint)
- [3] Atkinson F V and Peletier L A, Ground state and Dirichlet problems for  $-\Delta u = f(u)$  in  $\mathbb{R}^2$ , *Arch. Ration. Mech. Anal.* **96** (1986) 147–165
- [4] Brezis H and Nirenberg L, Positive solutions of nonlinear elliptic equations involving critical exponents, *Comm. Pure Appl. Math.* **36** (1983) 437–477
- [5] Carleson L and Chang A, On the existence of an extremal function for an inequality of J Moser, *Bull. Sc. Math. 2° Serie* **110** (1986) 113–127
- [6] Moser J, A sharp form of an inequality by N Trudinger, *Indiana Univ. Math. J.* **20** (1971) 1077–1092
- [7] Rabinowitz P H, Minimax methods in critical point theory with applications to differential equations, *Regional Conference Series in Mathematics* (AMS) Vol. 65
- [8] Zeev-Nehari, On a class of nonlinear second order differential equations, *Trans. AMS* **95** (1960) 101–123