arXiv:1108.2420v1 [quant-ph] 11 Aug 2011

Locally Accessible Information of Multisite Quantum Ensembles Violates Monogamy

Aditi Sen(De) and Ujjwal Sen

Harish-Chandra Research Institute, Chhatnag Road, Jhunsi, Allahabad 211 019, India

Locally accessible information is a useful information-theoretic physical quantity of an ensemble of multiparty quantum states. We find it has properties akin to quantum as well as classical correlations of single multiparty quantum states. It satisfies monotonicity under local quantum operations and classical communication. However we show that it does not follow monogamy, an important property usually satisfied by quantum correlations, and actually violates any such relation to the maximal extent. Violation is obtained even for locally indistinguishable, but globally orthogonal, multisite ensembles. The results assert that while single multiparty quantum states are monogamous with respect to their shared quantum correlations, ensembles of multiparty quantum states may not be so. The results have potential implications for quantum communication systems.

The science of quantum information [1] had its origins in thermodynamics [2] and foundations of quantum mechanics [3], and it has since been successfully applied to computation [4] and communication sciences [5], and has also found interesting links to many-body physics [6]. The field typically deals with a system of many different subsystems, and quantum correlations [7], in its multifarious versions, between the different subsystems form the backbone of these applications. The excitement in quantum information is even more because numerous laboratories around the globe can realize entanglement in a variety of different physical systems. While these different measures of quantum correlation have quite diverse motivations for their introduction and have otherwise dissimilar regions of utility, they do share certain intuitively satisfying criteria [8].

An important property that is expected to be satisfied by quantum correlation measures is monotonicity (more precisely, non-increasing), in some form, under local quantum operations and classical communication (LOCC).

Another prominent criterion that is usually expected from a measure of quantum correlation is the so-called "monogamy". Such a property is expected to be active in any measure of quantum correlation for any quantum state of a multiparty system, where each of the parties possess one of the subsystems. Monogamy requires that for any quantum state of a multiparty system, if two parties (i.e. the subsystems in possession of the respective parties) are highly correlated according to a certain measure of quantum correlation, then these parties would not have a substantial amount of that quantum correlation measure with any other third party (subsystem). See [9, 10].

None of these dual fundamental properties – monogamy and monotonicity – are expected to be satisfied by a classical correlation measure of a multiparty quantum state.

The concept of accessible information of a quantum ensemble is one that predates the usually accepted beginnings of the field of quantum information, and is defined as the maximal amount of classical information that one can obtain from a quantum ensemble by using arbitrary quantum measurements. The "Holevo bound" for accessible information obtained about 40 years earlier, provides us with an important piece of information: bit per qubit, i.e. the amount of classical information that can be incorporated in a two-dimensional quantum system (qubit) is one bit (binary digit) [11].

A related concept is that of locally accessible information and is the maximal amount of classical information that one can obtain from a multiparty quantum ensemble, when only *local* quantum measurement strategies (LOCC-based measurement strategies) are allowed – quantum measurements at all the subsystems and classical communication between them. The first result in this direction was very surprising: It was shown that there exists sets of orthogonal product states that are not locally distinguishable - "quantum nonlocality without entanglement" [12–14]. On the other hand, it turned out that any two orthogonal pure states of arbitrary dimensions and number of parties are locally distinguishable irrespective of its quantum correlation content [15]. Furthermore, an example was given of an ensemble of bipartite quantum states which is locally distinguishable, but on *reducing* its average quantum correlation content, the ensemble becomes locally indistinguishable – "more nonlocality with less entanglement" [16]. See also [17] in this regard.

Notwithstanding these counterintuitive properties of locally accessible information, the latter does have a certain amount of direct proportionality with quantum correlation, as was found in [18]. The Holevo bound states that using a quantum ensemble of a system of n qubits, it is not possible to send more than n bits of classical information. Ref. [18] demonstrates a local version of this result: Using a quantum ensemble of a bipartite system of n qubits, it is not possible to send more than $n - \overline{E}$ bits of classical information. Here \overline{E} denotes the average, over the ensemble, of an entanglement measure that satisfies certain basic postulates.

For an ensemble of multipartite quantum states, its locally accessible information is an important and useful information-theoretic physical quantity of the ensemble. In this paper, we find that locally accessible information possesses properties that are similar to those of quantum as well as of classical correlations of single multisite quantum states. In particular, it certainly satisfies the monotonicity postulate: locally accessible information of an ensemble of quantum states is nonincreasing under LOCC on the ensemble. However, we show here that the same physical quantity is polygamous, even for multiparty quantum ensembles that are LOCC indistinguishable. We also find locally distinguishable ensembles that violate any monogamy relation to the maximal extent.

For an ensemble of multiparty quantum states, shared between the N + 1 parties, A, B_1, B_2, \dots, B_N , the accessible information $I_{acc}^{LOCC,A:B_i}$ of the ensemble, when we have access to only the parties A and B_i (i = any one of $1, 2, \dots, N$), and when only LOCC is allowed between Aand B_i , monogamy would dictate that [9]

$$\sum_{i=1}^{N} I_{acc}^{LOCC,A:B_i} \le I_{acc}^{LOCC,A:B_1B_2...B_N}.$$
 (1)

We show that there exist locally indistinguishable (but globally orthogonal) ensembles that violate this relation.

Moreover, note that for an arbitrary ensemble shared between the N + 1 parties, we certainly have $I_{acc}^{LOCC,A:B_i} \leq \log_2 \Gamma$, where Γ denotes the cardinality of the ensemble [19]. Monogamy of locally accessible information would provide a bound on $\sum_{i=1}^{N} I_{acc}^{LOCC,A:B_i}$ that is strictly lower than $N \log_2 \Gamma$. Quite contrarily, we show that there exists ensembles of multiparty quantum states, for which the sum $\sum_{i=1}^{N} I_{acc}^{LOCC,A:B_i}$ violates monogamy to the maximal extent – it attains the value $N \log_2 \Gamma$.

The results have potential applications in quantum communication networks and in dealing with the information dynamics in quantum computation devices, especially in distributed quantum computing. Moreover, since we demonstrate the results for arbitrary N, the violation of monogamy obtained, persists for macroscopic systems. This, in particular, has potential implications for the way in which the quantum-to-classical transition is currently being viewed. Despite its importance in the field of secret quantum communication, quantum correlations of an ensemble of multiparty quantum states is not as yet a well-developed field, and the results obtained has potential of producing an axiomatic formalism for such a quantity (see [20] in this regard), just like the one existing for quantum correlations of a single quantum state.

Let us begin by briefly presenting a formal definition of accessible information and locally accessible information. Suppose that an observer, Alice, obtains the classical message i, and it is known that the message appears with probability p_i . She wants to send it to another observer, Bob. To this end, Alice encodes the information i in a quantum state ρ_i , and sends the quantum state to Bob. Bob receives the ensemble $\{p_i, \rho_i\}$, and wants to obtain as much information as possible about i. He performs a quantum measurement, that produces the result m, with probability q_m . Let the corresponding post-measurement ensemble be $\{p_{i|m}, \rho_{i|m}\}$. The classical information that is gathered about the index i by the quantum measurement, can be quantified by the mutual information between the message index i and the measurement outcome m [21]:

$$I(i:m) = H(\{p_i\}) - \sum_{m} q_m H(\{p_{i|m}\}).$$
 (2)

Here $H(\{r_{\alpha}\}) = -\sum_{\alpha} r_{\alpha} \log_2 r_{\alpha}$ is the Shannon entropy of the probability distribution $\{r_{\alpha}\}$. Note that the mutual information can be seen as the difference between the initial disorder and the (average) final disorder. Bob will be interested to obtain the maximal information, which is the maximum of I(i:m) over all measurement strategies. This quantity is called the accessible information,

$$I_{acc} = \max I(i:m), \tag{3}$$

where the maximization is over all measurement strategies. The Holevo bound provides a universal upper bound [11] (see also [18, 22]) on this quantity:

$$I_{acc}\left(\{p_i, \rho_i\}\right) \le \chi\left(\{p_i, \rho_i\}\right) \equiv S\left(\overline{\rho}\right) - \sum_i p_i S\left(\rho_i\right). \quad (4)$$

Here $\overline{\rho} = \sum_{i} p_i \rho_i$ is the average ensemble state, and

$$S(\varsigma) = -\operatorname{tr}(\varsigma \log_2 \varsigma) \tag{5}$$

is the von Neumann entropy of the quantum state ς . The bound is universal in the sense that it is valid for arbitrary ensembles. A weaker version of this result states that the accessible information for an ensemble of n qubit states is bounded above by n bits.

There also exists a universal lower bound on accessible information, and is given by [23, 24]

$$I_{acc}\left(\{p_i, \rho_i\}\right) \ge Q(\overline{\rho}) - \sum_i p_i Q(\rho_i),\tag{6}$$

where the "subentropy" Q is given by

6

$$Q(\varsigma) = -\sum_{k} \prod_{l \neq k} \frac{\lambda_k}{\lambda_k - \lambda_l} \lambda_k \log_2 \lambda_k, \tag{7}$$

with λ_k 's being the eigenvalues of the state ς , and where one must consider the limit as the eigenvalues become equal, in the degenerate case.

To arrive at the concept of locally accessible information, let us again suppose that Alice has a message i, and that again it is known that the message happens with probability p_i . But now, Alice encodes the message i in a *bipartite* quantum state ρ_i . She sends one part of the bipartite state to an observer called Bob₁ (\mathcal{B}_1), and the other part to an observer called Bob₂ (\mathcal{B}_2). The Bobs therefore receive the ensemble $\{p_i, \rho_i^{\mathcal{B}_1 \mathcal{B}_2}\}$, and their task is to gather as much information as possible about the index i, by using only LOCC. The maximal mutual information in this case is the locally accessible information,

$$I_{acc}^{LOCC} = \max I(i:m), \tag{8}$$

where the maximization is now over all LOCC-based measurement strategies. A universal upper bound on the locally accessible information is given by [18]

$$I_{acc}^{LOCC}\left(\{p_i, \varrho_i^{\mathcal{B}_1\mathcal{B}_2}\}\right) \leq \chi^{LOCC}\left(\{p_i, \varrho_i^{\mathcal{B}_1\mathcal{B}_2}\}\right)$$
$$\equiv S\left(\overline{\varrho}^{\mathcal{B}_1}\right) + S\left(\overline{\varrho}^{\mathcal{B}_2}\right) - \max_{k=1,2}\sum_i p_i S\left(\varrho_i^{\mathcal{B}_k}\right). (9)$$

Here $\overline{\varrho}^{\mathcal{B}_1} = \operatorname{tr}_{\mathcal{B}_2} \sum_i p_i \varrho_i^{\mathcal{B}_1 \mathcal{B}_2}$, and similarly for $\overline{\varrho}^{\mathcal{B}_2}$. Also, $\varrho_i^{\mathcal{B}_1} = \operatorname{tr}_{\mathcal{B}_2} \varrho_i^{\mathcal{B}_1 \mathcal{B}_2}$, and similarly for $\varrho_i^{\mathcal{B}_2}$. Again, a weaker version of this result is available, which states that the locally accessible information of a bipartite ensemble of n qubits is bounded above by $n - \overline{E}$, where $\overline{E} = \sum_i p_i E\left(\varrho_i^{\mathcal{B}_1 \mathcal{B}_2}\right)$ is the average entanglement E of the ensemble states. Here E is any measure of bipartite entanglement that satisfies $E(\zeta^{\mathcal{B}_1 \mathcal{B}_2}) \leq \max_{k=1,2} S\left(\zeta^{\mathcal{B}_k}\right)$ for all bipartite quantum states $\zeta^{\mathcal{B}_1 \mathcal{B}_2}$, where $\zeta^{\mathcal{B}_1} = \operatorname{tr}_{\mathcal{B}_2} \zeta^{\mathcal{B}_1 \mathcal{B}_2}$, and similarly for $\zeta^{\mathcal{B}_2}$.

Similarly, as in the case of accessible information with global operations, there also exist a universal lower bound on locally accessible information, and is given by [25, 26]

$$I_{acc}^{LOCC}\left(\{p_i, \varrho_i^{\mathcal{B}_1\mathcal{B}_2}\}\right) \ge \Lambda^{LOCC}\left(\{p_i, \varrho_i^{\mathcal{B}_1\mathcal{B}_2}\}\right)$$
$$\equiv Q_L\left(\overline{\varrho}^{\mathcal{B}_1\mathcal{B}_2}\right) - \sum_i p_i Q_L\left(\varrho_i^{\mathcal{B}_1\mathcal{B}_2}\right)(10)$$

Here $\overline{\varrho}^{\mathcal{B}_1 \mathcal{B}_2} = \sum_i p_i \varrho_i^{\mathcal{B}_1 \mathcal{B}_2}$, and the "local subentropy" Q_L is given by

$$Q_L(\zeta) = -d_{\mathcal{B}_1} d_{\mathcal{B}_2} \int d\alpha d\beta \langle \alpha | \langle \beta | \zeta | \alpha \rangle | \beta \rangle \log_2 \langle \alpha | \langle \beta | \sigma | \alpha \rangle | \beta \rangle,$$
(11)

for a bipartite state ζ of dimensions $d_{\mathcal{B}_1} \otimes d_{\mathcal{B}_2}$.

With these concepts in hand, we will now probe the status of monogamy for locally accessible information.

Case I. We begin by considering the following set of three-qubit Greenberger-Horne-Zeilinger (GHZ) states [27]:

$$|\psi_0^+\rangle_{AB_1B_2} = \frac{1}{\sqrt{2}}(|000\rangle + |111\rangle),$$

$$|\psi_0^-\rangle_{AB_1B_2} = \frac{1}{\sqrt{2}}(|000\rangle - |111\rangle).$$
(12)

Therefore, A and B_1 possess the quantum state

$$\frac{1}{2}(|00\rangle\langle 00| + |11\rangle\langle 11|), \tag{13}$$

irrespective of whether the three parties, A, B_1 , and B_2 share the state $|\psi_0^+\rangle$ or $|\psi_0^-\rangle$. Consequently,

$$I_{acc}^{LOCC,A:B_1} = 0. (14)$$

Similarly, $I_{acc}^{LOCC,A:B_2}$ is also vanishing, so that

$$I_{acc}^{LOCC,A:B_1} + I_{acc}^{LOCC,A:B_2} = 0$$
(15)

in this case. And, $I_{acc}^{LOCC,A:B_1B_2} = 1$ here, so that monogamy is satisfied in this case. The two GHZ states considered are distinguishable by LOCC between *all the three parties*. However, the complete orthogonal basis of GHZ states spanning the three-qubit Hilbert space, which is locally indistinguishable, also satisfies (actually saturates) the monogamy relation.

Case II. In complete contrast to the previous case, in this case study, we provide three concrete ensembles, each of which contains a different amount of average entanglement, such that each of them will violate any monogamy relation to the maximal extent. Each of the examples are three-qubit ensembles of two elements each. Let the three observers be again called A, B_1 , and B_2 . The maximal value that $I_{acc}^{LOCC,A:B_1}$ can attain for a two element ensemble is unity. So is the case for $I_{accc}^{LOCC,A:B_2}$. The first ensemble (\mathcal{E}_1) consists of the GHZ states

$$|\psi_0^+\rangle_{AB_1B_2} = \frac{1}{\sqrt{2}}(|000\rangle + |111\rangle),$$

$$|\psi_3^+\rangle_{AB_1B_2} = \frac{1}{\sqrt{2}}(|011\rangle - |100\rangle).$$
(16)

Forgetting about B_2 , the ensemble consists of the states

$$\rho_{0+}^{AB_1} = \frac{1}{2} (|00\rangle \langle 00| + |11\rangle \langle 11|),
\rho_{1+}^{AB_1} = \frac{1}{2} (|01\rangle \langle 01| + |10\rangle \langle 10|).$$
(17)

This ensemble can be exactly distinguished by LOCC between A and B_1 , by measurement in the computational basis at A and B_1 and communication of the results (say, by a phone call), so that

$$I_{acc}^{LOCC,A:B_1}(\mathcal{E}_1) = 1.$$
(18)

The ensemble \mathcal{E}_1 is invariant under a swap operation between B_1 and B_2 , so that we also have

$$I_{acc}^{LOCC,A:B_2}(\mathcal{E}_1) = 1.$$
(19)

Consequently, we have

$$I_{acc}^{LOCC,A:B_1}(\mathcal{E}_1) + I_{acc}^{LOCC,A:B_2}(\mathcal{E}_1) = 2, \qquad (20)$$

and 2 is the maximal value that the sum on the left-handside can attain (for an arbitrary two-element ensemble), because the individual algebraic maxima are unity. This therefore is a violation of any monogamy relation that one can envisage for locally accessible information. The states in the ensemble \mathcal{E}_1 are genuinely multiparty entangled [28]. However, the violation of monogamy is not related to this fact, as borne out by the next two ensembles. Let us therefore consider the ensemble \mathcal{E}_2 that consists of the states

$$|\psi^{+}\rangle_{AB_{1}B_{2}} = |0\rangle \frac{1}{\sqrt{2}} (|00\rangle + |11\rangle),$$

 $|\psi^{-}\rangle_{AB_{1}B_{2}} = |1\rangle \frac{1}{\sqrt{2}} (|00\rangle - |11\rangle).$ (21)

Again we have

$$I_{acc}^{LOCC,A:B_1}(\mathcal{E}_2) + I_{acc}^{LOCC,A:B_2}(\mathcal{E}_2) = 2, \qquad (22)$$

and again 2 is the maximal value that the sum on the lefthand-side can attain, because the individual algebraic maxima are unity. The states of this ensemble are still entangled, although not genuinely multisite entangled. The third and last ensemble (\mathcal{E}_3) consists of the states $|000\rangle$ and $|111\rangle$, which once more violates any monogamy relation that one can write down, and this ensemble is devoid of any quantum correlations in its element states. Note that just like in **Case I**, all the ensembles are distinguishable by LOCC between all the three parties.

Case III. The violations in **Case II** are all obtained for tripartite ensembles. However, this is not a necessary restriction, and ensembles of an arbitrary number of parties can be shown to maximally violate the corresponding monogamy, as seen in the following example. We present the example by again using a two-element ensemble, where each element is genuinely multiparty entangled [28]. Let us therefore consider the ensemble (\mathcal{E}_{cat}) containing the GHZ states (also called cat states)

$$\begin{aligned} |\psi\rangle_{AB_{1}B_{2}...B_{N}}^{cat} &= \frac{1}{\sqrt{2}} (|000...0\rangle + |111...1\rangle), \\ |\phi\rangle_{AB_{1}B_{2}...B_{N}}^{cat} &= \mathbb{I}_{A} \otimes \sigma_{B_{1}}^{x} \otimes ... \otimes \sigma_{B_{N}}^{x} |\psi\rangle_{AB_{1}B_{2}...B_{N}}^{cat}, \end{aligned}$$
(23)

where σ^x is the Pauli spin-flip operator. Tracing out all the parties except A and B_1 , we obtain the ensemble

$$\rho_{\psi^{cat}}^{AB_{1}} = \frac{1}{2} (|00\rangle\langle00| + |11\rangle\langle11|),
\rho_{\phi^{cat}}^{AB_{1}} = \frac{1}{2} (|01\rangle\langle01| + |10\rangle\langle10|).$$
(24)

This ensemble can be exactly distinguished, and so we have

$$I_{acc}^{LOCC,A:B_1}(\mathcal{E}_{cat}) = 1.$$
⁽²⁵⁾

However, the ensemble (\mathcal{E}_{cat}) is invariant with respect to a swap between any two of the B_i s (i = 1, 2, ..., N), and so we have

$$I_{acc}^{LOCC,A:B_i}(\mathcal{E}_{cat}) = 1, \quad \forall i = 1, 2, \dots, N.$$
 (26)

However, each of the $I_{acc}^{LOCC,A:B_i}(\mathcal{E}_{cat})$ can reach a maximum of unity, as \mathcal{E}_{cat} is a two-element ensemble. Therefore, for any monogamy relation to be nontrivial, we must have the sum $\sum_{i=1}^{N} I_{acc}^{LOCC,A:B_i}$ strictly less than N. However, the sum is actually equal to N for the ensemble \mathcal{E}_{cat} .

Case IV. The ensembles that we have considered until now are all qubit ensembles, and have two elements in them. Neither of these conditions are necessary. For example, the ensemble \mathcal{E}_T , consisting of the three threequtrit quantum states (shared between A, B_1 , and B_2 respectively),

$$|000\rangle + |111\rangle,$$

$$|011\rangle + |122\rangle,$$

$$|100\rangle + |200\rangle,$$

(27)

also violates any monogamy relation to the maximal extent. Also, the ensembles that have been considered above do not form complete bases of the corresponding multiparty Hilbert space. Again, this condition is not a necessity, as the ensemble \mathcal{E}_P formed by the eight three-qubit quantum states $|ijk\rangle$ (i, j, k = 0, 1), form a complete basis, and also violates the monogamy relation for I_{acc}^{LOCC} . It is also not necessary to consider orthogonal ensembles to violate monogamy, as has been done until now. This can be seen by considering the ensemble consisting of the states $|000\rangle$ and $|\vec{n}\vec{n}\vec{n}\rangle$, where $|\vec{n}\rangle$ is a qubit state slightly different from $|1\rangle$.

Case V. We now consider the ensemble \mathcal{E}_{shifts} of the following four three-qubit, (globally) orthogonal, quantum states

$$|01+\rangle, |1+0\rangle, |+01\rangle, |---\rangle, (28)$$

where $|\pm\rangle = \frac{1}{\sqrt{2}}(|0\rangle \pm |1\rangle)$. This set was discovered in Ref. [13], and it was shown that the set forms an unextendible product basis, in the sense that there are no product states in the orthogonal complement of the subspace spanned the elements in \mathcal{E}_{shifts} . It was shown there that this ensemble cannot be distinguished by LOCC between the three parties. It was thereafter connected to the phenomenon of bound entanglement [29]. We will now show that the ensemble \mathcal{E}_{shifts} violates monogamy. Let the three parties possessing the states in \mathcal{E}_{shifts} be called A, B_1 , and B_2 respectively. Leaving out B_2 , the ensemble consists of the states

$$|01\rangle, |1+\rangle, |+0\rangle, |--\rangle.$$
 (29)

Let us try to find the locally accessible information for this ensemble. Let us begin by noting that the universal upper and lower bounds on locally accessible information imply that

$$1 - \frac{1}{2}\log_2 \mathbf{e} \le I_{acc}^{LOCC,A:B_1}(\mathcal{E}_{shifts}) \le 2, \qquad (30)$$

that is

$$0.27865 \le I_{acc}^{LOCC,A:B_1}(\mathcal{E}_{shifts}) \le 2.$$
(31)

Here and hereafter, all numerical values are rounded off to the fifth decimal place. The upper bound does not help us in violating any monogamy relation. And the lower bound is too weak for our purposes. We now find a much stronger lower bound. This is obtained by using the following LOCC measurement strategy, which includes a single bit of classical communication from A to B_1 . Suppose that A measures in the basis $\mathbb{Z} = \{|0\rangle, |1\rangle\}$, and sends the result to B_1 over a classical channel. If the result is $|0\rangle$, the observer in possession of B_1 measures in the basis \mathbb{Z} , and otherwise he measures in the basis $\mathbb{X} = \{|+\rangle, |-\rangle\}$. The mutual information, between the ensemble index and the measurement index, that is obtained by following this LOCC measurement strategy, can be shown to be

$$I_{M}^{LOCC,A:B_{1}}(\mathcal{E}_{shifts}) \geq \frac{13}{4} - \frac{1}{8} [3 \log_{2} 3 + 5 \log_{2} 5] \\\approx 1.20443.$$
(32)

The bound is much better than the universal lower bound, and indeed will help us to violate monogamy. Now

$$I_{acc}^{LOCC,A:B_1}(\mathcal{E}_{shifts}) \ge I_M^{LOCC,A:B_1}(\mathcal{E}_{shifts}) \ge 1.20443.$$
(33)

Also, the ensemble obtained from \mathcal{E}_{shifts} by leaving out B_1 , is the same as that obtained by leaving out B_2 , up to a swap operation, and we know that locally accessible information is invariant under the swap operation. Consequently, we have

$$I_{acc}^{LOCC,A:B_2}(\mathcal{E}_{shifts}) \ge 1.20443, \tag{34}$$

so that

$$I_{acc}^{LOCC,A:B_1}(\mathcal{E}_{shifts}) + I_{acc}^{LOCC,A:B_2}(\mathcal{E}_{shifts}) \ge 2.40887.$$
(35)

However, since there are four elements in the ensemble \mathcal{E}_{shifts} , we have

$$I_{acc}^{LOCC,A:B_1B_2}(\mathcal{E}_{shifts}) \le 2.$$
(36)

Consequently, the monogamy relation is violated for the locally indistinguishable ensemble \mathcal{E}_{shifts} by more than 20%.

In conclusion, we have shown that locally accessible information of multisite quantum ensembles can violate monogamy, even maximally. Violation can appear even for locally indistinguishable, but globally orthogonal, multiparty quantum ensembles. This is despite the fact that this physically important quantity does satisfy monotonicity under local operations and classical communication.

None of the dual fundamental properties of monogamy and monotonicity are expected to be satisfied by a classical correlation measure of a multiparty system, quantum or classical. Indeed, just like a single ball can be either green or blue in color, a set \mathcal{S} of ten (or twenty) balls can be either all green or all blue. [Similarly, the spin states of a set of ten spin-1/2 particles can be either all up, in the z-direction, or all down.] The marginals of Sconsisting of any two balls is again either both green or both blue, irrespective of the number of balls in S – a clear violation of monogamy. Also, monotonicity is violated by any classical correlation measure, as can be seen in the following scenario. Suppose that two white balls are sent to two cities, so that initially there are no correlations in this bipartite system. The receivers of the balls are then instructed to color them to a some single color and to choose that color from among green, blue, and red, and to fix the color by phone call between the parties. So finally, classical correlation is present, whatever be its value (that depends on the exact measure (and its normalization) used), in the bipartite system.

While classical correlation measures are not expected to satisfy these dual properties, quantum correlations are. Locally accessible information, therefore, contains elements of both the worlds.

On the application front, the results may have implications for quantum communication networks, where classical information is transferred by using quantum means. Another potential candidate for application is distributed quantum computing, where efficient transfer of information is vital for a robust and competent performance of the system. Yet another possible region of application is in secure information transfer, where quantum correlations of quantum ensembles is an important, though not very well-understood, physical quantity, and the results obtained may provide inputs towards an axiomatic formalism of this quantity.

On the fundamental side, we note that the violations are obtained for systems with an arbitrary number of subsystems, and so are valid even for macroscopic systems. This can have implications for researches in the quantumto-classical transition: Macroscopic systems, whether classical or quantum, have inherent physical quantities that violate monogamy, but satisfy monotonicity. Lastly, the results may be important to understand the counterintuitive properties obtained in Refs. [12–16], and may indicate that these nonintuitive properties are a product of the violation of monogamy of locally accessible information.

- M.A. Nielsen and I.L. Chuang, *Quantum Computation and Quantum Information* (Cambridge University Press, Cambridge, 2000).
- [2] See C.H. Bennett, Internat. J. Theoret. Phys. 21, 905 (1982);
 S. Lloyd, Phys. Rev. A 56, 3374 (1997);
 P. Horodecki, R. Horodecki, and M. Horodecki, Acta Phys. Slov. 48, 141 (1998);
 L Henderson and V Vedral, J. Phys. A: Math. Gen. 34, 6899 (2001);
 H. Ollivier

and W.H. Zurek, Phys. Rev. Lett. 88, 017901 (2001);
R. Horodecki, M. Horodecki, and P. Horodecki, Phys.
Rev. A 63, 022310 (2001); V. Vedral and E. Kashefi,
Phys. Rev. Lett. 89, 037903 (2002); J. Oppenheim, M.
Horodecki, P. Horodecki, and R. Horodecki, Phys. Rev.
Lett. 89, 180402 (2002); M. Horodecki, K. Horodecki,
P. Horodecki, R. Horodecki, J. Oppenheim, A. Sen(De),
and U. Sen, Phys. Rev. Lett. 90, 100402 (2003); M.

Horodecki, P. Horodecki, R. Horodecki, J. Oppenheim,
A. Sen(De), U. Sen, and B. Synak-Radtke, Phys. Rev. A
71, 062307 (2005); F.G.S.L. Brandao and M.B. Plenio,
Nat. Phys. 4, 873 (2008); K. Maruyama, F. Nori, and V.
Vedral, Rev. Mod. Phys. 81, 1 (2009); F.G.S.L. Brandao and M.B. Plenio, Commun. Math. Phys. 295, 829 (2010); L. del Rio, J. Aberg, R. Renner, O. Dahlsten, and V. Vedral, Nature 474, 61 (2011), and references therein.

- [3] A. Einstein, B. Podolsky, and N. Rosen, Phys. Rev. 47, 777 (1935); J.S. Bell, *Speakable and Unspeakable in Quantum Mechanics* (Cambridge University Press, Cambridge, 2004), and references thereto.
- [4] A. Ekert and R. Jozsa, Rev. Mod. Phys. 68, 733 (1996), and references therein.
- [5] C.H. Bennett and G. Brassard, in Proceedings of the International Conference on Computers, Systems and Signal Processing, Bangalore, India (IEEE, NY, 1984); A.K. Ekert, Phys. Rev. Lett. (1991); C.H. Bennett and S.J. Wiesner, *ibid.* **69**, 2881 (1992); C.H. Bennett, G. Brassard, C. Crépeau, R. Josza, A. Peres, and W.K. Wootters, Phys. Rev. Lett. *ibid.*, 1895 (1993). For a recent review, see e.g. A. Sen(De) and U. Sen, Physics News **40**, 17 (2010) (arXiv:1105.2412 [quant-ph]).
- [6] M. Lewenstein, A. Sanpera, V. Ahufinger, B. Damski, A. Sen(De), and U. Sen, Adv. Phys. 56, 243 (2007); L. Amico, R. Fazio, A. Osterloh, and V. Vedral, Rev. Mod. Phys. 80, 517 (2008).
- [7] R. Horodecki, P. Horodecki, M. Horodecki, and K. Horodecki, Rev. Mod. Phys. 81, 865 (2009).
- [8] See e.g. M. Horodecki, P. Horodecki, and R. Horodecki, Phys. Rev. Lett. 84, 2014 (2000), and references therein.
- [9] V. Coffman, J. Kundu, and W.K. Wootters, Phys. Rev. A 61, 052306 (2000).
- [10] See also M. Koashi and A. Winter, Phys. Rev. A 69, 022309 (2004); G. Adesso, A. Serafini, and F. Illuminati, Phys. Rev. A 73, 032345 (2006); T.J. Osborne and F. Verstraete, Phys. Rev. Lett. 96, 220503 (2006); T. Hiroshima, G. Adesso, and F. Illuminati, Phys. Rev. Lett. 98, 050503 (2007); M. Seevinck, Phys. Rev. A 76, 012106 (2007); S. Lee and J. Park, Phys. Rev. A 79, 054309 (2009); A. Kay, D. Kaszlikowski, and R. Ramanathan, Phys. Rev. Lett. 103, 050501 (2009); M. Hayashi and L. Chen, Phys. Rev. A 84, 012325 (2011), and references therein.
- [11] J.P. Gordon, in Proc. Int. School Phys. "Enrico Fermi, Course XXXI", ed. P.A. Miles, pp. 156 (Academic Press, NY 1964); L.B. Levitin, in Proc. VI National Conf. Inf. Theory, Tashkent, pp. 111 (1969); A.S. Holevo, Probl. Pereda. Inf. 9, 3 1973 [Probl. Inf. Transm. 9, 110 (1973)]; H.P. Yuen and M. Ozawa, Phys. Rev. Lett. 70, 363 (1993); H.P. Yuen in Quantum Communication, Computing, and Measurement, ed. O. Hirota et al. (Plenum, NY 1997)).
- [12] C.H. Bennett, D.P. DiVincenzo, C.A. Fuchs, T. Mor, E. Rains, P.W. Shor, J.A. Smolin, and W.K. Wootters, Phys. Rev. A 59, 1070 (1999).

- [13] C.H. Bennett, D.P. DiVincenzo, T. Mor, P.W. Shor, J.A. Smolin, and B.M. Terhal, Phys. Rev. Lett. 82, 5385 (1999).
- [14] D.P. DiVincenzo, T. Mor, P.W. Shor, J.A. Smolin, and B.M. Terhal, Comm. Math. Phys. 238, 379 (2003).
- [15] J. Walgate, A.J. Short, L. Hardy, and V. Vedral, Phys. Rev. Lett. 85, 4972 (2000); S. Virmani, M.F. Sacchi, M.B. Plenio, and D. Markham, Phys. Lett. A 288, 62 (2001); Y.-X. Chen and D. Yang, Phys. Rev. A 64, 064303 (2001); 65, 022320 (2002); J. Walgate and L. Hardy, Phys. Rev. Lett. 89, 147901 (2002).
- [16] M. Horodecki, A. Sen(De), U. Sen, and K. Horodecki, Phys. Rev. Lett. **90**, 047902 (2003).
- [17] A. Peres and W.K. Wootters, Phys. Rev. Lett. 66, 1119 (1991); W.K. Wootters, arXiv:quant-ph/0506149, and references therein.
- [18] P. Badziąg, M. Horodecki, A. Sen(De), and U. Sen, Phys. Rev. Lett. **91**, 117901 (2003).
- [19] Throughout the paper, we calculate the all the quantities on amounts of information transfer in bits (binary digits), and this requires the use of logarithm with base 2 in the definition mutual information.
- [20] A quantum correlation measure for an ensemble of quantum states (including single party ones) was proposed in M. Horodecki, A. Sen(De), U. Sen, Phys. Rev. A **75**, 062329 (2007). See also M.-Y. Ye, Y.-K. Bai, X.-M. Lin, and Z.D. Wang, Phys. Rev. A **81**, 014303 (2010) in this regard. See C.A. Fuchs and A. Peres, arXiv:quant-ph/9512023; B.-G. Englert, Phys. Rev. Lett. **77**, 2154 (1996) for an earlier discussion concerning single-party ensembles.
- [21] T.M. Cover and J.A. Thomas, *Elements of Information Theory* (Wiley, New York, 1991).
- [22] B. Schumacher, M. Westmoreland, and W.K. Wootters, Phys. Rev. Lett. **76**, 3452 (1996); M. Horodecki, J. Oppenheim, A. Sen(De), and U. Sen, *ibid.* **93**, 170503 (2004).
- [23] R. Jozsa, D. Robb, and W.K. Wootters, Phys. Rev. A 49, 668 (1994).
- [24] See also S.R. Nichols and W.K. Wootters, QIC 3, 1 (2003); F. Mintert and K. Życzkowski, Phys. Rev. A 69, 022317 (2004).
- [25] A. Sen(De), U. Sen, and M. Lewenstein, Phys. Rev. A 74, 052332 (2006).
- [26] W. Matthews, S. Wehner and A. Winter, Commun. Math. Phys. 291, 813 (2009).
- [27] D.M. Greenberger, M.A. Horne, and A. Zeilinger, in *Bells Theorem, Quantum Theory, and Conceptions of the Universe*, ed. M. Kafatos, (Kluwer, Dordrecht, 1989).
- [28] A pure multiparty quantum state is said to be genuinely multiparty entangled if the state is bipartite entangled across any partition of the set of observers into two groups.
- [29] P. Horodecki, Phys. Lett. A 232, 333 (1997); M. Horodecki, P. Horodecki, and R. Horodecki, Phys. Rev. Lett. 80, 5239 (1998).