

Changing epidemiology of hepatitis A virus in Indian children

Vidya Arankalle¹
 Monjori Mitra²
 Sheila Bhawe³
 Apurba Ghosh²
 Sundaram Balasubramanian⁴
 Suparna Chatterjee⁵
 Jaydeep Choudhury⁶
 Amarjeet Chitkara⁷
 Ganesh Kadhe⁸
 Amey Mane⁸
 Sucheta Roy⁸

¹Department of Virology, National Institute of Virology, Pashan, Pune, Maharashtra, India; ²Department of Pediatrics, Institute of Child Health, Kolkata, West Bengal, India; ³Department of Pediatrics, KEM Hospital and Research Center, Pune, Maharashtra, India; ⁴Department of Pediatrics, Kanchi Kamkodi Child Trust Hospital (KKCTH), Nungambakkam, Chennai, Tamil Nadu, India; ⁵Department of Pharmacology, Institute of Post Graduate Medical Education and Research (IPGMER) and Seth Sukhlal Karnani Memorial (SSKM) Hospital, Kolkata, West Bengal, India; ⁶Department of Pediatrics, Sri Aurobindo Seva Kendra (EEDF Hospital), Kolkata, West Bengal, India; ⁷Department of Pediatrics, Sarvodaya Childcare, Pitampura, Delhi, India; ⁸Medical Affairs Department, Wockhardt Ltd, Mumbai, Maharashtra, India

Correspondence: Monjori Mitra
 Department of Pediatrics, Institute of Child Health, 11, Dr Biresh Guha Street, Kolkata, West Bengal, India
 Email monjorim@medclinsearch.com

Abstract: Previous studies from India have observed an increased incidence of hepatitis A virus (HAV) infection in the adult and adolescent population compared with children, indicating a shift in epidemiology of HAV. However, no HAV seroprevalence study has been conducted in India over the past decade. This prospective, multicenter, cross-sectional study was conducted in 928 children (aged 18 months to 10 years), to estimate the age-related seroprevalence of HAV across different regions of India. The present study also evaluated the impact of various factors such as age, socioeconomic class (SEC), education, source of drinking water, and excreta disposal on HAV seroprevalence. Overall, 348 (37.5%) children were seropositive for anti-HAV antibodies. Seroprevalence of HAV in the 6- to 10-year age group (50.3%) was higher ($P=0.000$) than in the 18-month to 6-year age group (30.3%). SEC and educational status of the parents were significantly associated with HAV seropositivity ($P=0.000$ for both). An increase in seroprevalence was observed with an increase in age in all SECs ($P<0.000-0.001$ for all) except upper SEC ($P=0.124$). Fewer subjects using a private toilet within the house (33.1%) were seropositive than subjects using an open field (75%) for excreta disposal. In conclusion, the present study observed a transition in the seroprevalence of HAV, with a higher seropositivity in older children. Age, SEC, source of drinking water, and education status of parents were significantly associated with HAV seroprevalence.

Keywords: children, hepatitis A, infectious disease, seroprevalence

Introduction

Hepatitis A is an acute, self-limiting disease of the liver caused by the hepatitis A virus (HAV), and there are >1.5 million cases worldwide annually.¹ The World Health Organization (WHO) estimates that the global burden of viral hepatitis A increased from 177 million in 1990 to 212 million in 2005.² Overall, 35,245 HAV-related deaths were estimated in 2005.²

India is hyper-endemic for HAV infection. Studies conducted in the 2000s observed that nearly 90% of adolescents, adults, and most children acquired immunity to HAV infection in their preschool years.³ However, recent studies have indicated a shift in epidemiology of HAV infection over the past decade.^{4,5} Chitambar et al⁶ demonstrated a significant ($P<0.05$) shift in epidemiology of HAV infection among children aged 5–10 years from 96.7% in 1983 to 85.1% in 1995. Murhekar et al⁷ showed a decrease in anti-HAV seropositivity among children aged <10 years between 1989 and 1999 (decrease in seropositivity: 80% in 1989 to 60% in 1999; $P=0.019$). Anti-HAV antibodies were detected in 100% of newborns 15 years ago as a result of a rise in maternal exposure to HAV. Recent studies have shown a fall in the number of cases

to 50%–60%.⁸ Acharya et al⁹ opined that improved hygienic and socioeconomic conditions led to a shift in epidemiology of HAV infection.

Incidence of HAV is closely related to socioeconomic development, with prevalence of anti-HAV varying from <50% in high-income regions (Western Europe, Australia, New Zealand, Canada, the USA, Japan, Republic of Korea, Singapore) to as high as >90% in low-income regions (sub-Saharan Africa, parts of South Asia). Most middle-income regions in Asia, Latin America, Eastern Europe, and the Middle East have a mix of intermediate (>50% immunity by 15 years) and low (<50% immunity by 30 years) prevalence of anti-HAV antibodies.¹⁰ Syed et al¹¹ suggested a shift in HAV infection epidemiology in India, like other countries in southeast Asia and People's Republic of China, over the past 20 years from high to moderate endemicity and from moderate to low endemicity due to community living conditions and occupation. In India, there is a growing body of evidence that children from low socioeconomic classes (SECs) continue to have a high prevalence of anti-HAV antibodies.^{12–15} Dutta et al¹⁶ reported considerably lower HAV seroprevalence among children (0–12 years) whose parents were educated and in children with improved sanitary conditions. Arankalle et al¹³ reported that, over a 16-year observation period, lower SEC was associated with a 23-fold higher risk of HAV infection. HAV seropositivity was 64.5% in a population from a higher SEC compared with 85.4% from a lower SEC.¹⁴

Hepatitis A is a vaccine-preventable disease. Although studies conducted in the past have provided adequate evidence of changing HAV epidemiology,^{13,17} recent data are required for national vaccine strategy policy making. The present multicenter study evaluated seroprevalence of HAV among children across different geographical locations of India. The impact of various factors, such as age, SEC, source of drinking water, and education status of parents, on HAV seroprevalence was investigated in Indian children between 18 months and 10 years of age.

Materials and methods

Ethics statement

The study protocol was reviewed and approved by the individual Independent Ethics Committees/Institutional Review Boards (IECs/IRBs) (Ethics Committee of Childs Trust Medical Research Foundation, Chennai; Ethics Committee King Edward Memorial [KEM] Hospital and Research Center, Pune; and Ethics Committee of Institute of Child Health [ICH], Kolkata). The study was conducted in accordance

with the guidelines of the International Conference on Harmonisation of Technical Requirements for Registration of Pharmaceuticals for Human Use/Good Clinical Practice (GCP), and Indian Council of Medical Research (ICMR), and the Declaration of Helsinki.¹⁸

Study characteristics

This was a prospective, cross-sectional, multicenter study conducted in five different centers (ICH, Kolkata; Sree Aurobindo Seva Kendra–Economic Entrepreneurship Development Foundation [EEDF], Kolkata; KEM Hospital, Pune; Kanchi Kamakoti Childs Trust Hospital [KKCTH] Chennai; and Sarvodaya Child Care [SCC], New Delhi) across four cities (New Delhi, Kolkata, Pune, and Chennai) between September 2011 and July 2012.

Inclusion and exclusion criteria

Children (18 months to 10 years) seeking medical treatment for a health problem or minor non-infective illness or attending a follow-up visit or health examination and requiring blood draw for any indication were eligible for the study. All subjects and/or their parents gave written consent to participate in the study. Children with a history of hepatitis A vaccination or current symptoms of viral hepatitis/chronic hepatitis were excluded from the study.

Data collection

A thorough physical examination of the eligible children was performed. The following details were documented:

- medical history;
- area of residence (such as having a home/homeless, rural/urban, if urban whether in an urban slum area/urban non-slum area [an urban non-slum area is defined as a statutory place with a municipality, corporation, cantonment board, or notified town area committee; an urban slum area is defined as a compact area with at least a population of 300/60–70 households of poorly built congested tenements, in an unhygienic environment, usually with inadequate infrastructure lacking proper sanitary and drinking water facilities]);
- home sanitary conditions, such as source of drinking water (tap water, river water, well water, or hand pump) and drinking water treatment (filtered water, boiled water, ultraviolet [UV]/reverse osmosis [RO]-based purified water, or untreated water);
- excreta disposal (private toilet within the house, private toilet outside house, public toilet, or open field);

- education status (professional/honors, graduate/post-graduate, intermediate/post-high school diploma, high school certificate, middle school certificate, primary school certificate, or illiterate); and
- occupation of parents/guardians (professional, semi-professional, clerical/shop owner/farmer, skilled worker, semi-skilled worker, unskilled worker, or unemployed).

Kuppuswamy's scale was used for evaluating the SEC by summing up the score for occupation of the parents/guardians; education of parents/guardians; and family income.¹⁹

Seroprevalence

Blood samples of subjects were collected and the serum samples were screened for the presence and quantitation of total HAV antibodies using enzyme-linked immunosorbent assay (ELISA) AxSYM® HAVAB®-M 2.0 (Abbott Laboratories, Abbot Park, IL, USA). All the tests were conducted centrally in collaboration with the National Institute of Virology, Pune, India. Total anti-HAV levels >0.80 mIU mL⁻¹ were considered to be seropositive.

Statistical analysis

All demographic and socioeconomic parameters were summarized using descriptive statistics. Association between seroprevalence and various factors of sanitation and SEC (separate and combined) was analyzed using the χ^2 test. Binary logistic regression using the likelihood model was performed for seroprevalence and various factors. A *P*-value of <0.05 was considered to be statistically significant. The overall effect of all parameters with HAV seroprevalence were evaluated with 2 log-likelihood, Cox and Snell *R*², and Nagelkerke *R*² tests. Statistical analysis was performed using SPSS 16 software (IBM Corporation, Armonk, NY, USA).

Results

The demographic details of subjects are presented in Table 1. A total of 928 children (N=272 aged 18 months to 3 years; N=322 aged 3–6 years; N=334 aged 6–10 years) were enrolled. Based on the Kuppuswamy SEC scoring system, 69 belonged to a lower SEC, 209 to an upper lower SEC, 250 to a lower middle SEC, 269 to an upper middle SEC, and 131 to an upper SEC. Overall, seropositivity varied between different centers: New Delhi had the highest number of seropositive cases (32/56; 57.1%), followed by Pune (101/240; 42.1%). Heterogeneous pockets of susceptible individuals co-existed in different areas of Kolkata. The seropositivity

Table 1 Demographic profile of subjects

Study center (N)	Age (year)	Female, n (%)	Height (cm)	Weight (kg)
EEDF (72)	5.4±2.6	26 (36.1)	111.8±18	19.0±7.5
ICH (300)	4.6±2.4	109 (36.3)	104.5±21.8	15.4±6.2
KKCTH (260)	4.6±2.3	92 (35.4)	104.3±22.1	17.3±10.2
KEM (240)	4.8±2.4	108 (45)	102.7±21.6	18.7±15.7
SCC (56)	5.2±3	19 (33.9)	107.4±27.5	19.7±12.4
Total (928)	4.7±2.4	354 (38.1)	106.1±22.2	17.3±11

Note: Data are presented as mean ± SD unless otherwise indicated.

Abbreviations: EEDF, Economic Entrepreneurship Development Foundation (Sree Aurobindo Seva Kendra; Kolkata); ICH, Institute of Child Health (Kolkata); KEM, King Edward Memorial Hospital (Pune); KKCTH, Kanchi Kamakoti Childs Trust Hospital (Chennai); SCC, Sarvodaya Child Care (New Delhi); SD, standard deviation.

varied from 17% (12/72) to 48.3% (145/300) in the two centers of Kolkata. In Chennai, seropositivity was observed in 22.3% (58/260) of the subjects.

The number of seropositive cases increased with age across all study centers (Figure 1). The overall seroprevalence of HAV in the group aged 6–10 years (50.3%) was significantly higher (*P*=0.000) than in the group aged 18 months to 6 years (30.3%). The seroprevalence of HAV in different age groups (18 months to 6 years and 6–10 years) and SECs showed a significant increase in all age groups across different SECs (*P*<0.000–0.001 for all) except upper SEC (*P*=0.124).

With respect to SEC, the heterogeneity of seroprevalence in upper and upper middle classes in different centers were 18%, 32%, 28%, 18%, and 54% for EEDF, ICH, KEM, KKCTH, and SCC centers, respectively, reflecting endemicity differences in various regions of India. The pooled seropositivity across SECs was 26%. The overall association of SEC with seroprevalence achieved statistical significance (*P*=0.000), with the highest impact in Kolkata (upper and upper middle SEC 90/29; other SEC 210/116; *P*=0.000).

The role of other variables, such as educational status of the parents, source of drinking water, and sanitary conditions, with HAV seropositivity (Figure 2) was also analyzed. The difference in seroprevalence of HAV in subjects from an educated background (35.7%) and subjects from an illiterate background (52.3%) was statistically significant across all the centers (*P*=0.000). The seroprevalence of HAV was higher in subjects using a hand pump as the source of drinking water (50%) than in those using other sources of drinking water (well water 22.2%; river water 22.7%; tap water 37%). However; the source of drinking water for most subjects was tap water (86.8%) and was not significantly associated with a higher HAV seroprevalence (*P*=0.45) (Figure 2).




Figure 1 Age-specific prevalence (all centers).
Abbreviation: HAV, hepatitis A virus.

Fewer subjects from a household with a private toilet within the house (33.1%) were seropositive compared with those using an open field for defecation (75%). The association of HAV seroprevalence was statistically significant ($P=0.001$) with excreta disposal (Figure 2).

HAV seropositivity was not related to the sex of the subjects, either combined in all centers or when considered for each study center separately. The area of residence did

not show a significant and clear association with HAV seroprevalence in most of the study centers. Only one of the study centers (KEM, Pune) showed a statistically significant ($P=0.001$) association between area of residence and HAV seroprevalence. The binary logistic regression analysis revealed that SEC, age, source of drinking water, and education had a positive impact (68.9%) on the seroprevalence of HAV (Table 2).




Figure 2 Overall impact of all factors on HAV seroprevalence.
Notes: *Overall association of SEC with seroprevalence was statistically significant ($P=0.000$); #difference in HAV seroprevalence among subjects from educated background and from illiterate background was statistically significant ($P=0.000$); *association of HAV seroprevalence with excreta disposal was statistically significant ($P=0.001$).
Abbreviations: HAV, hepatitis A virus; SEC, socioeconomic class.

Table 2 Association between seroprevalence and the determining factors for each center and combining all the centers (level of significance)

Factors	Coefficient	Significance	Exp(B)
Socioeconomic class	-0.639	0.003**	0.528
Age	0.842	0.000**	2.322
Sex	0.255	0.094	1.290
Source of drinking water	0.905	0.030*	2.471
Toilet facility	-0.315	0.236	0.730
Place of residence	-0.083	0.615	0.920
Education	-0.296	0.040*	0.744
Constant	-0.974	0.088	0.378
Model summary	1,087.864 ^a	0.127 ^b	0.173 ^c

Notes: ^a2 Log likelihood; ^bCox and Snell R Square; ^cNagelkerke R Square; * $P < 0.05$; ** $P < 0.01$.

Abbreviation: Exp(B), exponentiation of the B coefficient.

Discussion

This study has demonstrated the current pattern of HAV seroprevalence in Indian children aged below 10 years, taking various SECs into consideration. A higher seroprevalence of HAV was observed among the older children than among the younger children (1.5–2 years 20.6%; 9–10 years 60.9%; $P = 0.000$), demonstrating a shift in HAV epidemiology as observed in previous studies.^{12,14,20,21}

The results of a previous surveillance study demonstrated 98% seropositive cases in the group aged 6–10 years in 1992 as compared with 50.4% in the present study.¹² When compared with the seroprevalence in different cities among different age stratifications over a period of 10–15 years, a significant decline amongst all age groups of approximately 30% was noted. This difference in the number of HAV-seropositive cases over age clearly demonstrates a shift in epidemiology. This may be due to an increase in non-immune subjects over the years, resulting in changing epidemiology that might lead to a profound impact on the magnitude and severity of the disease.

Studies worldwide have shown an association between higher SEC and decreased HAV seroprevalence.²² In India, various surveillance studies over the years in different cities documented a decline in anti-HAV prevalence, mostly because of the transition in demography. Improved hygiene through better education and sanitary conditions has led to an overall improvement in SEC. Dhawan et al¹⁴ demonstrated a significant effect of SEC on HAV seroprevalence; approximately 64.5% in the upper SEC and 85.3% in the lower SEC in the year 1998. The present study demonstrated seroprevalence of 25.7% in the upper SEC and 46.4% in the lower SEC. Although the association between seroprevalence and SEC remains the same as in the previous study, there has been significant decline of seropositivity in both SECs,

confirming a decrease in hepatitis A transmission over the years.

Seroprevalence of HAV varied amongst different cities and within different communities of the same city. Heterogeneous pockets of susceptible individuals co-existed in different areas of Kolkata. This difference in anti-HAV prevalence was related to educational status and toilet facility. This indicated that awareness and education has an impact on the water hygiene and sanitary habits of the population, irrespective of SEC. Economic conditions and drinking water source did not show a significant effect on seroprevalence. In all centers, tap water was the primary source of drinking water, hence an association between source of drinking water and seroprevalence could not be established. However, that most subjects enrolled in the present study had access to tap water as a source of drinking water confirms improved facilities, leading to decreased transmission.

Although several studies have evaluated the seroprevalence of HAV, the effect of educational background has not been considered by most studies conducted in India. A study conducted in Turkey showed a statistically significant ($P = 0.000$) association between HAV seroprevalence and educational status of the mother (university graduate 23.1%; illiterate 92.1%) and father (university graduate 31.8%; illiterate 92.3%).²³ In the present study, the lowest HAV seroprevalence was observed in EEDF, Kolkata, where the educational background of most parents was higher than in other centers (above secondary/total number of subjects, 61/72).

Hepatitis A remains asymptomatic among children below the age of 5 years; however, the severity of this infection increases with age and the disease assumes epidemic proportions. India has experienced HAV outbreaks in children,²⁴ children and young adults,²⁵ and adults.²⁶ The World Health Organization (WHO) recommends large-scale hepatitis A vaccination in countries with improving SEC that have moved from high to intermediate hepatitis A endemicity.¹ Documentation of large proportions of children still susceptible to HAV up to the age of 10 years, and evidence of outbreaks of HAV in the recent past, indicates the need for routine hepatitis A vaccination. Further, international data show that hepatitis A mass vaccination has significantly decreased the incidence of hepatitis A. Vaccination programs have decreased the incidence of HAV infection to a large extent in countries such as the USA (rate of HAV infection per 1,000,000 population reduced from 12 in 1995 to 1.5 in 2005), Israel (resulted in seroprotective antibody concentrations among 100% of children), and Argentina

(average reduction rate between 1998 and 2002 of 88%; $P < 0.001$).^{27–29}

Conclusion

The present study demonstrates heterogeneity in the endemicity of HAV across different pockets of geographical distribution, various centers, and even within centers based on age, educational status, drinking water, and overall SEC status. The rate of seroprevalence has declined across all age group segments. Although the trend is higher in the age group 6–10 years, it has also shown a decrease of 30% within the last decade. This transition in the rate of endemicity has been reported to increase the disease morbidity and mortality, which is otherwise in high endemic zones that are relatively asymptomatic.³⁰

The diverse variation in the rates of seropositivity across various cities in different locations of India in the urban settings confirms the need for routine vaccination. Considering the diversity of endemicity, there is scope for a single-dose vaccine schedule for both inactivated and live attenuated hepatitis A vaccine, as highlighted by the WHO for developing countries.¹

Acknowledgments

This epidemiological research study was sponsored by Wockhardt Limited. All authors participated in the analysis and interpretation of data and critical revision of the manuscript. The authors extend acknowledgement to Professors Monoranjan Pal and Premananda Bharti from the Indian Statistical Institute for statistical analysis. The manuscript was read, revised, and approved by all authors. The authors acknowledge Knowledge Isotopes Pvt, Ltd (<http://www.knowledgeisotopes.com>) for the writing support.

Disclosure

SB, AG, SB, AC, and JC have received a research grant for conducting the research study. SC and MM are employees of Medclin Research Pvt, Ltd, a Clinical Research Organization. MM is the corresponding author and has received payment from Wockhardt Limited for project management services provided by Medclin Research Pvt, Ltd. GK, AM, and SR are employees of Wockhardt Limited. VA has received a research grant for conducting the laboratory investigation.

References

- No authors listed. WHO position paper on hepatitis A vaccines – Jun 2012. *Wkly Epidemiol Rec.* 2012;87:261–276.
- Jacobsen KH, Wiersma ST. Hepatitis A virus seroprevalence by age and world region, 1990 and 2005. *Vaccine.* 2010;28:6653–6657.
- Jacobsen K. *The Global Prevalence of Hepatitis a Virus Infection and Susceptibility: A Systematic Review. Immunization, Vaccines and Biologicals.* Geneva: World Health Organization; 2009. Available from: http://whqlibdoc.who.int/hq/2010/WHO_IVB_10.01_eng.pdf. Accessed December 26, 2012.
- Kar P. Is there a change in seroepidemiology of hepatitis A infection in India? *Indian J Med Res.* 2006;123:727–729.
- Mathur P, Arora NK. Epidemiological transition of hepatitis A in India: issues for vaccination in developing countries. *Indian J Med Res.* 2008;128:699–704.
- Chitambar SD, Chadha MS, Joshi MS, Arankalle VA. Prevalence of hepatitis A antibodies in western Indian population: changing pattern. *Southeast Asian J Trop Med Public Health.* 1999;30:273–276.
- Murhekar MV, Sehgal SC, Murhekar KM, Padbhidri SP, Chitambar SD, Arankalle VA. Changing scenario of hepatitis A virus and hepatitis E virus exposure among the primitive tribes of Andaman and Nicobar Islands, India over the 10-year period 1989–1999. *J Viral Hepat.* 2009;9:315–321.
- Franco E, Meleleo C, Serino L, Sorbara D, Zaratti L. Hepatitis A: epidemiology and prevention in developing countries. *World J Hepatol.* 2012;4:68–73.
- Acharya SK, Madan K, Dattagupta S, Panda SK. Viral hepatitis in India. *Natl Med J India.* 2006;19:203–217.
- World Health Organization. *Evidence Based Recommendation for Use of Hepatitis A Vaccines in Immunization Services: Background Paper for SAGE Discussions.* Geneva: World Health Organization; 2011. Available from: http://www.who.int/immunization/sage/1_HepABackground_17Oct_final2_nov11.pdf. Accessed December 26, 2012.
- Syed R, Mohammed A, Sindiri P, et al. Seroepidemiology of hepatitis A virus in Hyderabad, South India. *J Med Allied Sci.* 2012;2:58–61.
- Arankalle VA, Tsarev SA, Chadha MS, et al. Age-specific prevalence of antibodies to hepatitis A and E viruses in Pune, India, 1982 and 1992. *J Infect Dis.* 1995;171:447–450.
- Arankalle VA, Chadha MS, Chitambar SD, Walimbe AM, Chobe LP, Gandhe SS. Changing epidemiology of hepatitis A and hepatitis E in urban and rural India (1982–1998). *J Viral Hepat.* 2001;8:293–303.
- Dhawan PS, Shah SS, Alvares JF, et al. Seroprevalence of hepatitis A virus in Mumbai, and immunogenicity and safety of hepatitis A vaccine. *Indian J Gastroenterol.* 1998;17:16–18.
- Acharya SK, Batra Y, Bhatkal B, et al. Seroepidemiology of hepatitis A virus infection among school children in Delhi and north Indian patients with chronic liver disease: implications for HAV vaccination. *J Gastroenterol Hepatol.* 2003;18:822–827.
- Dutta AK, Aggarwal A, Kapoor AK, Ray GN, Batra S. Seroepidemiology of hepatitis A in Delhi. *Indian J Pediatr.* 2000;67:77–79.
- Gupta A, Chawla Y. Changing epidemiology of hepatitis A infection. *Indian J Med Res.* 2008;128:7–9.
- Krleza-Jerić K, Lemmens T. 7th revision of the Declaration of Helsinki: good news for the transparency of clinical trials. *Croat Med J.* 2009;50:105–110.
- Kumar N, Gupta N, Kishore J. Kuppaswamy's socioeconomic scale: updating income ranges for the year 2012. *Indian J Public Health.* 2012;56:103–104.
- Aggarwal R, Naik S, Yachha SK, Naik SR. Seroprevalence of antibodies to hepatitis A virus among children in Northern India. *Indian Pediatr.* 1999;36:1248–1250.
- Mall ML, Rai RR, Philip M, et al. Seroepidemiology of hepatitis A infection in India: changing pattern. *Indian J Gastroenterol.* 2001;20:132–135.
- Jacobsen KH, Koopman JS. Declining hepatitis A seroprevalence: a global review and analysis. *Epidemiol Infect.* 2004;132:1005–1022.
- Altinkaynak S, Selimoğlu MA, Ertekin V, Kiliçaslan B. Epidemiological factors affecting hepatitis A seroprevalence in childhood in a developing Country. *Eurasian J Med.* 2008;40:25–28.

24. Chadha MS, Arankalle VA, Jadi RS, Joshi MV, Thakare JP, et al. An outbreak of Chandipura virus encephalitis in the eastern districts of Gujarat state, India. *Am J Trop Med Hyg.* 2005;73:566–570.
25. Chobe LP, Arankalle VA. Investigation of a hepatitis A outbreak from Shimla Himachal Pradesh. *Indian J Med Res.* 2009;130:179–184.
26. Arankalle VA, Sarada Devi KL, Lole KS, Shenoy KT, Verma V, Haneephabi M. Molecular characterization of hepatitis A virus from a large outbreak from Kerala, India. *Indian J Med Res.* 2006;123:760–769.
27. Wasley A, Miller JT, Finelli L; Centers for Disease Control and Prevention (CDC). Surveillance for acute viral hepatitis: United States, 2005. *MMWR Surveill Summ.* 2007;56:1–24.
28. Dagan R, Greenberg D, Weber F. Immunogenicity of an inactivated hepatitis A pediatric vaccine: three-year post-booster follow-up. *Vaccine.* 2005;23:5144–5148.
29. Vacchino MN. Incidence of hepatitis A in Argentina after vaccination. *J Viral Hepat.* 2008;15 Suppl 2:47–50.
30. Verma R, Khanna P. Hepatitis A vaccine should receive priority in National Immunization Schedule in India. *Hum Vaccin Immunother.* 2012;8:1132–1134.

Vaccine: Development and Therapy

Publish your work in this journal

Vaccine: Development and Therapy is an international, peer-reviewed, open access journal that spans the spectrum of vaccine design and development through to clinical applications. The journal is characterized by the rapid reporting of application notes, reviews, original research and clinical studies in all therapeutic areas. Clinical outcomes, patient safety,

Submit your manuscript here: <http://www.dovepress.com/vaccine-development-and-therapy-journal>

and programs for the development and effective, safe, and sustained use of vaccines will be a feature of the journal. The manuscript management system is completely online and includes a very quick and fair peer-review system. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Dovepress