

Two Mode Quantum Systems: Invariant Classification of Squeezing Transformations and Squeezed States

Arvind*

Department of Physics

Indian Institute of Science, Bangalore - 560 012, India

B. Dutta

Jawaharlal Nehru Centre for Advanced Scientific Research

IISc Campus, Bangalore - 560 012, India

N. Mukunda[†]

Centre for Theoretical Studies and Department of Physics

Indian Institute of Science, Bangalore - 560 012, India

and

R. Simon

Institute of Mathematical Sciences, C. I. T. Campus, Madras - 600 113

(August 21, 2002)

03.65.Fd, 42.50.Dv, 42.50.Lc

Typeset using REVTeX

Abstract

A general analysis of squeezing transformations for two mode systems is given based on the four dimensional real symplectic group $Sp(4, \mathbb{R})$. Within the framework of the unitary metaplectic representation of this group, a distinction between compact photon number conserving and noncompact photon number nonconserving squeezing transformations is made. We exploit the $Sp(4, \mathbb{R}) - SO(3, 2)$ local isomorphism and the $U(2)$ invariant squeezing criterion to divide the set of all squeezing transformations into a two parameter family of distinct equivalence classes with representative elements chosen for each class. Familiar two mode squeezing transformations in the literature are recognized in our framework and seen to form a set of measure zero. Examples of squeezed coherent and thermal states are worked out. The need to extend the heterodyne detection scheme to encompass all of $U(2)$ is emphasized, and known experimental situations where all $U(2)$ elements can be reproduced are briefly described.

I. INTRODUCTION

The theoretical analysis [1] and experimental [2], [3], [4] realization of squeezed states of radiation continue to receive a great deal of attention. While much of the work so far has concerned itself with single mode situations [5], [6] some analysis of two-mode states has also been presented [7], [8]. Other nonclassical effects of radiation beyond second order have also received attention in the literature [9]. More recently, a general invariant squeezing criterion for n -mode systems has been developed by some of us elsewhere [10].

The purpose of the present paper is to study squeezing transformations for two-mode systems, and to develop a classification scheme for them motivated by the above mentioned invariant squeezing criterion. Basic to all such discussions is the four-dimensional real symplectic group $Sp(4, \mathbb{R})$, of real linear homogeneous canonical transformations, and the unitary metaplectic representation of this group acting on the Hilbert space of states for a two-mode quantum system. The structure of the noncompact group $Sp(4, \mathbb{R})$ (and its n -mode counterpart $Sp(2n, \mathbb{R})$) leads to a natural separation of its elements into passive (compact), and active (noncompact) types. Here the adjectives *passive* and *active* mean total photon number conserving and nonconserving respectively. This group theoretical framework gives us an unambiguous way of defining precisely the family of squeezing transformations; they are the active elements of $Sp(4, \mathbb{R})$ and they do not form a subgroup. The action of the maximal compact (passive) subgroup $U(2)$ of $Sp(4, \mathbb{R})$ on the set of squeezing transformations by conjugation leads to a natural equivalence relation, leading to the emergence of equivalence classes and convenient representative elements as well. In studying the physical properties of a state subjected to squeezing, therefore, we are able to isolate the dependence on intrinsic squeezing parameters and separate them from other passive factors. As might be expected, the single *squeeze factor* encountered in the studies of single mode states gets enlarged here to *two independent intrinsic* squeeze factors; and it turns out that the two mode squeezing transformations so far studied in the literature form a very small subset of all the independent available possibilities.

The material in this paper is arranged as follows: Section II sets up the basic kinematics for two mode systems, the Fock and Schrödinger representations and the actions of $Sp(4, \mathbb{R})$ on the canonical variables and states. The variance matrix for a general state and its change under $Sp(4, \mathbb{R})$ are derived. After identifying the maximal compact or passive $U(2)$ subgroup of $Sp(4, \mathbb{R})$, the $U(2)$ invariant squeezing criterion for two mode systems is discussed. Section III introduces the generators for the metaplectic representation of $Sp(4, \mathbb{R})$ and brings out the connection to the $SO(3, 2)$ Lie algebra. The photon number conserving compact generators and the remaining noncompact ones are clearly identified. Based on the polar decomposition theorem for general elements of $Sp(4, \mathbb{R})$ we are then led to a precise definition of squeezing transformations: These are single exponentials of linear combinations of the noncompact metaplectic generators. We then proceed to break up the set of all squeezing transformations into equivalence classes under $U(2)$ action. We find that these classes form a continuous two parameter family describable by points in an octant in a two dimensional plane. The families of Caves-Schumaker transformations and essentially single mode transformations correspond to one dimensional lines bounding this octant and so are of measure zero. Section IV applies our formalism to two mode squeezed coherent and thermal states. In Section V we see how the heterodyne detection scheme fits into our analysis. We argue that it is necessary to experimentally realise all elements of the $U(2)$ subgroup of $Sp(4, \mathbb{R})$; the heterodyne scheme only handles a one parameter subset of $U(2)$. Two examples of two-mode situations where all elements of $U(2)$ can be experimentally realised are briefly described. Section VI contains some concluding remarks.

II. SYMPLECTIC GROUP FOR TWO MODES AND THE SQUEEZING CRITERION

We consider two orthogonal modes of the radiation field, with annihilation operators $a_j, j = 1, 2$, and corresponding creation operators $a_j^\dagger$. These two modes could, for example, be two different frequencies for the same or different propagation directions and polarizations,

two different propagation directions at a common frequency, two different polarization states of plane waves degenerate in frequency and direction of propagation, etc. We arrange these operators in the form of a four-component column vector:

$$\xi^{(c)} = (\xi_a^{(c)}) = \begin{pmatrix} a_1 \\ a_2 \\ a_1^\dagger \\ a_2^\dagger \end{pmatrix},$$

$$a = 1, 2, 3, 4. \quad (2.1)$$

The superscript (c) indicates that the entries here are complex, i.e., nonhermitian, operators. For discussion of quadrature squeezing, however, we need to also deal with the hermitian quadrature components of these operators. Therefore we define another column vector ξ with four hermitian entries, related to $\xi^{(c)}$ by a fixed numerical matrix, as follows:

$$\xi = (\xi_a) = \begin{pmatrix} q_1 \\ q_2 \\ p_1 \\ p_2 \end{pmatrix},$$

$$q_j = \frac{1}{\sqrt{2}}(a_j + a_j^\dagger), \quad p_j = \frac{-i}{\sqrt{2}}(a_j - a_j^\dagger);$$

$$\xi^{(c)} = \Omega \xi, \quad \xi = \Omega^{-1} \xi^{(c)},$$

$$\Omega = (\Omega^{-1})^\dagger = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 0 & i & 0 \\ 0 & 1 & 0 & i \\ 1 & 0 & -i & 0 \\ 0 & 1 & 0 & -i \end{pmatrix}. \quad (2.2)$$

The canonical commutation relations can now be written either in terms of ξ or (in two ways) in terms of $\xi^{(c)}$:

$$[\xi_a, \xi_b] = i\beta_{ab},$$

$$[\xi_a^{(c)}, \xi_b^{(c)}] = \beta_{ab},$$

$$\begin{aligned}
[\xi_a^{(c)}, \xi_b^{(c)\dagger}] &= (\Sigma_3)_{ab}, \\
(\beta_{ab}) &= \begin{pmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ -1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \end{pmatrix}, \\
((\Sigma_3)_{ab}) &= \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & -1 \end{pmatrix}.
\end{aligned} \tag{2.3}$$

A general real linear homogeneous transformation on the q 's and p 's is described by a 4×4 real matrix S acting as follows:

$$\begin{aligned}
\xi &\rightarrow \xi' = S\xi \\
\xi'_a &= S_{ab}\xi_b.
\end{aligned} \tag{2.4}$$

If the ξ'_a are to satisfy the same commutation relations as the ξ_a , the condition on S is:

$$S \beta S^T = \beta. \tag{2.5}$$

This is the defining property for the elements of the group $Sp(4, \mathfrak{R})$, which is a noncompact group:

$$Sp(4, \mathfrak{R}) = \left\{ S = 4 \times 4 \text{ real matrix} \mid S \beta S^T = \beta \right\}. \tag{2.6}$$

Note that $\beta = -\beta^{-1}$ itself is an element of $Sp(4, \mathfrak{R})$ whereas Σ_3 is not. Further, $S \in Sp(4, \mathfrak{R})$ implies $-S, S^T, S^{-1} = \beta S^T \beta^T \in Sp(4, \mathfrak{R})$ and $\det S = 1$ for every $S \in Sp(4, \mathfrak{R})$.

The action of $Sp(4, \mathfrak{R})$ on the nonhermitian operators $\xi_a^{(c)}$ is by a (generally) complex matrix $S^{(c)}$ related to S by conjugation with Ω :

$$\begin{aligned}
S \in Sp(4, \mathfrak{R}) : \quad \xi' = S\xi &\Rightarrow \\
\xi^{(c)'} &= S^{(c)} \xi^{(c)}, \\
S^{(c)} &= \Omega S \Omega^\dagger.
\end{aligned} \tag{2.7}$$

We denote the Hilbert space on which ξ and $\xi^{(c)}$ act by $\mathcal{H}$. Since the hermiticity properties and commutation relations of the ξ_a are maintained by the transformation (2.4) for any $S \in Sp(4, \mathbb{R})$, and since the ξ_a act irreducibly on $\mathcal{H}$, it follows from the Stone-von Neumann theorem [11] that it should be possible to construct a unitary operator $\mathcal{U}(S)$ on $\mathcal{H}$ implementing (2.4) via conjugation:

$$\begin{aligned} S \in Sp(4, \mathbb{R}) : S_{ab}\xi_b &= \mathcal{U}(S)^{-1} \xi_a \mathcal{U}(S), \\ \mathcal{U}(S)^\dagger \mathcal{U}(S) &= \mathbf{1} \text{ on } \mathcal{H}. \end{aligned} \quad (2.8)$$

This $\mathcal{U}(S)$, for each S , is clearly arbitrary upto (and only upto by virtue of irreducibility) an S -dependent phase. Even after making use of this freedom, however, it turns out that we cannot choose the operators $\mathcal{U}(S)$ for various S so as to give us a true unitary representation of $Sp(4, \mathbb{R})$ on $\mathcal{H}$. Rather, after maximum simplification, they give us a two-valued representation of this group [12]:

$$S_1, S_2 \in Sp(4, \mathbb{R}) : \mathcal{U}(S_1) \mathcal{U}(S_2) = \pm \mathcal{U}(S_1 S_2). \quad (2.9)$$

Alternatively we can regard the $\mathcal{U}(S)$, chosen so as to obey this composition law, as providing a true and faithful unitary representation of the four-dimensional metaplectic group $Mp(4)$, which in turn is a two-fold cover of $Sp(4, \mathbb{R})$. In the literature the operators $\mathcal{U}(S)$ are often said to provide the metaplectic representation of $Sp(4, \mathbb{R})$. The connection between $Mp(4)$ and $Sp(4, \mathbb{R})$ has some similarity with the one between $SU(2)$ and $SO(3)$ familiar in angular momentum theory.

Now we consider physical states of the two-mode system, the action of $Sp(4, \mathbb{R})$ on them, and the statement of a suitable squeezing criterion. Let ρ be the density operator of any (pure or mixed) state of the two-mode radiation field. With no loss of generality we may assume that the means $\text{Tr}(\rho \xi_a)$ of ξ_a vanish in this state. (Any non-zero values for these means can always be reinstated by a suitable phase space displacement which has no effect on the squeezing properties). Squeezing involves the set of all second order noise moments of the quadrature operators q_j and p_j . To handle them collectively we define the variance or noise matrix V for the state ρ as follows:

$$\begin{aligned}
V &= (V_{ab}), \\
V_{ab} &= V_{ba} = \frac{1}{2} \text{Tr}(\rho \{\xi_a, \xi_b\}).
\end{aligned} \tag{2.10}$$

This definition is valid for a system with any number of modes. For a two-mode system it can be written explicitly in terms of q_j and p_j as:

$$V = \begin{pmatrix} \langle q_1^2 \rangle & \langle q_1 q_2 \rangle & \frac{1}{2} \langle \{q_1, p_1\} \rangle & \langle q_1 p_2 \rangle \\ \langle q_1 q_2 \rangle & \langle q_2^2 \rangle & \langle q_2 p_1 \rangle & \frac{1}{2} \langle \{q_2, p_2\} \rangle \\ \frac{1}{2} \langle \{q_1, p_1\} \rangle & \langle q_2 p_1 \rangle & \langle p_1^2 \rangle & \langle p_1 p_2 \rangle \\ \langle q_1 p_2 \rangle & \frac{1}{2} \langle \{q_2, p_2\} \rangle & \langle p_1 p_2 \rangle & \langle p_2^2 \rangle \end{pmatrix} \tag{2.11}$$

This matrix is real symmetric positive definite and obeys additional inequalities expressing the Heisenberg uncertainty principle of quantum mechanics [10].

When the state ρ is transformed to a new state ρ' by the unitary operator $\mathcal{U}(S)$ for some $S \in Sp(4, \mathbb{R})$, we see easily from eqs (2.8, 2.10) that the variance matrix V undergoes a symmetric symplectic transformation:

$$\begin{aligned}
S \in Sp(4, \mathbb{R}) : \rho' &= \mathcal{U}(S) \rho \mathcal{U}(S)^{-1} \Rightarrow \\
V' &= S V S^T.
\end{aligned} \tag{2.12}$$

This transformation law for V preserves all the properties mentioned at eq. (2.11).

Towards setting up a squeezing criterion, we identify an important subgroup of $Sp(4, \mathbb{R})$, namely its maximal compact subgroup $K \equiv U(2)$. This consists of matrices $S \in Sp(4, \mathbb{R})$ having a specific block form determined by two-dimensional unitary matrices belonging to $U(2)$:

$$\begin{aligned}
K = U(2) &= \left\{ S(X, Y) \in Sp(4, \mathbb{R}) \mid U = X - iY \in U(2) \right\} \subset Sp(4, \mathbb{R}), \\
S(X, Y) &= \begin{pmatrix} X & Y \\ -Y & X \end{pmatrix}.
\end{aligned} \tag{2.13}$$

]Here X and Y are the real and imaginary parts of U ; and the unitary condition $U^\dagger U = \mathbf{1}$ on U ensures that $S(X, Y)$ obeys the symplectic condition (2.5). We also recognise that

this subgroup K is the intersection of the symplectic and the orthogonal groups in four real dimensions, namely

$$K = U(2) = Sp(4, \mathfrak{R}) \cap O(4) :$$

$$S(X, Y) (\beta \text{ or } \mathbf{1}) S(X, Y)^T = \beta \text{ or } \mathbf{1}. \quad (2.14)$$

The complex form $S^{(c)}(X, Y)$ corresponding to $S(X, Y)$, and which acts on $\xi^{(c)}$, is rather simple; it is given by

$$S^{(c)}(X, Y) \equiv S^{(c)}(U)$$

$$= \begin{pmatrix} X - iY & 0 \\ 0 & X + iY \end{pmatrix} \quad (2.15)$$

Thus the maximal compact subgroup of $Sp(4, \mathfrak{R})$ mixes a_1 and a_2 unitarily, *but does not mix the a_j with the $a_j^\dagger$* . This fact may be explained by saying that the subgroup $U(2)$ of $Sp(4, \mathfrak{R})$ consists of *passive*, or total photon number conserving, transformations.

We see that the *physical requirement* that the total number of photons be conserved singles out a unique maximal compact subgroup of $Sp(4, R)$ out of many equivalent ones. In contrast, elements of $Sp(4, \mathfrak{R})$ outside the subgroup $U(2)$ are *non-compact* elements which do *not* conserve total photon number, and so describe *active* transformations. (These properties of *compact* and *non-compact* elements of $Sp(4, \mathfrak{R})$ will become transparent when we identify their generators in the following Section).

As has been discussed in detail elsewhere [10], [13], for a multimode system it is physically reasonable to set up a definition of squeezing which is invariant under the subgroup of passive transformations of the full symplectic group. For the present case of two-mode systems, we evidently need a $U(2)$ -invariant squeezing criterion. That is, our definition must be such that if a state ρ with variance matrix V is found to be squeezed, then the state $\mathcal{U}(S(X, Y)) \rho \mathcal{U}(S(X, Y))^{-1}$ with variance matrix $V' = S(X, Y) V S(X, Y)^T$ must also be squeezed, for any $U = X - iY \in U(2)$.

Conventionally a state is said to be squeezed if any one of the diagonal elements of V is less than $1/2$. The diagonal elements correspond, of course, to fluctuations in the “chosen”

set of quadrature components of the system. The $U(2)$ -invariant definition is as follows: the state ρ is a quadrature squeezed state if either some diagonal element of V is less than $1/2$ (and then we say that the state is manifestly squeezed), or some diagonal element of $V' = S(X, Y) V S(X, Y)^T$ for some $U = X - iY \in U(2)$ is less than $1/2$:

$$\begin{aligned} \rho \text{ is a squeezed state } &\Leftrightarrow \\ &\left(S(X, Y) V S(X, Y)^T \right)_{aa} < \frac{1}{2} \\ &\text{for some } a \text{ and some } X - iY \in U(2). \end{aligned} \tag{2.16}$$

That is, running over $S(X, Y) \in K$ is the same as running over all possible sets of quadrature components. We may say that since any element of $U(2)$ passively mixes the two modes, the appropriate $S(X, Y) \in K$ which achieves the above inequality for some a (assuming the given V permits the same) just chooses the right combination of quadratures to make the otherwise possibly hidden squeezing manifest.

To implement this definition in practice, it would appear that even if a state is intrinsically squeezed, we may have to explicitly find a suitable $U(2)$ transformation which when applied to V makes the squeezing manifest. This however could be complicated. Here the point to be noticed and appreciated is that diagonalisation of a noise matrix V generally requires a real orthogonal transformation belonging to $SO(4)$ which may not lie in $U(2) = O(4) \cap Sp(4, \mathbb{R})$. It is therefore remarkable that, as shown in [10], the $U(2)$ -invariant squeezing criterion (2.16) can be expressed in terms of the spectrum of eigenvalues of V , namely:

$$\rho \text{ is a squeezed state } \Leftrightarrow \ell(V) = \text{least eigenvalue of } V < \frac{1}{2}. \tag{2.17}$$

That is, while the diagonalization of V is in general not possible within $K = U(2)$ which is a proper subgroup of $O(4)$, any one particular (and hence the smallest) eigenvalue of V can be made to become one of the diagonal elements of V transformed by an appropriate $S(X, Y) \in K$. In other words any quadrature component can be taken to any other quadrature component by a suitable element of $U(2)$. We shall in the sequel work with the

$U(2)$ -invariant squeezing criterion (2.16, 2.17).

III. THE $Sp(4, \mathbb{R}) - SO(3, 2)$ CONNECTION AND CLASSIFICATION OF TWO-MODE SQUEEZING TRANSFORMATIONS

We have shown in the previous Section that the group $Sp(4, \mathbb{R})$ of linear canonical transformations contains two kinds of elements: passive total photon-number conserving elements belonging to the maximal compact subgroup $K = U(2)$; and active *noncompact* elements lying outside this subgroup, and which do not conserve total photon number. It is clear from the $U(2)$ -invariant squeezing criterion (2.16, 2.17) that the former elements cannot produce squeezing. This is because the corresponding changes in the variance matrix, $V' = S(X, Y) V S(X, Y)^T$, being similarity transformations preserve the eigenvalue spectrum of V ; hence $\ell(V) \geq 1/2$ implies $\ell(V') \geq 1/2$ and conversely for every $S(X, Y) \in U(2)$. The *noncompact* elements of $Sp(4, \mathbb{R})$, on the other hand, while they do not form a subgroup, have the potential to produce a squeezed state starting from a nonsqueezed state. Thus they may be called squeezing transformations. The following questions then naturally arise: what are the really distinct squeezing transformations which are not related to each other by just passive transformations, and how can they be invariantly labelled or parametrised?

To answer these questions it is useful to work at the level of the Lie algebra $\underline{Sp(4, \mathbb{R})}$ of $Sp(4, \mathbb{R})$, and the hermitian generators of the unitary metaplectic representation $\mathcal{U}(S)$ acting on $\mathcal{H}$. As is well known, these generators are basically all possible hermitian symmetric quadratic expressions in the canonical variables q and p . They may be expressed more transparently for our purposes in terms of a 's and $a^\dagger$'s. While it is possible to set up the generators in a uniform manner for all $Sp(2n, \mathbb{R})$, in the two-mode case we can exploit the fact that the group $Sp(4, \mathbb{R})$ is locally isomorphic to the de Sitter group $SO(3, 2)$ [14], as they share the same Lie algebra. We therefore choose the basis for $\underline{Sp(4, \mathbb{R})}$ to make this explicit; it helps us visualise geometrically the analysis to follow [15]. We define the ten generators $Q, J_r, K_r, L_r, r = 1, 2, 3$ of $Sp(4, \mathbb{R})$ as:

$$Q = \frac{1}{2}(N + 1) = \frac{1}{2}(a_1^\dagger a_1 + a_2^\dagger a_2 + 1); \quad (3.1a)$$

$$\begin{aligned} J_1 &= \frac{1}{2}(a_1^\dagger a_2 + a_2^\dagger a_1), \\ J_2 &= \frac{i}{2}(a_2^\dagger a_1 - a_1^\dagger a_2), \\ J_3 &= \frac{1}{2}(a_1^\dagger a_1 - a_2^\dagger a_2); \end{aligned} \quad (3.1b)$$

$$\begin{aligned} K_1 &= \frac{1}{4}(a_1^{\dagger 2} + a_1^2 - a_2^{\dagger 2} - a_2^2), \\ K_2 &= -\frac{i}{4}(a_1^{\dagger 2} - a_1^2 + a_2^{\dagger 2} - a_2^2), \\ K_3 &= -\frac{1}{2}(a_1^\dagger a_2^\dagger + a_1 a_2); \end{aligned} \quad (3.1c)$$

$$\begin{aligned} L_1 &= \frac{i}{4}(a_1^{\dagger 2} - a_1^2 - a_2^{\dagger 2} + a_2^2), \\ L_2 &= \frac{1}{4}(a_1^{\dagger 2} + a_1^2 + a_2^{\dagger 2} + a_2^2), \\ L_3 &= -\frac{i}{2}(a_1^\dagger a_2^\dagger - a_1 a_2). \end{aligned} \quad (3.1d)$$

These particular quadratic expressions in a_j and $a_j^\dagger$ are chosen because the commutation relations have a suggestive simple form:

$$\begin{aligned} [J_r, J_s] &= i\epsilon_{rst} J_t, \\ [Q, J_r] &= 0; \end{aligned} \quad (3.2a)$$

$$\begin{aligned} [J_r, K_s \text{ or } L_s] &= i\epsilon_{rst}(K_t \text{ or } L_t), \\ [Q, K_r \pm iL_r] &= \mp(K_r \pm iL_r); \end{aligned} \quad (3.2b)$$

$$\begin{aligned} [K_r, K_s] &= [L_r, L_s] = -i\epsilon_{rst} J_t, \\ [K_r, L_s] &= i\delta_{rs} Q. \end{aligned} \quad (3.2c)$$

Comparing and combining the above two sets of equations we see that the four operators Q and J_r are the photon-number conserving generators of $U(2)$ (being respectively the generators of the $U(1)$ and $SU(2)$ parts of $U(2)$); while the six operators K_r and L_r are the *noncompact* generators of squeezing transformations.

The connection to $SO(3, 2)$ becomes evident by regarding these generators as the various components of an antisymmetric set $M_{AB} = -M_{BA}$, where the indices A, B go over the range

1, 2, \dots, 5:

$$\begin{aligned} Q &= M_{45}, & J_r &= \frac{1}{2}\epsilon_{rst}M_{st}; \\ K_r &= M_{r4}, & L_r &= M_{r5}. \end{aligned} \tag{3.3}$$

Then the commutation relations (3.2) take on the de Sitter form

$$\begin{aligned} [M_{AB}, M_{CD}] &= i(g_{AC}M_{BD} - g_{BC}M_{AD} + g_{AD}M_{CB} - g_{BD}M_{CA}), \\ g_{AB} &= \text{diag.}(+1, +1, +1, -1, -1). \end{aligned} \tag{3.4}$$

In this picture, Q, J_r are generators of rotations in the 4-5 plane and in the 1-2-3 subspace respectively, together making up the maximal compact $SO(2) \times SO(3)$ subgroup of $SO(3, 2)$. With respect to the Fock or two mode photon number basis $|n_1, n_2\rangle$ for $\mathcal{H}$ the situation is that all these vectors with a fixed total number $n = n_1 + n_2$, $(n + 1)$ in all, are eigenvectors of Q with a common eigenvalue $(n + 1)/2$; and they simultaneously provide the *spin* $n/2$ representation of the $SO(3)$ subgroup generated by J_r . On the other hand, the K_r and $L_r, r = 1, 2, 3$ are noncompact *Lorentz boost* generators in the $r - 4$ and $r - 5$ planes respectively.

With this algebraic background, we go on to consider two-mode squeezing transformations. It is a well known fact [15] [16] that each matrix $S \in Sp(4, \mathbb{R})$ can be decomposed, globally and uniquely, into the product of two particular kinds of $Sp(4, \mathbb{R})$ matrices: one factor belongs to the subgroup K , the other to a subset Π defined in the following way:

$$\Pi = \left\{ S \in Sp(4, \mathbb{R}) \mid S^T = S = \text{positive definite} \right\} \subset Sp(4, \mathbb{R}). \tag{3.5}$$

We shall hereafter denote elements in Π by $P, P', \dots$. The decomposition mentioned above is then:

$$\begin{aligned} S \in Sp(4, \mathbb{R}) : \quad S &= P S(X, Y), \\ P &\in \Pi, \\ X - iY &\in U(2), \end{aligned} \tag{3.6}$$

with both factors being uniquely determined by S . This is the form in the present context of the familiar polar decomposition formula [17] for a general matrix. For the metaplectic operators $\mathcal{U}(S)$ we have the corresponding statement

$$\begin{aligned}\mathcal{U}(S) &= \mathcal{U}(P) \mathcal{U}(S(X, Y)), \\ \mathcal{U}(P) &= \exp \left[i(\text{real linear combination of } \vec{K} \text{ and } \vec{L}) \right], \\ \mathcal{U}(S(X, Y)) &= \exp \left[i(\text{real linear combination of } Q \text{ and } \vec{J}) \right].\end{aligned}\tag{3.7}$$

We may now identify precisely the most general squeezing transformation within the $Sp(4, \mathbb{R})$ framework, as the operator $\mathcal{U}(P)$ characterised by two numerical three-dimensional vectors $\vec{k}, \vec{l}$ appearing as coefficients of $\vec{K}$ and $\vec{L}$ in the exponent:

$$\mathcal{U}(\vec{k}, \vec{l}) = \exp \left[i(\vec{k} \cdot \vec{K} + \vec{l} \cdot \vec{L}) \right].\tag{3.8}$$

Thus we reserve the name *squeezing transformations* for these elements of Π within $Sp(4, \mathbb{R})$, represented in the metaplectic representation by a single exponential factor. (It is well to keep in mind that Π is not a subgroup, so the product of two such single exponential squeezing transformations is in general not a similar single exponential. This is analogous to the well known fact that the product of two $SO(3, 1)$ Lorentz boosts is not just a boost but a boost followed or preceded by a rotation called the Wigner rotation [18]).

We may relate the decomposition (3.7) of a general metaplectic transformation to a general quadratic Hamiltonian quite directly. Any such Hamiltonian containing both photon conserving and nonconserving terms with possibly time dependent coefficients would lead via the Schrödinger equation to a unitary finite time evolution operator which can be uniquely decomposed into the product form (3.7). Thus, integration of the Schrödinger equation leads in general to a specific passive factor and another specific squeezing transformation. In case the Hamiltonian is time independent and a combination only of the generators $\vec{K}$ and $\vec{L}$, this evolution operator is already of the form $\mathcal{U}(P)$.

Since we have a $U(2)$ -invariant squeezing criterion, as we have seen, elements of $U(2)$ have no effect on the squeezed or nonsqueezed status of any given state. This means that

the $U(2)$ transform, by conjugation, of a squeezing transformation is another squeezing transformation which should be regarded as equivalent to the first one. It is clear that, in any case, any equivalence relation among squeezing transformations as defined by us above should be based on processes which take one squeezing transformation to another.

Now from the commutation relations (3.2b) we can see that the generators $\vec{K}$, $\vec{L}$, and the squeezing transformations $\mathcal{U}(\vec{k}, \vec{l})$ defined in eq. (3.8), behave as follows under conjugation by elements of $U(2)$:

$$\begin{aligned} e^{i\theta Q} (\vec{K} \pm i\vec{L}) e^{-i\theta Q} &= e^{\mp i\theta} (\vec{K} \pm i\vec{L}), \\ e^{i\vec{\alpha} \cdot \vec{J}} (K_r \text{ or } L_r) e^{-i\vec{\alpha} \cdot \vec{J}} &= R_{sr}(\vec{\alpha}) (K_s \text{ or } L_s); \end{aligned} \quad (3.9a)$$

$$\begin{aligned} e^{i\theta Q} \mathcal{U}(\vec{k}, \vec{l}) e^{-i\theta Q} &= \mathcal{U}(\vec{k}', \vec{l}'), \\ \begin{pmatrix} \vec{k}' \\ \vec{l}' \end{pmatrix} &= \begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix} \begin{pmatrix} \vec{k} \\ \vec{l} \end{pmatrix}; \end{aligned} \quad (3.9b)$$

$$\begin{aligned} e^{i\vec{\alpha} \cdot \vec{J}} \mathcal{U}(\vec{k}, \vec{l}) e^{-i\vec{\alpha} \cdot \vec{J}} &= \mathcal{U}(\vec{k}'', \vec{l}''), \\ k_r'' \text{ or } l_r'' &= R_{rs}(\vec{\alpha}) (k_s \text{ or } l_s); \end{aligned} \quad (3.9c)$$

$$\begin{aligned} R_{rs}(\vec{\alpha}) &= \delta_{rs} \cos \alpha + \alpha_r \alpha_s \frac{1 - \cos \alpha}{\alpha^2} + \epsilon_{rst} \alpha_t \frac{\sin \alpha}{\alpha}, \\ \alpha &= |\vec{\alpha}|. \end{aligned} \quad (3.9d)$$

Here we have listed separately the effects of $U(1)$ and $SU(2)$ within $U(2)$ on the generators and the squeezing transformations. The questions raised at the start of this Section can now be posed more precisely: If the set of squeezing transformations $\mathcal{U}(\vec{k}, \vec{l})$ is separated into distinct nonoverlapping equivalence classes based on the $U(2)$ action (3.9), how can we conveniently choose $U(2)$ -invariant parameters to label these classes, and then pick a convenient representative element from each class? We answer these questions in this sequence.

It is clear that we need to construct a complete set of independent expressions in $\vec{k}$ and $\vec{l}$, invariant under both $U(1)$ and $SU(2)$ actions (3.9b, 3.9c). We begin by defining the matrix M of scalar products among $\vec{k}$ and $\vec{l}$, which is then $SU(2)$ invariant:

$$M(\vec{k}, \vec{l}) = \begin{pmatrix} \vec{k} \cdot \vec{k} & \vec{k} \cdot \vec{l} \\ \vec{k} \cdot \vec{l} & \vec{l} \cdot \vec{l} \end{pmatrix}. \quad (3.10)$$

This is a real, symmetric positive semi-definite matrix. With respect to $U(1)$ action, we see from eq. (3.9b) that $M(\vec{k}, \vec{l})$ undergoes a similarity transformation by the rotation matrix of angle θ :

$$M(\vec{k}', \vec{l}') = R(\theta) M(\vec{k}, \vec{l}) R(\theta)^{-1},$$

$$R(\theta) = \begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix}. \quad (3.11)$$

One now sees that there are two independent $U(2)$ invariants that can be formed:

$$\mathfrak{S}_1(\vec{k}, \vec{l}) = \det M(\vec{k}, \vec{l}) = |\vec{k} \wedge \vec{l}|^2,$$

$$\mathfrak{S}_2(\vec{k}, \vec{l}) = \text{Tr } M(\vec{k}, \vec{l}) = |\vec{k}|^2 + |\vec{l}|^2; \quad (3.12)$$

and it is easily checked that there are no other invariants independent of these.

Next let us tackle the problems of finding convenient parameters and representative squeezing transformations for the $U(2)$ equivalence classes, one for each class. We see from eq. (3.12) that if $\mathfrak{S}_1 > 0$ (i.e. $\mathfrak{S}_1 \neq 0$) then the two vectors $\vec{k}$ and $\vec{l}$ are both nonzero and nonparallel; while if $\mathfrak{S}_1 = 0$ they are parallel (and one of them could vanish). These are therefore clearly different geometrical situations. Starting with the matrix $M(\vec{k}, \vec{l})$, we see from its $U(1)$ transformation law (3.11) that by a suitable choice of the angle θ we can arrange the transformed matrix $M(\vec{k}', \vec{l}')$ to be diagonal, and in the case of unequal eigenvalues to place the larger eigenvalue in the first position. This means that in each equivalence class of squeezing transformations there certainly are elements $\mathcal{U}(\vec{k}, \vec{l})$ for which $\vec{k} \cdot \vec{l} = 0$ and $|\vec{k}| \geq |\vec{l}|$. This still leaves us the freedom of action by $SU(2)$. We may now exploit this freedom to put the (mutually perpendicular) vectors $\vec{k}$ and $\vec{l}$ into a convenient geometrical configuration. A look at the forms of the noncompact generators $\vec{K}$ and $\vec{L}$ in eq. (3.1) suggests that we choose $\vec{k}$ and $\vec{l}$ as follows:

$$\vec{k} = (0, a, 0), \quad \vec{l} = (b, 0, 0), \quad a \geq b. \quad (3.13)$$

(A further reason for this choice will emerge shortly). $\mathfrak{S}_1$ and $\mathfrak{S}_2$ can now be evaluated in terms of a, b to obtain the relations

$$\begin{aligned}\mathfrak{S}_1(\vec{k}, \vec{l}) &= a^2 b^2, \\ \mathfrak{S}_2(\vec{k}, \vec{l}) &= a^2 + b^2, \\ a &\geq b \geq 0, \quad (a, b) \neq (0, 0).\end{aligned}\tag{3.14}$$

With this parametrisation we can now say: there is a two-fold infinity of distinct equivalence classes of squeezing transformations for two-mode systems, each class corresponding uniquely and unambiguously to a point (a, b) in the octant $a \geq b \geq 0$ in the real $a - b$ plane, excluding the origin. Different points in the octant correspond to intrinsically distinct equivalence classes. Within an equivalence class determined by a point (a, b) , of course, one can connect different squeezing transformations $\mathcal{U}(\vec{k}, \vec{l})$ by conjugation with suitable $U(2)$ elements. Given a squeezing transformation $P \in \Pi \subset Sp(4, \mathfrak{R})$ its class (a, b) is determined by solving the equations

$$\begin{aligned}\text{Tr}(P) &= 2[\cosh \frac{(a-b)}{2} + \cosh \frac{(a+b)}{2}] \\ \text{Tr}(P^2) &= 2[\cosh(a-b) + \cosh(a+b)]\end{aligned}\tag{3.15}$$

subject to the conditions on a and b appearing in eq. (3.14).

Then we have the following convenient two-mode squeezing transformation representing the equivalence class (a, b) :

$$\begin{aligned}\mathcal{U}^{(0)}(a, b) &= \mathcal{U}^{(0)}(a, 0) \mathcal{U}^{(0)}(0, b), \\ \mathcal{U}^{(0)}(a, 0) &= \exp(iaK_2) \\ &= \exp\left[\frac{-ia}{2}(q_1p_1 + p_2q_2)\right], \\ \mathcal{U}^{(0)}(0, b) &= \exp(ibL_1) \\ &= \exp\left[\frac{ib}{2}(q_1p_1 - q_2p_2)\right].\end{aligned}\tag{3.16}$$

The two factors $\mathcal{U}^{(0)}(a, 0)$ and $\mathcal{U}^{(0)}(0, b)$ commute and may be written in either order, since according to eq. (3.2c) the noncompact generators K_2 and L_1 commute.

Finally one can easily calculate the symplectic matrix $S^{(0)}(a, b) \in Sp(4, \mathbb{R})$, corresponding to the metaplectic operator $\mathcal{U}^{(0)}(a, b)$, by using eq. (2.8). The result is:

$$\begin{aligned} \mathcal{U}^{(0)}(a, b)^{-1} \xi \mathcal{U}^{(0)}(a, b) &= S^{(0)}(a, b) \xi, \\ S^{(0)}(a, b) &= \text{diag.} \left(e^{(a-b)/2}, e^{(a+b)/2}, e^{-(a-b)/2}, e^{-(a+b)/2} \right). \end{aligned} \quad (3.17)$$

Now we can clarify that the particular choice (3.13) was dictated by the desire to have $S^{(0)}(a, b)$ diagonal. This element of $Sp(4, \mathbb{R})$ describes independent reciprocal scalings of the standard quadrature components of each mode. This amounts to showing geometrically that it is possible to diagonalise every $P \in \Pi$ using conjugation by $U(2)$.

We illustrate our classification scheme of two-mode squeezing transformations by giving two examples. The extensively studied Caves-Schumaker [6] transformation uses the operator

$$\mathcal{U}^{(\text{C-S})}(z) = \exp \left(z a_1^\dagger a_2^\dagger - z^* a_1 a_2 \right). \quad (3.18)$$

By appearance, this attempts to involve or entangle the two modes maximally. In our notation this squeezing transformation corresponds to the generator combination

$$\begin{aligned} z a_1^\dagger a_2^\dagger - z^* a_1 a_2 &= i(\vec{k} \cdot \vec{K} + \vec{l} \cdot \vec{L}), \\ \vec{k} &= -2(0, 0, \text{Im } z), \\ \vec{l} &= 2(0, 0, \text{Re } z). \end{aligned} \quad (3.19)$$

Thus the invariant parameters a and b have values

$$a = 2|z|, \quad b = 0. \quad (3.20)$$

The Caves-Schumaker squeezing transformations, and their $U(2)$ conjugates, all taken together, form a one-parameter family or one-dimensional line, in the $a - b$ octant. In that sense they are a set of measure zero.

Another interesting case is a squeezing transformation that refers essentially to a single mode but masquerades as a two-mode transformation:

$$\mathcal{U}(z; \alpha, \beta) = \exp \left[z(\alpha^* a_1^\dagger + \beta^* a_2^\dagger)^2 - z^*(\alpha a_1 + \beta a_2)^2 \right],$$

$$|\alpha|^2 + |\beta|^2 = 1. \quad (3.21)$$

After some simple algebra we find

$$\mathcal{U}(z; \alpha, \beta) = \exp \left[i(\vec{k} \cdot \vec{K} + \vec{l} \cdot \vec{L}) \right],$$

$$\vec{k} + i\vec{l} = 2z \left(-i(\alpha^{*2} - \beta^{*2}), (\alpha^{*2} + \beta^{*2}), 2i\alpha^* \beta^* \right). \quad (3.22)$$

The associated invariants and parameters are

$$\mathfrak{S}_1 = 16|z|^4, \quad \mathfrak{S}_2 = 8|z|^2,$$

$$a = b = 2|z|. \quad (3.23)$$

These equivalence classes thus lie along the line $a = b$ in the octant, again a one-parameter family of zero measure. Our results are depicted in Figure 1.

We note here that the size of an equivalence class depends sensitively upon the point (a, b) . Since for $a \neq 0, b \neq 0$ and $a \neq b$ none of the generators of $U(2)$ or a linear combination of them commute with $aK_2 + bL_1$, we have a full four parameter equivalence class. For the cases $a \neq 0, b = 0$, and $a = b$, respectively, the vanishing of the commutators $[J_2, K_2]$ and $[Q + J_3, K_2 + L_1]$ leads to reduction of the dimensionality of the equivalence class to three.

We conclude this Section with a few comments. Each point (a, b) in the octant denotes an equivalence class of squeezing transformations, whose dependences on a and b would be of physical significance and would show up in a variety of $U(2)$ -invariant properties. The two-mode transformations so far discussed in the literature lie basically along the two lines shown in Figure 1. In this sense, *most* of the intrinsically distinct two-mode transformations, their effects on various states, etc., remain to be explored. Those equivalence classes (a, b) for which $a > b$ involve the two modes in an essential way. We may say purely qualitatively that the *distance* of the point (a, b) from the line $a = b$, or perhaps better the expression $(1 - b/a)$, is a measure of the extent to which two independent modes are involved in the transformation. In this sense, as remarked earlier, the Caves-Schumaker transformations involve two modes maximally.

IV. SQUEEZED COHERENT AND THERMAL STATES FOR TWO MODES

The general two mode coherent state with complex two-component displacement $\tilde{\alpha} = (\alpha_1, \alpha_2)$ is defined by

$$\begin{aligned} |\tilde{\alpha}\rangle &= \exp\left(\tilde{\alpha} \cdot \tilde{a}^\dagger - \tilde{\alpha}^* \cdot \tilde{a}\right) |0, 0\rangle \\ &= \exp\left(-\frac{1}{2}|\alpha_1|^2 - \frac{1}{2}|\alpha_2|^2\right) \exp\left(\alpha_1 a_1^\dagger + \alpha_2 a_2^\dagger\right) |0, 0\rangle. \end{aligned} \quad (4.1)$$

For this state the means of the quadrature components ξ_a do not vanish in general:

$$\langle \tilde{\alpha} | \xi | \tilde{\alpha} \rangle = \sqrt{2} (\text{Re } \alpha_1, \text{Re } \alpha_2, \text{Im } \alpha_1, \text{Im } \alpha_2)^T. \quad (4.2)$$

The variance matrix is however independent of $\tilde{\alpha}$:

$$V(|\tilde{\alpha}\rangle) = V(|\tilde{0}\rangle) = \frac{1}{2} \mathbf{1}_{4 \times 4}. \quad (4.3)$$

The most general squeezed coherent state is obtained by applying $\mathcal{U}(P)$ for some $P \in \Pi \subset Sp(4, \mathbb{R})$ to $|\tilde{\alpha}\rangle$ for some $\tilde{\alpha}$. This $\mathcal{U}(P)$ is conjugate, via some $U(2)$ element, to $\mathcal{U}^{(0)}(a, b)$ for some a, b . Now the effect of a $U(2)$ transformation on $|\tilde{\alpha}\rangle$ is to give us another coherent state $|\tilde{\alpha}'\rangle$, $\tilde{\alpha}'$ being the $U(2)$ transform of $\tilde{\alpha}$. But the variance matrix is in any case $\tilde{\alpha}$ -independent. To examine the $U(2)$ -invariant squeezing condition, therefore, it suffices to examine the particular class of squeezed coherent states

$$|\tilde{\alpha}; a, b\rangle = \mathcal{U}^{(0)}(a, b) |\tilde{\alpha}\rangle. \quad (4.4)$$

From eqs (2.12, 3.17), the calculation of the variance matrix for this state is trivial, and it is in fact diagonal:

$$\begin{aligned} V(|\tilde{\alpha}; a, b\rangle) &= S^{(0)}(a, b) V(|\tilde{\alpha}\rangle) S^{(0)}(a, b) \\ &= \frac{1}{2} S^{(0)}(2a, 2b) \\ &= \frac{1}{2} \text{diag.} \left(e^{(a-b)}, e^{(a+b)}, e^{(b-a)}, e^{-(a+b)} \right). \end{aligned} \quad (4.5)$$

Since a and b are nonnegative, and in addition $a + b > 0$, we see that the least eigenvalue of this variance matrix is

$$\ell(V(|\tilde{\alpha}; a, b\rangle)) = \frac{1}{2} e^{-(a+b)} < \frac{1}{2}. \quad (4.6)$$

These states are thus always squeezed.

If we apply any passive $U(2)$ transformation $S(X, Y)$ to any one of the states $|\tilde{\alpha}; a, b\rangle$ defined above, the variance matrix will in general change as $V \rightarrow V' = S(X, Y) V S(X, Y)^T$; but its eigenvalue spectrum, and in particular $\ell(V)$, remains unaltered. Thus all the states symbolically written as $\mathcal{U}(U(2)) |\tilde{\alpha}; a, b\rangle$, for various $U(2)$ elements, are squeezed to the same extent as $|\tilde{\alpha}; a, b\rangle$, and have $\ell(V)$ given by eq. (4.6).

The Schrödinger wave functions for the subfamily of squeezed coherent states (4.4) are particularly simple, since they are products of single mode squeezed coherent state wave-functions:

$$\begin{aligned} \langle q'_1, q'_2 | \tilde{\alpha}; a, b \rangle &= \psi^{(0)}(q'_1; \alpha_1, a - b) \psi^{(0)}(q'_2; \alpha_2, a + b) \\ \psi^{(0)}(q'; \alpha, a) &= \frac{e^{-a/4}}{\pi^{1/4}} \exp \left[i\alpha \operatorname{Im} \alpha - \frac{1}{2} \left(q' e^{-a/2} - \sqrt{2}\alpha \right)^2 \right]. \end{aligned} \quad (4.7)$$

(For a general state $\mathcal{U}(U(2)) |\tilde{\alpha}; a, b\rangle$, we do not expect such a product form). When we set $b = 0$ (Caves-Schumaker limit), both factors show the same amount of squeezing; while when we set $a = b$ (essentially single mode situation) we see squeezing only in the factor referring to the second mode. These features are as we would have expected.

The next example we look at is the case of a two-mode thermal state subjected to squeezing. The motivation in making this choice is that the starting density operator is explicitly $U(2)$ invariant. The normalized density operator corresponding to inverse temperature $\beta = \hbar\omega/kT$ is described in the Fock and Schrödinger representations by:

$$\begin{aligned} \rho_0(\beta) &= (1 - e^{-\beta})^2 \exp \left[-\beta(a_1^\dagger a_1 + a_2^\dagger a_2) \right] \\ &= (1 - e^{-\beta})^2 \sum_{n_1, n_2=0}^{\infty} e^{-\beta(n_1+n_2)} |n_1, n_2\rangle \langle n_1, n_2|; \end{aligned} \quad (4.8a)$$

$$\begin{aligned} \rho_0(q'_1, q'_2, q''_1, q''_2; \beta) &= \frac{2}{\pi} \tanh^2 \frac{\beta}{2} \exp \left[-\frac{1}{2} \left(\tanh \frac{\beta}{2} + \coth \frac{\beta}{2} \right) (q_1'^2 + q_2'^2 + q_1''^2 + q_2''^2) \right. \\ &\quad \left. - \left(\tanh \frac{\beta}{2} - \coth \frac{\beta}{2} \right) (q_1' q_1'' + q_2' q_2'') \right]; \end{aligned} \quad (4.8b)$$

with $U(2)$ invariance expressed by

$$e^{i\theta Q} \rho_0(\beta) e^{-i\theta Q} = e^{i\vec{\alpha} \cdot \vec{J}} \rho_0(\beta) e^{-i\vec{\alpha} \cdot \vec{J}} = \rho_0(\beta). \quad (4.9)$$

Therefore it suffices to examine the properties of the density operator obtained by conjugating $\rho_0(\beta)$ with $\mathcal{U}^{(0)}(a, b)$:

$$\rho(\beta; a, b) = \mathcal{U}^{(0)}(a, b) \rho_0(\beta) \mathcal{U}^{(0)}(a, b)^{-1}. \quad (4.10)$$

The most general squeezed thermal state is evidently

$$\mathcal{U}(U(2)) \rho(\beta; a, b) \mathcal{U}(U(2))^{-1}, \quad (4.11)$$

but this has the same squeezing properties as $\rho(\beta; a, b)$.

For the thermal state $\rho_0(\beta)$ the variance matrix is well known [10]:

$$V(\rho_0(\beta)) = \frac{1}{2} \coth \frac{\beta}{2} \mathbf{1}_{4 \times 4} \quad (4.12)$$

Therefore for the particular set of squeezed thermal states (4.10), we have diagonal variance matrices:

$$\begin{aligned} V(\rho(\beta; a, b)) &= S^{(0)}(a, b) V(\rho_0(\beta)) S^{(0)}(a, b)^T \\ &= \frac{1}{2} \coth \frac{\beta}{2} S^{(0)}(2a, 2b) \\ &= \frac{1}{2} \coth \frac{\beta}{2} \text{diag.} \left(e^{(a-b)}, e^{(a+b)}, e^{(b-a)}, e^{-a-b} \right). \end{aligned} \quad (4.13)$$

The least eigenvalue is evidently

$$\ell(V) = \frac{1}{2} \coth \frac{\beta}{2} e^{-(a+b)}, \quad (4.14)$$

so for a given temperature, squeezing sets in when

$$a + b > \ln \coth \frac{\beta}{2}. \quad (4.15)$$

In Figure 1, this region consists of all points in the $a - b$ octant *to the right of* the line $a + b = \ln \coth \beta/2$, which is a line perpendicular to the line $a = b$ and at a distance $\ln \coth(\beta/2)/\sqrt{2}$ from the origin.

V. DETECTION SCHEMES AND ROLE OF $U(2)$ TRANSFORMATIONS

We have so far not specified in any detail the two orthogonal modes of radiation being subjected to squeezing. Let us at this point consider a situation well studied experimentally by the heterodyne detection scheme [19]. Here the two modes differ only slightly in frequency, but are otherwise similar. In this kind of experimental arrangement, what is actually measured is the fluctuation of a certain photocurrent, and this in turn gives the fluctuation or variance of the q -quadrature component of a particular (passive) combination of the original modes. The combinations of a_1 and a_2 that are involved form the one-parameter family

$$a(\psi) = \frac{1}{\sqrt{2}} (a_1 + a_2) e^{-i\psi/2}, \quad 0 \leq \psi < 4\pi. \quad (5.1)$$

This can be regarded as the first component of a $U(2)$ -transformed pair a'_1, a'_2 :

$$\begin{pmatrix} a'_1 \\ a'_2 \end{pmatrix} = \frac{1}{\sqrt{2}} \begin{pmatrix} e^{-i\psi/2} & e^{-i\psi/2} \\ -e^{i\psi/2} & e^{i\psi/2} \end{pmatrix} \begin{pmatrix} a_1 \\ a_2 \end{pmatrix},$$

$$a'_1 \equiv a(\psi). \quad (5.2)$$

The hermitian quadrature component whose fluctuation is measured is

$$\begin{aligned} q(\psi) &= \frac{1}{\sqrt{2}} (a(\psi) + a(\psi)^\dagger) \\ &= \frac{1}{\sqrt{2}} (q_1 + q_2) \cos \frac{\psi}{2} + \frac{1}{\sqrt{2}} (p_1 + p_2) \sin \frac{\psi}{2}. \end{aligned} \quad (5.3)$$

The only experimentally adjustable parameter here is the angle ψ . The family of $U(2)$ elements realised in the heterodyne scheme is thus only the one-parameter set given in eq. (5.2) parametrised by ψ , and belonging to $SU(2)$:

$$U_H(\psi) = \frac{1}{\sqrt{2}} \begin{pmatrix} e^{-i\psi/2} & e^{-i\psi/2} \\ -e^{i\psi/2} & e^{i\psi/2} \end{pmatrix} \in SU(2). \quad (5.4)$$

We notice that this is *not* a one-parameter subgroup of $SU(2)$; in particular even the identity element of the group is not contained here.

With this description of the heterodyne setup in our framework, let us see to what extent it can be used to detect $U(2)$ -invariant squeezing. Now a general two-mode state ρ with

variance matrix V , even if it is squeezed in the intrinsic sense of eq. (2.17), may not be manifestly squeezed. That is, it may happen that $V_{aa} \geq 1/2$ for all $a = 1, \dots, 4$. As our discussion in Section (II) shows, we need to be able to *experimentally* realise a general $U(2)$ -transformation applied to the state ρ , and change its variance matrix to a form where one of its diagonal entries (say the leading one) becomes less than $1/2$. However the heterodyne method is generally unable to do this job for us, as it can only realise the one-parameter subset of $SU(2)$ transformations $U_H(\psi)$ for $0 \leq \psi < 4\pi$.

In the two examples of squeezed coherent states and squeezed thermal states studied in the previous Section, we have a family of states related to each other by conjugation with $U(2)$ for each point in the $a-b$ plane. Each equivalence class has appropriate dimensionality depending upon the point (a, b) as explained in Section III. It turns out that for each (a, b) the heterodyning scheme can detect squeezing in only a one parameter subset of the family of states. Although heterodyning detection covers the whole $a-b$ plane it does not reach all the states corresponding to each point in the $a-b$ plane. For example, in the representative chosen in eqs (4.4, 4.10), for which the variance matrix is already diagonal, the squeezing cannot be detected by this scheme because of the absence of the identity in U_H ! It should be possible to detect squeezing in these states by a suitably modified scheme. We wish to emphasize that there is a definite need to be able to experimentally implement the most general element of $U(2)$. This would allow the experimenter to detect the degree of squeezing unambiguously, if the state is squeezed, without any prior knowledge of the elements of the initial variance matrix.

Having stressed the need to implement arbitrary $U(2)$ transformations on the two modes of radiation in order to reach the proper quadrature to exhibit squeezing, we now describe how it can be achieved in some situations. We discuss two particular cases of the two modes involved, the first when the two modes have the same frequency but different directions of propagation, and the second when the modes have the same frequency and direction of propagation but different polarisations.

The experimental setup for the first case is shown in Figure 2. We achieve an arbitrary

$U(2)$ transformation on the two modes by using a Mach-Zehnder interferometer with two 50:50 beam splitters (BS_1 and BS_2) and appropriate phase shifters [20]. The input modes with annihilation operators a_1 and a_2 are subjected to equal and opposite phase shifts by angles ϕ and $-\phi$, then the modes are mixed in the beam splitter BS_1 , the mixed modes again undergo equal and opposite phase shifts by angles θ and $-\theta$ and further mixing through the beam splitter BS_2 . Finally they undergo unequal phase shifts by angles ψ_1 and ψ_2 . If the annihilation operators at the output are a'_1 and a'_2 then all the above operations when combined are implemented through the transformation

$$\begin{pmatrix} a'_1 \\ a'_2 \end{pmatrix} = \begin{pmatrix} e^{i(\phi+\psi_1)} \cos \theta & -ie^{-i(\phi-\psi_1)} \sin \theta \\ -ie^{i(\phi+\psi_2)} \sin \theta & e^{-i(\phi-\psi_2)} \cos \theta \end{pmatrix} \begin{pmatrix} a_1 \\ a_2 \end{pmatrix} \quad (5.5)$$

relating the two sets of annihilation operators. The above matrix is the most general $U(2)$ transformation matrix. We note here that if $\psi_2 = -\psi_1$ then the transformation matrix is the most general $SU(2)$ transformation. Going from $SU(2)$ to $U(2)$ is just a matter of overall phase and can also be achieved by free propagation.

For the second case when the two modes differ only in polarisation we achieve the arbitrary $U(2)$ transformation by using two quarter wave plates (Q_1, Q_2) and a half wave plate H as shown in Figure 3. The detailed discussion of this set up is given in [21]. It turns out that the configuration Q-H-Q is not the only one but Q-Q-H and H-Q-Q also accomplish the same result, as shown in [21]. We basically have three elements, a quarter wave plate Q_1 , a half wave plate H and a quarter wave plate Q_2 all three of them being coaxially mounted and with their slow axes in the $x - y$ plane making angles of α, β and γ respectively with the $\hat{x}$ axis. The two modes having annihilation operators a_1 and a_2 moving along the $\hat{z}$ direction pass through this arrangement. If the annihilation operators at the output are a'_1 and a'_2 then they are related to the operators at the input by an $SU(2)$ transformation given in terms of α, β and γ . By changing these parameters one can reproduce any desired $SU(2)$ element. As has been pointed out earlier going to $U(2)$ now is just a matter of free propagation.

In both the above cases, by going to the proper $U(2)$ element we can make the squeezing (if present) manifest and bring it to the leading diagonal element of the variance matrix i.e. in the quadrature $q'_1 = \frac{1}{\sqrt{2}}(a'_1 + a_1^{\dagger'})$. The squeezing in this quadrature can now be measured by any standard one mode detection method.

These remarks show on the one hand the way the heterodyning scheme fits into our general analysis, and on the other hand the need to devise new schemes capable of realising all elements of $U(2)$, tailored to the definition of the two modes involved.

VI. CONCLUDING REMARKS

We have presented a classification scheme for two-mode squeezing transformations, based on the structure of the real four dimensional symplectic group $Sp(4, \mathbb{R})$, and the separation of its elements into passive (compact) and active (noncompact) types. The structure and action of the maximal compact subgroup $U(2)$ in $Sp(4, \mathbb{R})$, and the $U(n)$ -invariant squeezing criterion formulated elsewhere for a general n -mode system, have guided our considerations. All our work is in the metaplectic unitary representation of $Sp(4, \mathbb{R})$; and the local isomorphism $Sp(4, \mathbb{R}) \approx SO(3, 2)$ has led to a convenient geometric picturization of the situation.

As emphasized in Section III the squeezing transformations $\mathcal{U}(P)$, $P \in \Pi \subset Sp(4, \mathbb{R})$, do not form a subgroup of $Sp(4, \mathbb{R})$. The breakup of these transformations into equivalence classes, based on the effect of conjugation by elements of $U(2)$ is the only natural available classification procedure. This is because the definition of equivalence classes for any set of objects has to be based on an equivalence relation defined on that set. Thus, we have treated two elements $P, P' \in \Pi$ as intrinsically equivalent if

$$P' = S(X, Y) P S(X, Y)^T \quad \text{for some} \quad X - iY \in U(2). \quad (6.1)$$

It should however be realised that the detailed effects of action by $\mathcal{U}(P)$ and $\mathcal{U}(P')$ on a *general* initial two mode state ρ_0 , as seen in the changes caused in the variance matrix $V(\rho_0)$, need not be identical. Since this is a subtle and important point we spell it out in detail. Starting from a general state ρ_0 , action by a squeezing transformation leads to a new state

$$\rho = \mathcal{U}(P) \rho_0 \mathcal{U}(P)^{-1}. \quad (6.2)$$

As seen in Section III, any $\mathcal{U}(P)$ is expressible in terms of a representative element $\mathcal{U}^{(0)}(a, b)$ as

$$\mathcal{U}(P) = \mathcal{U}(S(X, Y)) \mathcal{U}^{(0)}(a, b) \mathcal{U}(S(X, Y))^{-1} \quad (6.3)$$

for suitable $X - iY \in U(2)$. Therefore we have

$$\rho = \mathcal{U}(S(X, Y)) \mathcal{U}^{(0)}(a, b) \mathcal{U}(S(X, Y))^{-1} \rho_0 \mathcal{U}(S(X, Y)) \mathcal{U}^{(0)}(a, b)^{-1} \mathcal{U}(S(X, Y))^{-1}. \quad (6.4)$$

This leads by eq. (2.12) to the relation

$$V(\rho) = S(X, Y) S^{(0)}(a, b) S(X, Y)^T V(\rho_0) S(X, Y) S^{(0)}(a, b) S(X, Y)^T \quad (6.5)$$

between the two variance matrices. Now the first and last factors on the right hand side here have no influence on the spectrum, and so on the least eigenvalue, of $V(\rho)$. Therefore the squeezed or nonsqueezed nature of ρ is actually determined by the least eigenvalue of the matrix

$$S(X, Y)^T V(\rho) S(X, Y) = S^{(0)}(a, b) S(X, Y)^T V(\rho_0) S(X, Y) S^{(0)}(a, b). \quad (6.6)$$

But now the right hand side is in *general* dependent not only on the invariant parameters a, b but also on X, Y . In the examples studied in Section IV, namely where ρ_0 is a coherent state or an isotropic thermal state, $V(\rho_0)$ happens to be a multiple of the identity matrix, so that on the right hand side of eq. (6.6) the dependence on X, Y cancels. But this need not happen in general. Thus for instance if we take for ρ_0 an anisotropic thermal state with unequal temperatures for the two modes, we have only $U(1) \times U(1)$, rather than $U(2)$, invariance for this ρ_0 ; so the least eigenvalue $\ell(V(\rho))$ of $V(\rho)$ will depend on a, b and on two out of the four $U(2)$ parameters present in $X - iY$. One can easily convince oneself that the only situation where $S(X, Y)^T V(\rho_0) S(X, Y) = V(\rho_0)$ independent of X and Y is when $V(\rho_0)$ is a multiple of the unit matrix; and the isotropic thermal states do reproduce all such

cases. Therefore a more detailed study of the effect of a general squeezing transformation on initial states ρ_0 with nontrivial $V(\rho_0)$ is of considerable interest.

A related important question is the following: Let us take two squeezing transformations $P_1, P_2 \in \Pi$ belonging to equivalence classes $(a_1, b_1), (a_2, b_2)$ respectively, which could coincide. The product $P_1 P_2$ will in general be of the form $S(X, Y) P$ with $S(X, Y) \in U(2)$ and $P \in \Pi$. If P belongs to the equivalence class (a, b) we wish to determine this class in terms of P_1 and P_2 . Using the fact $Tr(P^2) = Tr((P_1 P_2)(P_1 P_2)^T)$ and eq. (3.15) we arrive at the following relations

$$\begin{aligned} 2[\cosh(a - b) + \cosh(a + b)] &= Tr((P_1 P_2)(P_1 P_2)^T) \\ 2[\cosh 2(a - b) + \cosh 2(a + b)] &= Tr(((P_1 P_2)(P_1 P_2)^T)^2) \end{aligned} \quad (6.7)$$

which can be solved to find (a, b) . We note here that (a, b) do depend not only upon (a_1, b_1) and (a_2, b_2) but also on the actual elements chosen from each of these classes. So we do not have a notion of class multiplication among these equivalence classes.

Finally we call attention to our considerations in Section V and to the need for being able to experimentally implement or realize general passive elements of the subgroup $U(2)$ of $Sp(4, \mathbb{R})$ for each given choice of the independent modes in a two-mode system. Once this is achieved, for any given state we can bring out in an explicit or manifest fashion its squeezing nature (provided it is squeezed) by altering its variance matrix and making the least eigenvalue appear in the leading position on the diagonal. This also means that we would be able to experimentally measure the fluctuation in the quadrature variable isolating the least eigenvalue. As extensively discussed elsewhere, these considerations which exploit the richness of the geometry underlying the symplectic group do not require complete diagonalization of the variance matrix at all [10].

In the following paper we shall examine $U(2)$ -invariant properties of two mode squeezed states which go beyond the level of second order moments of the quadrature operators.

REFERENCES

* email arvind@physics.iisc.ernet.in

† Also at Jawaharlal Nehru Centre for Advanced Scientific Research, Indian Institute of Science Campus, Bangalore - 560 012, India.

- [1] D. Stoler, Physical Review D **1**, 3217 (1970); H. P. Yuen, Physical Review A **13**, 2226 (1976); D. F. Walls, Nature **306**, 141 (1983); J. N. Hollenhorst, Physical Review D **19**, 1669 (1979).
- [2] R. E. Slusher, L. W. Hollberg, B. Yurke, J. C. Mertz, and J.F. Valley, Physical Review Letters **55**, 2409 (1985); R. M. Shelby, M. D. Levenson, S. H. Perlmutter, R. G. DeVoe, and D. F. Walls, Physical Review Letters **57**, 691 (1986); Ling-An Wu, J. H. Kimble, J. L. Hall, and H. Wu, Physical Review Letters **57**, 2520 (1986); M. W. Maeda, P. Kumar, and J. H. Shapiro, Optics Letters **12**, 161 (1987).
- [3] B. L. Schumaker, S. H. Perlmutter, R. M. Shelby and M. D. Levenson, Physical Review Letters **58**, 357 (1987); M. G. Raizen, L. A. Oroscio, M. Xiao, T. L. Boyd and H. J. Kimble, Physical Review Letters **59**, 198 (1987); A. Heidmann, R. J. Horowicz, S. Reynaud, E. Giacobino, C. Fabre, and G. Camy, Physical Review Letters **59**, 2555 (1987); Z. Y. Ou, S. F. Pereira, H. J. Kimble and K. C. Peng, Physical Review Letters **68**, 3663 (1992).
- [4] P. Kumar, O. Aytur and J. Huang, Physical Review Letters **64**, 1015 (1990); A. Sizmann, R. J. Horowicz, E. Wagner and G. Leuchs, Optics Communications **80**, 138 (1990); M. Rosenbluh and R. M. Shelby, Physical Review Letters **66**, 153 (1991); K. Bergman and H. A. Haus, Optics Letters **16**, 663 (1991); E. S. Polzik, R. C. Carri and H. J. Kimble, Physical Review Letters **68**, 3020 (1992).
- [5] For recent reviews, see, R. Loudon and P. L. Knight, Journal of Modern Optics **34**, 709 (1987); M. C. Teich and B. E. A. Saleh, Quantum Optics **1**, 153 (1990), S. Rey-

- naud, A. Heidmann, E. Giacobino and C. Fabre, in *Progress in Optics*, Vol. 30, ed. E. Wolf (North-Holland, Amsterdam, 1992); C. Fabre, *Physics Reports* **219**, 215 (1992); H. J. Kimble, *Physics Reports* **219**, 227 (1992).
- [6] C. M. Caves and B. L. Schumaker, *Physical Review A* **31**, 3068 (1985); B. L. Schumaker and C. M. Caves, *Physical Review A* **31**, 3093 (1985); B. L. Schumaker, *Physics Reports* **135**, 317 (1986); C. M. Caves, *Physical Review D* **23**, 1693 (1981).
- [7] G. S. Agarwal, *Physical Review Letters* **57**, 827 (1986); G. S. Agarwal, *Journal of the Optical Society of America B* **5**, 1940 (1988).
- [8] A. K. Eckert and P. L. Knight, *American Journal Physics* **57**, 692 (1989); C. M. Caves, C. Zhu, G. J. Milburn and W. Schleich, *Physical Review A* **43**, 3854 (1991); M. Selvadhoray, M. Sanjay Kumar and R. Simon, preprint I. M. Sc. - 93/22 (1993).
- [9] C. K. Hong and L. Mandel, *Physical Review Letters* **54**, 323 (1985); *Physical Review A* **32**, 974 (1985); C. C. Gerry and P. J. Moyer, *Physical Review A* **38**, 5665 (1988); M. S. Kim, F. A. M. de Oliveira and P. L. Knight, *Physical Review A* **40**, 2494 (1989); J. J. Gong and P. K. Aravind, *Physical Review A* **46**, 1586 (1992); M. Hillery, *Physical Review A* **36**, 3796 (1987).
- [10] R. Simon, N. Mukunda and B. Dutta, *Physical Review A* **49**, 1567 (1994).
- [11] T. F. Jordan, *Linear operators in quantum mechanics*, (John Wiley, New York, 1974); G. Lion and M. Vergne, *The Weil Representation, Maslov Index, and Theta series*, (Birkhauser, Basel, 1980).
- [12] A. Weil, *Acta Mathematica* **111**, 143 (1964); V. Guillemin and S. Sternberg, *Symplectic Techniques in Physics*, Cambridge University Press, (1984); R. Simon and N. Mukunda, *The two-dimensional symplectic and metaplectic groups and their universal cover in Symmetries in Science VI: From the Rotation Group to Quantum Algebras*, B. Gruber (ed.), Plenum Press (to appear).

- [13] N. Mukunda, Current Science **59**, 1135 (1990); B. Dutta, N. Mukunda, R. Simon and A. Subramaniam, Journal of the Optical Society of America B **10**, 253 (1993).
- [14] The connection between the groups $Sp(4, \mathfrak{R})$ and $SO(3, 2)$ has been noticed and exploited in many physical contexts for a very long time. These include the infinite component relativistic wave equations constructed by Majorana, the new Dirac equation for a relativistic point particle with two internal harmonic oscillator degrees of freedom, the theory of relativistic objects with internal structure and classification of anisotropic Schell model beams in optics. See E. Majorana, Nuovo Cimento **9**, 335 (1932); P. A. M. Dirac, Journal of Mathematical Physics **4**, 901 (1963); N. Mukunda, Physical Review **183**, 1486 (1969); E. C. G. Sudarshan and N. Mukunda, Physical Review D **1**, 576 (1970); P. A. M. Dirac, Proceedings of the Royal Society A **322**, 435 (1971); *ibid* A **328**, 1 (1972); N. Mukunda, H. van Dam and L. C. Biedenharn, *Relativistic Models of Extended Hadrons obeying a mass spin trajectory*, Lecture Notes in Physics, Vol. 165, Springer-Verlag 1982; R. Simon, E. C. G. Sudarshan and N. Mukunda, Physical Review A **31**, 2419 (1985); Physical Review A **36**, 3868 (1987). A recent paper exploiting these ideas for squeezing problems is D. Han, Y. S. Kim, M. E. Noz and L. Yeh, Journal of Mathematical Physics **34**, 5493 (1993).
- [15] R. Gilmore, *Lie Groups, Lie Algebras and some of their applications*, John Wiley and Sons, New York USA(1974)
- [16] C. L. Siegel, American Journal of Mathematics **LXV**, 1 (1943), reprinted as C. L. Siegel *Symplectic Geometry*, Academic Press (New York) 1964.
- [17] Arvind, B. Dutta, N. Mukunda and R. Simon, *The real symplectic groups in quantum mechanics and optics*, IISc preprint in preparation.
- [18] E. P. Wigner, *Symmetries and Reflections*, Indiana University Press, 1967.
- [19] H. P. Yuen and J. H. Shapiro, IEEE Transactions on Information Theory **24**, 657 (1978);

- M. D. Levenson, R. M. Shelby, A. Aspect, M. Reid and D. F. Walls, *Physical Review A* **32**, 1550 (1985); M. J. Collett, R. Loudon and C. W. Gardiner, *Journal of Modern Optics* **34**, 881 (1987).
- [20] B. Yurke, S. L. McCall and J. R. Klauder, *Physical Review A* **33**, 4033 (1986); R. A. Campos, B. E. A. Saleh and M. C. Teich, *Physical Review A* **40**, 1371 (1989); R. F. Bishop and A. Vourdas, *Zeitschrift für Physik* **71**, 527 (1988); C. T. Lee, *Physical Review A* **42**, 4193 (1990); D. Han, Y. S. Kim and M. E. Noz, *Physical Review A* **41**, 6233 (1990).
- [21] R. Simon and N. Mukunda, *Physics Letters A* **143**, 165 (1990);

FIGURES




FIG. 1. Equivalence classes of two-mode squeezing transformations, Caves-Schumaker (C-S) and single mode limits, squeezed thermal region.




FIG. 2. Mach-Zehnder Interferometer implementing arbitrary $U(2)$ transformation on two modes at the same frequency but differing in their direction of propagation. BS_1 and BS_2 are 50:50 beam splitters and thick lines are phase shifters by angles indicated. a_1, a_2 are the annihilation operators at the input port and a'_1, a'_2 are the annihilation operators at the output port




FIG. 3. Implementation of arbitrary $SU(2)$ element on two modes with the same frequency and directions of propagation, but different polarisations. Here Q_1 , Q_2 are quarter wave plates and H is the half wave plate. α , β , and γ are the angles which the slow axes of Q_1 , H and Q_2 make with the $\hat{x}$ axis respectively. a_1 , a_2 are the annihilation operators at the input port and a'_1 , a'_2 are the annihilation operators at the output port