

## NONEXTENDED QUADRATIC FORMS OVER POLYNOMIAL RINGS OVER POWER SERIES RINGS

RAMAN PARIMALA

**ABSTRACT.** If  $R$  is a complete discrete valuation ring, then every quadratic space over  $R[T]$  is extended from  $R$ . We here show by an example that a corresponding result for higher-dimensional complete regular local rings is not valid.

It was proved in [3] that if  $R$  is any complete discrete valuation ring, every quadratic space over  $R[T]$  is extended from  $R$ . We show in this note that if  $R = \mathbf{R}[[X, Y]]$ , there exist anisotropic quadratic spaces over  $R[T]$  which are not extended from  $R$ . This is in contrast to a result of Mohan Kumar and Lindel in the linear case [2, Theorem 5.1, p. 150].

Let  $R = \mathbf{R}[[X, Y]]$ ,  $\mathbf{R}$  denoting the field of real numbers. Let  $\mathbf{H}$  be the quaternions over  $\mathbf{R}$  and  $A = \mathbf{H}[[X, Y]]$  the ring of formal power series in the variables  $X$  and  $Y$  over  $\mathbf{H}$ . We have an  $A[T]$ -linear map  $A[T]^2 \xrightarrow{\eta} A[T]$  defined by  $(1, 0) \rightarrow XT + i$ ,  $(0, 1) \rightarrow YT + j$  which is clearly a surjection. Let  $P$  be the kernel of  $\eta$ .

**PROPOSITION.** *The module  $P$  is nonfree projective over  $A[T]$ .*

**PROOF.** The first projection of  $A[T]^2$  onto  $A[T]$  maps  $P$  isomorphically onto the left ideal  $\mathfrak{A}$  of  $A[T]$  generated by  $1 + Y^2T^2$  and  $1 + (iX + jY)T - kXYT^2$  [4, p. 143]. We prove that  $P$  is not free by showing that  $\mathfrak{A}$  is not principal. We note first that  $\mathfrak{A}$  is not the unit ideal since it is generated modulo  $X$  by  $1 + jYT$  which is not a unit in  $\mathbf{H}[[Y]][T]$ . Suppose  $\mathfrak{A}$  is principal, generated by  $f$ . Then  $\deg_T f < 2$ . If  $\deg_T f = 0$ , then it follows that  $\mathfrak{A}$  is the unit ideal, which is not the case. If  $\deg_T f = 2$ , then  $1 + Y^2T^2$  and  $1 + (iX + jY)T - kXYT^2$  are unit left multiples of each other, which is again not possible. Let  $\deg_T f = 1$  and  $f = a + bT$ ,  $a, b \in \mathbf{H}[[X, Y]]$ . Then  $a$  is a unit and we assume  $a = 1$  so that  $f = 1 + bT$ ,  $b \in \mathbf{H}[[X, Y]]$ . We then have

$$1 + Y^2T^2 = (1 + cT)(1 + bT),$$

$$1 + (iX + jY)T - kXYT^2 = (1 + dT)(1 + bT).$$

From the first equation we get  $c = -b$ ,  $-b^2 = Y^2$ . This implies that  $b = \lambda Y$ ,  $\lambda \in \mathbf{H}[[X, Y]]$ . From the second equation we get  $d + b = iX + jY$ ,  $db = -kXY$  so that we have

$$(iX + jY)\lambda = -(kX + Y).$$

---

Received by the editors September 4, 1980.

AMS (MOS) subject classifications (1970). Primary 15A63, 18F25.

Key words and phrases. Projective modules, quadratic spaces, extendibility.

© 1981 American Mathematical Society  
0002-9939/81/0000-0502/\$01.50

If  $\lambda = \lambda_0 + \lambda_1 X + \lambda_2 Y + \dots$ ,  $\lambda_i \in \mathbf{H}$ , we have  $i\lambda_0 = -k$ ,  $j\lambda_0 = -1$ , a contradiction, which proves the proposition.

The reduced norm on  $P$  [1, Theorem 2.1] gives rise to a quadratic space of rank 4 and discriminant 1 over  $R[T]$ . This space is anisotropic and not extended from  $R$  (in fact indecomposable) in view of [1, Theorem 4.6].

#### REFERENCES

1. M. A. Knus, M. Ojanguren and R. Sridharan, *Quadratic forms and Azumaya algebras*, J. Reine Angew. Math. **303/304** (1978), 231–248.
2. T. Y. Lam, *Serre's conjecture*, Lecture Notes in Math., vol. 635, Springer-Verlag, Berlin and New York, 1978.
3. Raman Parimala, *Quadratic forms over polynomial rings over Dedekind domains*, Amer. J. Math. **100** (1978), 913–928.
4. S. Parimala and R. Sridharan, *Projective modules over polynomial rings over division rings*, J. Math. Kyoto Univ. **15** (1975), 129–148.

TATA INSTITUTE OF FUNDAMENTAL RESEARCH, HOMI BHABHA ROAD, BOMBAY 400 005, INDIA