

CpG-containing oligodeoxynucleotides as new generation adjuvants in DNA and protein vaccines

Since ages, Freund's Complete Adjuvant (FCA) has been used by immunologists to boost the immune response in experimental animals. Due to inflammatory side effects however, use of FCA is not permitted in humans and search is on for a better adjuvant. Recently several reports have shown that bacterial DNA acts as potent adjuvant, promoting a T-helper 1 (Th1) immune response. An immune response can be classified as a Th1 type characterized by the generation of cytokines like interferon gamma (IFN- γ) and interleukin-2 (IL-2), or a Th2 type in which cytokines like IL-4, IL-5, IL-10 and IL-13 are produced. Th1 cytokines influence the immune response and push it towards the generation of cell mediated immune (CMI) responses like cytotoxic T cells (CTL)

and DTH responses, as well as a switch over of immunoglobulin isotype to IgG2a. Th2 cytokines tilt the immune response towards production of other isotypes of immunoglobulins and block CMI responses. In many disease conditions, one of the two (Th1 or Th2) responses may impart protection while the other type may not be useful or even enhance the susceptibility of the host to the infection. For instance, in tuberculosis and AIDS, a Th1 immune response appears to be protective. Individuals infected with HIV may remain healthy as long as they have a Th1 type of immune response to the viral antigens. Symptoms of AIDS appear in these persons as their immune response shifts from a Th1 to a Th2 type (Clerici and Shearer 1993). The reason for this shift is not known. In this context, it is interesting that the use of bacterial DNA as adjuvant specifically promotes a Th1 response. These indications came as a spin off from the work being done on DNA vaccines where the gene encoding a given immunogenic protein is incorporated in a suitable vector and the modified vector is used as vaccine (Robinson and Torres 1997). In these experiments, administration of vector alone, as control, also resulted in potentiation of the immune response (Leclerc *et al* 1997). The effect was tracked down to the presence of CpG sequences in bacterial DNA. CpG sequences are also present in mammalian DNA but are heavily methylated and therefore not effective. Insect DNAs which have non-methylated CpG sequences are also potent adjuvants (Sun *et al* 1998). Synthetic oligodeoxynucleotides containing CpG sequences (CpG ODN) have been shown to be effective adjuvants when administered along with an antigen and tilt the immune response towards a Th1 type (Chu *et al* 1997). Davis *et al* (1998) recently demonstrated that immunization of mice with Hepatitis B surface antigen along with CpG ODN resulted in a strong Th1 response (antibody of IgG2a isotype and CTLs) but use of alum adjuvant with the same antigen resulted in a Th2 response (antibodies of IgG1 type and no CTLs). These results may mark a milestone in the evolution of vaccines (Krieg *et al* 1998). CpG sequences may either be incorporated into DNA vaccines or CpG ODNs mixed with protein vaccines to not only boost the immune response, but also promote a Th1 kind of response. The mechanism of action of CpG sequences in generating a Th1 response is not well understood. CpG containing bacterial DNA is known to induce the secretion of TNF and IL-12 by macrophages (Sparwasser *et al* 1997). NK cells generate IFN- γ in response to IL-12. As IFN- γ is known to inhibit a Th2 immune response, its early release may set the stage for a robust Th1 immune response.

References

- Chu R S, Torgoni O S, Krieg A M, Lehmann P V and Harding C V 1997 CpG oligodeoxynucleotides act as adjuvants that switch on T helper 1 immunity; *J. Exp. Med.* **186** 1623–1631
- Clerici M and Shearer G M 1993 TH1 to TH2 switch is a critical step in the etiology of HIV infection; *Immunol. Today* **14** 107–111
- Davis H L, Weeranta R, Waldschmidt T J, Tygrett L, Schorr J and Krieg A M 1998 CpG DNA is a potent enhancer of specific immunity in mice immunized with recombinant hepatitis B surface antigen; *J. Immunol.* **160** 870–876
- Krieg A M, Yi A K, Schorr J and Davis H L 1998 The role of CpG dinucleotides in DNA vaccines; *Trends Microbiol.* **6** 23–27
- Leclerc C, Deriaud E, Rojas M and Whalen R G 1997 The preferential induction of a Th1 immune response by DNA-based immunization is mediated by the immunostimulatory effect of plasmid DNA; *Cell. Immunol.* **179** 97–106
- Robinson H L and Torres C A 1997 DNA vaccines; *Semin. Immunol.* **9** 271–283
- Sparwasser T, Miethke T, Lipford G, Erdmann A, Heeg K and Wagner H 1997 Macrophages sense pathogens via DNA motifs: induction of tumor necrosis factor- α -mediated shock; *Eur. J. Immunol.* **27** 1671–1679
- Sun S, Kishimoto H and Sprent, J 1998 DNA as an adjuvant: capacity of insect DNA and synthetic oligodeoxynucleotides to augment T cell response to specific antigen; *J. Exp. Med.* **187** 1145–1150

RAJIV K SAXENA
SAUBIYA M SIDDIQUI
School of Life Sciences,
Jawaharlal Nehru University,
New Delhi 110 067, India
(Email, rajivksaxena@hotmail.com)

References

- Schlosser I J, Doeringsfeld M R, Elder J F and Arzayus L F 1998 Niche relationships of clonal and sexual fish in a heterogeneous landscape: *Ecology* **79** 953–968
- Vrijenhoek R C 1994 Unisexual fish: model systems for studying ecology and evolution; *Annu. Rev. Ecol. Syst.* **25** 71–96
- Bell, G H 1982 *The Masterpiece of Nature. The Evolution and Genetics of Sexuality* (London: Croom Helm)

RENEE M BORGES
*Centre for Ecological Sciences,
Indian Institute of Science,
Bangalore 560 012, India
(Email, renee@ces.iisc.ernet.in)*