



**Fermi National Accelerator Laboratory**

**FERMILAB-Pub-98/095-E**

**D0**

## **A Search for Heavy Pointlike Dirac Monopoles**

B. Abbott et al.  
The D0 Collaboration

*Fermi National Accelerator Laboratory  
P.O. Box 500, Batavia, Illinois 60510*

March 1998

Submitted to *Physical Review Letters*

## **Disclaimer**

*This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, expressed or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.*

## **Distribution**

*Approved for public release; further dissemination unlimited.*

## A Search for Heavy Pointlike Dirac Monopoles

B. Abbott,<sup>31</sup> M. Abolins,<sup>27</sup> B.S. Acharya,<sup>46</sup> I. Adam,<sup>12</sup> D.L. Adams,<sup>40</sup> M. Adams,<sup>17</sup>  
 S. Ahn,<sup>14</sup> H. Aihara,<sup>23</sup> G.A. Alves,<sup>10</sup> N. Amos,<sup>26</sup> E.W. Anderson,<sup>19</sup> R. Astur,<sup>45</sup>  
 M.M. Baarmand,<sup>45</sup> L. Babukhadia,<sup>2</sup> A. Baden,<sup>25</sup> V. Balamurali,<sup>35</sup> J. Balderston,<sup>16</sup>  
 B. Baldin,<sup>14</sup> S. Banerjee,<sup>46</sup> J. Bantly,<sup>5</sup> E. Barberis,<sup>23</sup> J.F. Bartlett,<sup>14</sup> A. Belyaev,<sup>29</sup>  
 S.B. Beri,<sup>37</sup> I. Bertram,<sup>34</sup> V.A. Bezzubov,<sup>38</sup> P.C. Bhat,<sup>14</sup> V. Bhatnagar,<sup>37</sup>  
 M. Bhattacharjee,<sup>45</sup> N. Biswas,<sup>35</sup> G. Blazey,<sup>33</sup> S. Blessing,<sup>15</sup> P. Bloom,<sup>7</sup> A. Boehnlein,<sup>14</sup>  
 N.I. Bojko,<sup>38</sup> F. Borcherding,<sup>14</sup> C. Boswell,<sup>9</sup> A. Brandt,<sup>14</sup> R. Brock,<sup>27</sup> A. Bross,<sup>14</sup>  
 D. Buchholz,<sup>34</sup> V.S. Burtovoi,<sup>38</sup> J.M. Butler,<sup>3</sup> W. Carvalho,<sup>10</sup> D. Casey,<sup>27</sup> Z. Casilum,<sup>45</sup>  
 H. Castilla-Valdez,<sup>11</sup> D. Chakraborty,<sup>45</sup> S.-M. Chang,<sup>32</sup> S.V. Chekulaev,<sup>38</sup> L.-P. Chen,<sup>23</sup>  
 W. Chen,<sup>45</sup> S. Choi,<sup>44</sup> S. Chopra,<sup>26</sup> B.C. Choudhary,<sup>9</sup> J.H. Christenson,<sup>14</sup> M. Chung,<sup>17</sup>  
 D. Claes,<sup>30</sup> A.R. Clark,<sup>23</sup> W.G. Cobau,<sup>25</sup> J. Cochran,<sup>9</sup> L. Coney,<sup>35</sup> W.E. Cooper,<sup>14</sup>  
 C. Cretsinger,<sup>42</sup> D. Cullen-Vidal,<sup>5</sup> M.A.C. Cummings,<sup>33</sup> D. Cutts,<sup>5</sup> O.I. Dahl,<sup>23</sup> K. Davis,<sup>2</sup>  
 K. De,<sup>47</sup> K. Del Signore,<sup>26</sup> M. Demarteau,<sup>14</sup> D. Denisov,<sup>14</sup> S.P. Denisov,<sup>38</sup> H.T. Diehl,<sup>14</sup>  
 M. Diesburg,<sup>14</sup> G. Di Loreto,<sup>27</sup> P. Draper,<sup>47</sup> Y. Ducros,<sup>43</sup> L.V. Dudko,<sup>29</sup> S.R. Dugad,<sup>46</sup>  
 D. Edmunds,<sup>27</sup> J. Ellison,<sup>9</sup> V.D. Elvira,<sup>45</sup> R. Engelmann,<sup>45</sup> S. Eno,<sup>25</sup> G. Eppley,<sup>40</sup>  
 P. Ermolov,<sup>29</sup> O.V. Eroshin,<sup>38</sup> V.N. Evdokimov,<sup>38</sup> T. Fahland,<sup>8</sup> M.K. Fatyga,<sup>42</sup> S. Feher,<sup>14</sup>  
 D. Fein,<sup>2</sup> T. Ferbel,<sup>42</sup> G. Finocchiaro,<sup>45</sup> H.E. Fisk,<sup>14</sup> Y. Fisyak,<sup>4</sup> E. Flattum,<sup>14</sup>  
 G.E. Forden,<sup>2</sup> M. Fortner,<sup>33</sup> K.C. Frame,<sup>27</sup> S. Fuess,<sup>14</sup> E. Gallas,<sup>47</sup> A.N. Galyaev,<sup>38</sup>  
 P. Gartung,<sup>9</sup> V. Gavrilov,<sup>28</sup> T.L. Geld,<sup>27</sup> R.J. Genik II,<sup>27</sup> K. Genser,<sup>14</sup> C.E. Gerber,<sup>14</sup>  
 Y. Gershtein,<sup>28</sup> B. Gibbard,<sup>4</sup> S. Glenn,<sup>7</sup> B. Gobbi,<sup>34</sup> A. Goldschmidt,<sup>23</sup> B. Gómez,<sup>1</sup>  
 G. Gómez,<sup>25</sup> P.I. Goncharov,<sup>38</sup> J.L. González Solís,<sup>11</sup> H. Gordon,<sup>4</sup> L.T. Goss,<sup>48</sup>  
 K. Gounder,<sup>9</sup> A. Goussiou,<sup>45</sup> N. Graf,<sup>4</sup> P.D. Grannis,<sup>45</sup> D.R. Green,<sup>14</sup> H. Greenlee,<sup>14</sup>  
 S. Grinstein,<sup>6</sup> P. Grudberg,<sup>23</sup> S. Grünendahl,<sup>14</sup> G. Guglielmo,<sup>36</sup> J.A. Guida,<sup>2</sup> J.M. Guida,<sup>5</sup>  
 A. Gupta,<sup>46</sup> S.N. Gurzhiev,<sup>38</sup> G. Gutierrez,<sup>14</sup> P. Gutierrez,<sup>36</sup> N.J. Hadley,<sup>25</sup> H. Haggerty,<sup>14</sup>  
 S. Hagopian,<sup>15</sup> V. Hagopian,<sup>15</sup> K.S. Hahn,<sup>42</sup> R.E. Hall,<sup>8</sup> P. Hanlet,<sup>32</sup> S. Hansen,<sup>14</sup>  
 J.M. Hauptman,<sup>19</sup> D. Hedin,<sup>33</sup> A.P. Heinson,<sup>9</sup> U. Heintz,<sup>14</sup> R. Hernández-Montoya,<sup>11</sup>  
 T. Heuring,<sup>15</sup> R. Hirosky,<sup>17</sup> J.D. Hobbs,<sup>45</sup> B. Hoeneisen,<sup>1,\*</sup> J.S. Hoftun,<sup>5</sup> F. Hsieh,<sup>26</sup>  
 Ting Hu,<sup>45</sup> Tong Hu,<sup>18</sup> T. Huehn,<sup>9</sup> A.S. Ito,<sup>14</sup> E. James,<sup>2</sup> J. Jaques,<sup>35</sup> S.A. Jerger,<sup>27</sup>  
 R. Jesik,<sup>18</sup> J.Z.-Y. Jiang,<sup>45</sup> T. Joffe-Minor,<sup>34</sup> K. Johns,<sup>2</sup> M. Johnson,<sup>14</sup> A. Jonckheere,<sup>14</sup>  
 M. Jones,<sup>16</sup> H. Jöstlein,<sup>14</sup> S.Y. Jun,<sup>34</sup> C.K. Jung,<sup>45</sup> S. Kahn,<sup>4</sup> G. Kalbfleisch,<sup>36</sup> J.S. Kang,<sup>20</sup>  
 D. Karmanov,<sup>29</sup> D. Karmgard,<sup>15</sup> R. Kehoe,<sup>35</sup> M.L. Kelly,<sup>35</sup> C.L. Kim,<sup>20</sup> S.K. Kim,<sup>44</sup>  
 B. Klima,<sup>14</sup> C. Klopfenstein,<sup>7</sup> J.M. Kohli,<sup>37</sup> D. Koltick,<sup>39</sup> A.V. Kostritskiy,<sup>38</sup> J. Kotcher,<sup>4</sup>  
 A.V. Kotwal,<sup>12</sup> J. Kourlas,<sup>31</sup> A.V. Kozelov,<sup>38</sup> E.A. Kozlovsky,<sup>38</sup> J. Krane,<sup>30</sup>  
 M.R. Krishnaswamy,<sup>46</sup> S. Krzywdzinski,<sup>14</sup> S. Kuleshov,<sup>28</sup> S. Kunori,<sup>25</sup> F. Landry,<sup>27</sup>  
 G. Landsberg,<sup>14</sup> B. Lauer,<sup>19</sup> A. Leflat,<sup>29</sup> H. Li,<sup>45</sup> J. Li,<sup>47</sup> Q.Z. Li-Demarteau,<sup>14</sup>  
 J.G.R. Lima,<sup>41</sup> D. Lincoln,<sup>14</sup> S.L. Linn,<sup>15</sup> J. Linnemann,<sup>27</sup> R. Lipton,<sup>14</sup> Y.C. Liu,<sup>34</sup>  
 F. Lobkowicz,<sup>42</sup> S.C. Loken,<sup>23</sup> S. Lökös,<sup>45</sup> L. Lueking,<sup>14</sup> A.L. Lyon,<sup>25</sup> A.K.A. Maciel,<sup>10</sup>  
 R.J. Madaras,<sup>23</sup> R. Madden,<sup>15</sup> L. Magaña-Mendoza,<sup>11</sup> V. Manankov,<sup>29</sup> S. Mani,<sup>7</sup>  
 H.S. Mao,<sup>14,†</sup> R. Markeloff,<sup>33</sup> T. Marshall,<sup>18</sup> M.I. Martin,<sup>14</sup> K.M. Mauritz,<sup>19</sup> B. May,<sup>34</sup>

A.A. Mayorov,<sup>38</sup> R. McCarthy,<sup>45</sup> J. McDonald,<sup>15</sup> T. McKibben,<sup>17</sup> J. McKinley,<sup>27</sup>  
 T. McMahon,<sup>36</sup> H.L. Melanson,<sup>14</sup> M. Merkin,<sup>29</sup> K.W. Merritt,<sup>14</sup> H. Miettinen,<sup>40</sup>  
 A. Mincer,<sup>31</sup> C.S. Mishra,<sup>14</sup> N. Mokhov,<sup>14</sup> N.K. Mondal,<sup>46</sup> H.E. Montgomery,<sup>14</sup>  
 P. Mooney,<sup>1</sup> H. da Motta,<sup>10</sup> C. Murphy,<sup>17</sup> F. Nang,<sup>2</sup> M. Narain,<sup>14</sup> V.S. Narasimham,<sup>46</sup>  
 A. Narayanan,<sup>2</sup> H.A. Neal,<sup>26</sup> J.P. Negret,<sup>1</sup> P. Nemethy,<sup>31</sup> D. Norman,<sup>48</sup> L. Oesch,<sup>26</sup>  
 V. Oguri,<sup>41</sup> E. Oliveira,<sup>10</sup> E. Oltman,<sup>23</sup> N. Oshima,<sup>14</sup> D. Owen,<sup>27</sup> P. Padley,<sup>40</sup> A. Para,<sup>14</sup>  
 Y.M. Park,<sup>21</sup> R. Partridge,<sup>5</sup> N. Parua,<sup>46</sup> M. Paterno,<sup>42</sup> B. Pawlik,<sup>22</sup> J. Perkins,<sup>47</sup>  
 M. Peters,<sup>16</sup> R. Piegaiia,<sup>6</sup> H. Piekarczyk,<sup>15</sup> Y. Pischalnikov,<sup>39</sup> B.G. Pope,<sup>27</sup> H.B. Prosper,<sup>15</sup>  
 S. Protopopescu,<sup>4</sup> J. Qian,<sup>26</sup> P.Z. Quintas,<sup>14</sup> R. Raja,<sup>14</sup> S. Rajagopalan,<sup>4</sup> O. Ramirez,<sup>17</sup>  
 L. Rasmussen,<sup>45</sup> S. Reucroft,<sup>32</sup> M. Rijssenbeek,<sup>45</sup> T. Rockwell,<sup>27</sup> M. Roco,<sup>14</sup> P. Rubinov,<sup>34</sup>  
 R. Ruchti,<sup>35</sup> J. Rutherford,<sup>2</sup> A. Sánchez-Hernández,<sup>11</sup> A. Santoro,<sup>10</sup> L. Sawyer,<sup>24</sup>  
 R.D. Schamberger,<sup>45</sup> H. Schellman,<sup>34</sup> J. Sculli,<sup>31</sup> E. Shabalina,<sup>29</sup> C. Shaffer,<sup>15</sup>  
 H.C. Shankar,<sup>46</sup> R.K. Shivpuri,<sup>13</sup> M. Shupe,<sup>2</sup> H. Singh,<sup>9</sup> J.B. Singh,<sup>37</sup> V. Sirotenko,<sup>33</sup>  
 W. Smart,<sup>14</sup> E. Smith,<sup>36</sup> R.P. Smith,<sup>14</sup> R. Snihur,<sup>34</sup> G.R. Snow,<sup>30</sup> J. Snow,<sup>36</sup> S. Snyder,<sup>4</sup>  
 J. Solomon,<sup>17</sup> M. Sosebee,<sup>47</sup> N. Sotnikova,<sup>29</sup> M. Souza,<sup>10</sup> A.L. Spadafora,<sup>23</sup> G. Steinbrück,<sup>36</sup>  
 R.W. Stephens,<sup>47</sup> M.L. Stevenson,<sup>23</sup> D. Stewart,<sup>26</sup> F. Stichelbaut,<sup>45</sup> D. Stoker,<sup>8</sup> V. Stolin,<sup>28</sup>  
 D.A. Stoyanova,<sup>38</sup> M. Strauss,<sup>36</sup> K. Streets,<sup>31</sup> M. Strovink,<sup>23</sup> A. Sznajder,<sup>10</sup>  
 P. Tamburello,<sup>25</sup> J. Tarazi,<sup>8</sup> M. Tartaglia,<sup>14</sup> T.L.T. Thomas,<sup>34</sup> J. Thompson,<sup>25</sup>  
 T.G. Trippe,<sup>23</sup> P.M. Tuts,<sup>12</sup> N. Varelas,<sup>17</sup> E.W. Varnes,<sup>23</sup> D. Vititoe,<sup>2</sup> A.A. Volkov,<sup>38</sup>  
 A.P. Vorobiev,<sup>38</sup> H.D. Wahl,<sup>15</sup> G. Wang,<sup>15</sup> J. Warchol,<sup>35</sup> G. Watts,<sup>5</sup> M. Wayne,<sup>35</sup>  
 H. Weerts,<sup>27</sup> A. White,<sup>47</sup> J.T. White,<sup>48</sup> J.A. Wightman,<sup>19</sup> S. Willis,<sup>33</sup> S.J. Wimpenny,<sup>9</sup>  
 J.V.D. Wirjawan,<sup>48</sup> J. Womersley,<sup>14</sup> E. Won,<sup>42</sup> D.R. Wood,<sup>32</sup> H. Xu,<sup>5</sup> R. Yamada,<sup>14</sup>  
 P. Yamin,<sup>4</sup> J. Yang,<sup>31</sup> T. Yasuda,<sup>32</sup> P. Yepes,<sup>40</sup> C. Yoshikawa,<sup>16</sup> S. Youssef,<sup>15</sup> J. Yu,<sup>14</sup>  
 Y. Yu,<sup>44</sup> Z. Zhou,<sup>19</sup> Z.H. Zhu,<sup>42</sup> D. Zieminska,<sup>18</sup> A. Zieminski,<sup>18</sup> E.G. Zverev,<sup>29</sup>  
 and A. Zylberstejn<sup>43</sup>

(DØ Collaboration)

<sup>1</sup> *Universidad de los Andes, Bogotá, Colombia*

<sup>2</sup> *University of Arizona, Tucson, Arizona 85721*

<sup>3</sup> *Boston University, Boston, Massachusetts 02215*

<sup>4</sup> *Brookhaven National Laboratory, Upton, New York 11973*

<sup>5</sup> *Brown University, Providence, Rhode Island 02912*

<sup>6</sup> *Universidad de Buenos Aires, Buenos Aires, Argentina*

<sup>7</sup> *University of California, Davis, California 95616*

<sup>8</sup> *University of California, Irvine, California 92697*

<sup>9</sup> *University of California, Riverside, California 92521*

<sup>10</sup> *LAFEX, Centro Brasileiro de Pesquisas Físicas, Rio de Janeiro, Brazil*

<sup>11</sup> *CINVESTAV, Mexico City, Mexico*

<sup>12</sup> *Columbia University, New York, New York 10027*

<sup>13</sup> *Delhi University, Delhi, India 110007*

<sup>14</sup> *Fermi National Accelerator Laboratory, Batavia, Illinois 60510*

<sup>15</sup> *Florida State University, Tallahassee, Florida 32306*

<sup>16</sup> *University of Hawaii, Honolulu, Hawaii 96822*

<sup>17</sup> *University of Illinois at Chicago, Chicago, Illinois 60607*

- <sup>18</sup> *Indiana University, Bloomington, Indiana 47405*
- <sup>19</sup> *Iowa State University, Ames, Iowa 50011*
- <sup>20</sup> *Korea University, Seoul, Korea*
- <sup>21</sup> *Kyungsoong University, Pusan, Korea*
- <sup>22</sup> *Institute of Nuclear Physics, Kraków, Poland*
- <sup>23</sup> *Lawrence Berkeley National Laboratory and University of California, Berkeley, California 94720*
- <sup>24</sup> *Louisiana Tech University, Ruston, Louisiana 71272*
- <sup>25</sup> *University of Maryland, College Park, Maryland 20742*
- <sup>26</sup> *University of Michigan, Ann Arbor, Michigan 48109*
- <sup>27</sup> *Michigan State University, East Lansing, Michigan 48824*
- <sup>28</sup> *Institute for Theoretical and Experimental Physics, Moscow, Russia*
- <sup>29</sup> *Moscow State University, Moscow, Russia*
- <sup>30</sup> *University of Nebraska, Lincoln, Nebraska 68588*
- <sup>31</sup> *New York University, New York, New York 10003*
- <sup>32</sup> *Northeastern University, Boston, Massachusetts 02115*
- <sup>33</sup> *Northern Illinois University, DeKalb, Illinois 60115*
- <sup>34</sup> *Northwestern University, Evanston, Illinois 60208*
- <sup>35</sup> *University of Notre Dame, Notre Dame, Indiana 46556*
- <sup>36</sup> *University of Oklahoma, Norman, Oklahoma 73019*
- <sup>37</sup> *University of Panjab, Chandigarh 16-00-14, India*
- <sup>38</sup> *Institute for High Energy Physics, Protvino 142284, Russia*
- <sup>39</sup> *Purdue University, West Lafayette, Indiana 47907*
- <sup>40</sup> *Rice University, Houston, Texas 77005*
- <sup>41</sup> *Universidade do Estado do Rio de Janeiro, Brazil*
- <sup>42</sup> *University of Rochester, Rochester, New York 14627*
- <sup>43</sup> *CEA, DAPNIA/Service de Physique des Particules, CE-SACLAY, Gif-sur-Yvette, France*
- <sup>44</sup> *Seoul National University, Seoul, Korea*
- <sup>45</sup> *State University of New York, Stony Brook, New York 11794*
- <sup>46</sup> *Tata Institute of Fundamental Research, Colaba, Mumbai 400005, India*
- <sup>47</sup> *University of Texas, Arlington, Texas 76019*
- <sup>48</sup> *Texas A&M University, College Station, Texas 77843*

(March 23, 1998)

## Abstract

We have searched for central production of a pair of photons with high transverse energies in  $p\bar{p}$  collisions at  $\sqrt{s} = 1.8$  TeV using  $70 \text{ pb}^{-1}$  of data collected with the DØ detector at the Fermilab Tevatron in 1994–1996. If they exist, virtual heavy pointlike Dirac monopoles could rescatter pairs of nearly real photons into this final state via a box diagram. We observe no excess of events above background, and set lower 95% C.L. limits of 610, 870, or 1580 GeV/c<sup>2</sup> on the mass of a spin 0, 1/2, or 1 Dirac monopole.

PACS numbers: 12.60.-i, 14.80.-j, 13.85.Rm

*Submitted to Phys. Rev. Lett.*

One of the open questions of particle physics is the existence of Dirac monopoles [1,2], hypothetical carriers of the magnetic charge proposed by P. Dirac to symmetrize the Maxwell equations and explain the quantization of electric charge. If such magnetic monopoles exist, then the elementary magnetic and electric charges ( $g$  and  $e$ ) must be quantized according to the following formula:

$$g = \frac{2\pi n}{e}, \quad n = \pm 1, \pm 2, \dots, \quad (1)$$

where  $n$  is an unknown integer. Here we assume that the elementary electric charge is that of an electron. If free quarks exist, Eq. (1) should be modified by replacing  $e$  with  $e/3$ , which effectively increases  $g$  by a factor of three.

Dirac monopoles are expected to couple to photons with a coupling constant  $\alpha_g = g^2/4\pi \approx 34 n^2$  which is at least three orders of magnitude larger than the corresponding photon coupling to the electric charge ( $\alpha_e = e^2/4\pi \approx 1/137$ ). Therefore such monopoles could give rise to photon-photon rescattering via the box diagram shown in Fig. 1 [3,4]. The contribution of this diagram for pointlike monopoles to diphoton production at hadron colliders was recently calculated [5] and shown to be significant even for monopole masses comparable to the collider beam energy.

Since the virtuality ( $Q^2$ ) of most incoming photons in the process of Fig. 1 is small [6], the interacting partons scatter at very small angles and therefore escape the detector through the beam pipe. Thus, a signature for monopoles at hadron colliders is the production of a pair of isolated photons with high transverse energies. This process gives a unique opportunity to find evidence for Dirac monopoles or to set limits on the monopole mass.




FIG. 1. Feynman diagram for  $\gamma\gamma$  production via a virtual monopole loop.

The only previous study of this nature was made by the L3 experiment at the LEP  $e^+e^-$  accelerator by searching for the  $Z \rightarrow \gamma\gamma\gamma$  decay via a similar monopole loop [7]. It resulted in the lower 95% confidence level (C.L.) mass limit of 510 GeV for pointlike spin 1/2 monopoles. Other accelerator experiments (see Ref. [8]) have looked for production

of monopoles by searching for the high ionization traces that would be produced by these particles, and would therefore be inherently restricted to monopole masses below the beam energy. A variety of experiments which look for monopoles in cosmic rays are sensitive to the relic monopole flux, rather than the monopole mass [8]. Indirect limits on the Dirac monopole mass can be derived from measurements of the top quark mass and the axial and vector couplings of the  $Z$  to charged leptons [4].

Despite numerous studies, QED with pointlike monopoles is still not a complete theory. For example, it is not clear whether such a theory can be constructed to be renormalizable to all orders [4]. Also, arguments exist (see, e.g. [9]) that Dirac monopoles must occupy a spatial volume of radius  $R \sim \mathcal{O}(g^2/M)$ , where  $M$  is the monopole mass, to accommodate the self energy implied by the large coupling. The non-observation of a new distance scale in QED or the SM for  $R < \mathcal{O}(1 \text{ TeV})$  requires the monopole mass to exceed  $\sim 100 \text{ TeV}$ . Further theoretical work on this subject therefore is required to define the regions of validity for a theory of pointlike monopoles. In such a theory, it is possible that hard interactions of a monopole with photons would be weakened substantially by the effects of a monopole form factor.

In this Letter we report on the results of a new search for Dirac monopoles with the  $D\phi$  detector (described in detail elsewhere [10]) operating at the Fermilab Tevatron proton-antiproton collider with beam energies of 900 GeV. The search is based on  $69.5 \pm 3.7 \text{ pb}^{-1}$  of data recorded in 1994–1996 using a trigger which required the presence of an electromagnetic (EM) object with transverse energy  $E_T$  above 40 GeV. This trigger did not require the presence of an inelastic collision, and therefore can be used to select low  $Q^2$  events typical of the process in Fig. 1.

The following offline selection criteria are: (i) at least two photons with  $E_T > 40 \text{ GeV}$  and pseudorapidity  $|\eta_\gamma| < 1.1$ ; (ii) missing transverse energy in the event  $\cancel{E}_T < 25 \text{ GeV}$ ; and (iii) no jets with  $E_T^j > 15 \text{ GeV}$  and  $|\eta^j| < 2.5$ . The jet veto requirement is used to select the low  $Q^2$  process in Fig. 1. The trigger is  $> 98\%$  efficient for this off-line selection.

In order to determine the hard scattering vertex, we calculate the most probable direction of each photon using the transverse and longitudinal segmentation of the EM calorimeter [11]. These directions determine the position of the interaction vertex along the beam axis. The resolution on the vertex position for this method is 14 cm, taken from  $Z \rightarrow ee$  decays where the vertex can also be determined with high precision using the tracking information. This EM-cluster-based vertex finding technique is preferred since for the event topology of Fig. 1 one does not expect charged particles, causing the tracking-based vertex finding to be biased significantly toward vertices from background interactions. We calculate kinematic parameters of the event based on the vertex obtained using the EM clusters.

Each photon is required to have: (i) energy isolation [11]  $I < 0.1$ ; (ii) more than 95% of the cluster energy deposited in the EM calorimeter; (iii) cluster shape consistent with that expected for a photon; and (iv) no tracks pointing toward the EM cluster from any of the event vertices.

The overall efficiency for photon identification is  $(73.0 \pm 1.2)\%$  per photon, as detailed in Table I. This includes the  $(92 \pm 1)\%$  probability of the photon not to convert in the material in front of the tracking chambers. The efficiency of criteria (i)–(iii) is determined using the  $Z \rightarrow ee$  events (with (ii) additionally checked using a GEANT [12] simulation of the  $D\phi$  detector for possible energy dependence); the efficiency of the no track requirement (iv)



TABLE I. Signal efficiency.

Cut	Efficiency (%)
per photon	
$I < 0.10$	$93.0 \pm 0.7$
EM fraction	$99.0 \pm 1.0$
Shape consistency	$94.7 \pm 0.8$
No tracks	$91.1 \pm 0.4$
No photon conversions	$92.0 \pm 1.0$
per event	
$\cancel{E}_T < 25$ GeV	$99.0 \pm 0.5$
Overall	$52.8 \pm 1.4$

was determined using simulated photons obtained by rotating the electromagnetic clusters from  $Z \rightarrow ee$  decays by  $\pi/2$  in azimuth [11]. The overall efficiency for the diphoton selection is  $(52.8 \pm 1.4)\%$ . This includes the efficiency of the  $\cancel{E}_T$  veto ( $99.0 \pm 0.5\%$ ) as well as the identification efficiency for a pair of photons.

The above selection criteria define our base sample which contains 90 candidate events.

The main backgrounds to photon scattering through a monopole loop are due to: (i) diagrams similar to Fig. 1 with other particles in the loop; (ii) QCD production of dijets ( $jj$ ) and direct photons ( $j\gamma$ ) (with jets misidentified as photons due to fragmentation into a leading  $\pi^0$  or  $\eta$  decaying into a pair of spatially close photons, reconstructed as one EM cluster), or direct diphotons ( $\gamma\gamma$ ); and (iii) Drell-Yan dielectron production with electrons misidentified as photons due to tracking inefficiency.

Background (i) is dominated by a virtual  $W$ -loop and has been shown to be negligible [13]. The other two background contributions are estimated from the data. The QCD background is determined using the  $j\gamma$  event sample collected with a single photon trigger, with the jet passing the same fiducial and kinematic cuts as the photon. We apply a jet-faking probability  $P(j \rightarrow \gamma)$  which we measure to be  $(10.5 \pm 1.5) \times 10^{-4}$  by counting the number of photons in multijet events, and find the QCD background to be  $25 \pm 8$  events. Direct photon and diphoton backgrounds are also included in this estimate. Their relative fractions are obtained from PYTHIA [14] Monte Carlo (MC). The 30% error assigned to the QCD background estimate reflects the uncertainty in the direct photon fractions and in the jet-faking-photon probability.

The Drell-Yan background is calculated from a sample of dielectron events passing the same fiducial and kinematic cuts as the signal sample. Multijet contamination of this sample is negligible since the probability for a jet to be misidentified as an electron is five times smaller than that for a photon. The probability for a dielectron pair to be misidentified as a diphoton pair is found to be  $(11 \pm 1)\%$  by comparing the number of events in the  $Z$  peak in the  $ee$  and  $\gamma\gamma$  samples passing loose kinematic cuts. The Drell-Yan background in the base sample is  $63 \pm 7$  events. The overall background in the base sample is  $88 \pm 11$  (syst) events, in good agreement with the 90 observed candidates.

To optimize the sensitivity of this search to the monopole loop contribution we apply a cut on the scalar sum of the transverse energies of all the photons in the event:  $S_T \equiv \sum_i E_T^{\gamma_i}$  [15].




FIG. 2. Data and expected background as a function of  $S_T^{\min}$  cut. Points are data, the upper hatched region corresponds to the QCD background, and the lower shaded region shows the Drell-Yan background. The  $\approx 15\%$  systematic error on the background is not shown.

We vary the  $S_T$  cut threshold ( $S_T^{\min}$ ) in 10 GeV steps to achieve an expected background of 0.4 events [16]. Such an optimization is based on the fact that for this expected background one has a 67% probability of observing no candidate events in the data in the absence of a signal. In such a case [8], the limits on the signal do not depend on the exact background value or its uncertainties. The agreement between the observed number of events and the predicted background as a function of  $S_T^{\min}$  is illustrated in Fig. 2. Note that since the plot shows the data and the backgrounds for  $S_T > S_T^{\min}$ , the points are highly correlated. The  $S_T^{\min} = 250$  GeV cut corresponds to a background of  $0.41 \pm 0.11$  events. The event in the base sample with highest  $S_T$  has  $S_T = 203$  GeV, well below this cut. Taking into account the selection efficiency we set an upper limit for the production cross section of two or more photons with  $\sum E_T^\gamma > 250$  GeV and  $|\eta^\gamma| < 1.1$ :

$$\sigma(p\bar{p} \rightarrow \geq 2\gamma)|_{S_T > 250 \text{ GeV}, |\eta^\gamma| < 1.1} < 83 \text{ fb} \quad (2)$$

at the 95% C.L. This limit is obtained using a Bayesian approach with a flat prior and with the uncertainties in the efficiency and the integrated luminosity properly taken into account.

Since the data are consistent with the background hypothesis, we can set limits on the production of pointlike Dirac monopoles. We calculate the acceptance for the monopole signal using a fast MC program that generates diphoton events from a monopole loop according to the calculated differential cross section  $d^3\sigma/dE_T^\gamma d\eta^{\gamma_1} d\eta^{\gamma_2}$  [5] with a subsequent parametric simulation of the DØ detector. The MC model takes into account the interaction vertex distribution; parton density distributions in the colliding protons and antiprotons, as described by the GRV [17] parton distribution functions (p.d.f.); smearing of photon momenta; and detector acceptance. Figures 3(a) and (b) show the expected signal  $S_T$  distribution and the

correlation between the photon pseudorapidities, respectively. The cuts used in this analysis are indicated in the figures. The overall acceptance for the monopole signal is found to be  $(51 \pm 1)\%$ , where the error reflects variations due to different p.d.f. (estimated by taking the acceptance difference using GRV and CTEQ4L [18]), and uncertainty in the detector response parametrization. The acceptance does not depend on the monopole mass for masses above the typical photon energy ( $\sim 300$  GeV) [6].

The total cross section of the heavy monopole production at the Tevatron is given by [5]:

$$\sigma(p\bar{p} \rightarrow \gamma\gamma + X) = 57 P \left( \frac{n}{M [\text{TeV}]} \right)^8 \text{ fb}, \quad (3)$$

where  $P$  is a spin dependent factor [19,13]:  $P = 0.085, 1.39,$  and  $159$  for monopole spin of  $0, 1/2,$  and  $1,$  respectively. The estimated error on this cross section due to choice of p.d.f. and to higher order QED effects is  $30\%$  [6]. Additional uncertainties are associated with the  $\gamma\gamma \rightarrow \gamma\gamma$  subprocess in Fig. 1 and with unitarity considerations. The coupling constant  $\alpha_g$  is replaced with an effective coupling [5] obtained by multiplying  $\alpha_g$  by a factor  $(E^\gamma/M)^2$ , where  $E^\gamma$  is the photon energy, typically  $300$  GeV at the Tevatron. Both unitarity and perturbation theory assumptions are satisfied when this factor is  $\ll 1$  [3,5].

Comparing the lower bound of the theoretical cross section corrected for acceptance with the cross section limit (2) set by this measurement, we obtain the following lower limits on the pointlike Dirac monopole mass (see Fig. 4):

$$M/n > \begin{cases} 610 \text{ GeV} & \text{for } S = 0 \\ 870 \text{ GeV} & \text{for } S = 1/2 \\ 1580 \text{ GeV} & \text{for } S = 1 \end{cases} .$$

These are currently the most stringent limits on heavy pointlike monopole mass. (We do assume, if more than one type of Dirac monopole exists, that there is no cancellation among the loop diagrams involving each monopole type.)

We note that the effective coupling exceeds 1 and unitarity is violated close to the experimental bound. For values  $E^\gamma/M > 1$ , the cross section will grow more slowly, approaching the usual  $1/M^2$  behavior of a QED process [6] which satisfies unitarity. Also, for lower monopole masses the effective parameter of the perturbation theory used in the calculations [5] becomes too large, and therefore one would expect a non-negligible contribution of the higher order diagrams with four, six, etc. photons in the final state. The latter effect is, however, largely compensated by our analysis cut on the sum of the photon transverse energies; if part of the signal cross section is due to the higher order diagrams, the above limits are unaffected.

When more complete theoretical calculations are available, limits on the monopole mass could be updated by comparing the modified cross section expression with the experimental limit (2). With current theory [5] the above limits are strictly valid only for monopole masses above several hundred GeV.

As a cross-check of the results of this search we have selected elastic or nearly elastic collisions by requiring no hits in the forward scintillating hodoscopes used for luminosity monitoring and triggering on the inelastic collisions [10]. This requirement drastically reduces the backgrounds. The remaining background in the base sample for elastic events




FIG. 3. a) Normalized  $S_T$  spectrum and b) photon pseudorapidities for the diphoton production via a heavy monopole loop. The arrow in a) and square in b) show the chosen cuts in the corresponding parameters.

is  $1.8 \pm 0.4$  events, dominated by diffractive Drell-Yan events and residual inelastic background due to inefficiency of the forward hodoscopes. We observe one candidate event in the base sample, consistent with this expected background rate. For  $S_T^{\min} \approx 100$  GeV the background is 0.4 events, and no candidates are observed. We use this sample only as a cross check because the efficiency of these selection requirements is significantly lower than that of the main analysis method, primarily because of multiple interactions.

In conclusion, we have performed a search for heavy pointlike Dirac monopoles by searching for pairs of photons with high transverse energies. Our data agree with the expected background from QCD and Drell-Yan production. No candidates pass the final cuts. Using theoretical calculations [5] we set 95% C.L. lower limits on the Dirac monopole mass for minimum magnetic charge ( $n = 1$ ) in the range 610 to 1580 GeV, depending on the monopole spin. These are the most stringent mass limits on heavy pointlike Dirac monopoles to date. Our cross section limit (2) is 83 fb, and may be applicable to the other production processes, such as that of dyons [4] or other exotic objects strongly coupled to photons.

We are grateful to I. Ginzburg and A. Schiller for many discussions and detailed cross section information and to U. Baur, B. Dobrescu, and A.S. Goldhaber for helpful discussions. We thank the staffs at Fermilab and collaborating institutions for their contributions to this work, and acknowledge support from the Department of Energy and National Science Foundation (U.S.A.), Commissariat à L’Energie Atomique (France), State Committee for Science and Technology and Ministry for Atomic Energy (Russia), CAPES and CNPq (Brazil), Departments of Atomic Energy and Science and Education (India), Colciencias (Colombia), CONACyT (Mexico), Ministry of Education and KOSEF (Korea), and CONICET and UBACyT (Argentina).




FIG. 4. The curved bands show the low and upper bounds on theoretical cross sections [5] for monopole spin,  $S = 0, 1/2$ , and  $1$ . The horizontal line shows the 95% CL experimental upper limit (2) on the cross section. The arrows indicate the lower 95% CL limits on the monopole mass at each spin value.

## REFERENCES

\* Visitor from Universidad San Francisco de Quito, Quito, Ecuador.

† Visitor from IHEP, Beijing, China.

- [1] P.A.M. Dirac, Proc. R. Soc. London, Ser. A **133**, 60 (1931).
- [2] J. Schwinger, Phys. Rev. **151**, 1055 (1966).
- [3] I.F. Ginzburg, S.L. Panfil, Sov. J. Nucl. Phys. **36**, 850 (1982).

- [4] A. De Rújula, Nucl. Phys. **B435** 257 (1995).
- [5] I.F. Ginzburg and A.Schiller, hep-ph/9802310, submitted to Phys. Rev. D.
- [6] I.F. Ginzburg, private communication.
- [7] L3 Collaboration, M. Acciarri *et al.*, Phys. Lett. **B345**, 609 (1995).
- [8] PDG Review of Particle Physics, Phys. Rev. D **54**, 166, 685-687 (1996).
- [9] A.S. Goldhaber, in Proceedings of the CRM-FIELDS-CAP Workshop “Solitons” at Queen’s University, Kingston, Ontario, July 1997 (Springer, New York 1998).
- [10] DØ Collaboration, S. Abachi *et al.*, Nucl. Instrum. Methods **A338**, 185 (1994).
- [11] DØ Collaboration, S. Abachi *et al.*, Phys. Rev. Lett. **78**, 3640 (1997); *ibid.* Phys. Rev. D **56** 6742 (1997).
- [12] R. Brun and F. Carminati, CERN Program Library Writeup W5013, 1993 (unpublished). We used GEANT version 3.15.
- [13] G. Jikia and A. Tkabaladze, Phys. Lett. **B323**, 453 (1994).
- [14] T. Sjöstrand, Comp. Phys. Comm. **82**, 74 (1994). We used PYTHIA version 5.7.
- [15] We use the  $S_T$  variable instead of the individual photon energies, since it is less sensitive to the next-to-leading order QED corrections to the diagram in Fig. 1. See discussion later in the main text.
- [16] DØ Collaboration, S. Abachi *et al.*, Phys. Rev. Lett. **79**, 4321 (1997); B. Abbott *et al.*, Phys. Rev. Lett. **80**, 2051 (1998).
- [17] M. Glück, E. Reya and A. Vogt, Z. Phys. **C67**, 433 (1995).
- [18] CTEQ Collaboration, H.L. Lai *et al.*, Phys. Rev. D **55**, 1280 (1997).
- [19] W. Heisenberg and H. Euler, Z. Phys. **38**, 714 (1936); V. Constantini, B. De Tollis, and G. Pistoni, Nuovo Cim. **2A**, 733 (1971); M. Baillagreon, G. Belanger, and F. Boudjema, Phys. Rev. D **51**, 4712 (1995); M. Baillagreon, F. Boudjema, E. Chopin, and V. Lafage, Z. Phys. **C67**, 431 (1996).