

Roles of reactive oxygen species in interactions between plants and pathogens

Nandini P. Shetty · Hans J. Lyngs Jørgensen ·
Jens Due Jensen · David B. Collinge ·
H. Shekar Shetty

Received: 23 May 2007 / Accepted: 3 March 2008
© KNPV 2008

Abstract The production of reactive oxygen species (ROS) by the consumption of molecular oxygen during host–pathogen interactions is termed the oxidative burst. The most important ROS are singlet oxygen ($^1\text{O}_2$), the hydroxyl radical ($\text{HO}_2\cdot$), the superoxide anion (O_2^-), hydrogen peroxide (H_2O_2), the hydroxyl radical ($\text{OH}\cdot$) and the closely related reactive nitrogen species, nitric oxide (NO). These ROS are highly reactive, and therefore toxic, and participate in several important processes related to defence and infection. Furthermore, ROS also play important roles in plant biology both as toxic by-products of aerobic metabolism and as key regulators of growth, development and defence pathways. In this review, we will assess the different roles of ROS in host–pathogen interactions with special emphasis on fungal and Oomycete pathogens.

Keywords Antimicrobial · Cell wall cross-linking · Hypersensitive response · Signal transduction · Gene expression · Successful pathogenesis · Hydrogen peroxide

Abbreviations

| | |
|------|----------------------------------|
| ROS | reactive oxygen species |
| SA | salicylic acid |
| ET | ethylene |
| MAPK | mitogen-activated protein kinase |
| SOD | superoxide dismutase |
| CWA | cell wall appositions |
| NO | nitric oxide |
| JA | jasmonic acid |
| HR | hypersensitive response |
| PCD | programmed cell death |

Introduction

Generation of ROS, especially hydrogen peroxide (H_2O_2), has been recorded in interactions with a variety of pathogens (Mellersh et al. 2002; Shetty et al. 2003; Thordal-Christensen et al. 1997; Unger et al. 2005). Avirulent pathogens often induce a biphasic ROS accumulation with a small, transient first phase, followed by a continuous phase of much higher intensity that correlates with disease resistance (Lamb and Dixon 1997; Torres et al. 2006). However, three phases of ROS accumulation have been observed in

N. P. Shetty (✉) · H. J. L. Jørgensen · J. D. Jensen ·
D. B. Collinge
Department of Plant Biology, Faculty of Life Sciences,
University of Copenhagen,
Thorvaldsensvej 40,
DK-1871 Frederiksberg, Denmark
e-mail: nps@life.ku.dk

H. S. Shetty
Department of Studies in Applied Botany,
Seed Pathology and Biotechnology, University of Mysore,
Manasgangotri,
Mysore 570 006, India

some cases, e.g., for *Blumeria graminis* f. sp. *hordei* infecting barley (Hückelhoven and Kogel 2003) and *Septoria tritici* infecting wheat (Shetty et al. 2003). These differences can be attributed to the more complicated development of these fungal pathogens and the influence of the host genotype, which presumably determine whether two or three phases of ROS accumulation occur. Virulent pathogens that avoid or suppress host recognition induce only the transient, first phase of this response (Bolwell et al. 2002). Elicitors of defence responses, often now referred to as microbe or pathogen-associated molecular patterns (PAMPs), also trigger an oxidative burst (Chisholm et al. 2006). There are several potential sources of ROS in plants and different sources of ROS may be activated within a species in different situations depending on the type of stress (Bolwell et al. 2002; Lamb and Dixon 1997). A variety of enzyme systems have been implicated in ROS generation following pathogen recognition, i.e., reduced form of nicotinamide adenine dinucleotide phosphate (NADPH) oxidase (Bedard et al. 2007; Carter et al. 2007; Grant et al. 2000), SOD (Auh and Murphy 1995; Deepak et al. 2006), oxalate oxidases (Hu et al. 2003; Zimmermann et al. 2006), peroxidases (Bindschedler et al. 2006; Bolwell et al. 2002), lipoxygenases (Babitha et al. 2004) and amine oxidases (Allan and Fluhr 1997; Cona et al. 2006; Walters 2003). Stress on of ROS-producing organelles during pathogenesis may also contribute to ROS production during host–pathogen interactions. Mitochondria are normally considered relatively unimportant ROS generators in photosynthesising tissue (Apel and Hirt 2004; Kuźniak and Skłodowska 2005). However, a recent review by Amirsadeghi et al. (2007) discusses evidence that mitochondria are a potential source of ROS in response to biotic stress. Chloroplasts (Kariola et al. 2005) and peroxisomes (Kuźniak and Skłodowska 2005) have also been shown to be important. In this review, we present recent knowledge on the roles of ROS during host–pathogen interactions with special emphasis on fungal and Oomycete pathogens.

Roles of ROS in host–pathogen interactions

ROS have been implicated in many different processes related to pathogen interactions with their hosts. In the initial phases of the interactions, this essentially

means involvement in defence processes, whereas at the later stages, during pathogen colonisation, the role of ROS may be more ambiguous.

ROS as antimicrobial agents

ROS, especially H_2O_2 , was suggested as an antimicrobial agent during the plant defence response (Apostol et al. 1989; Custers et al. 2004; Legendre et al. 1993; Walters 2003). However, the actual toxicity of ROS in a given plant–pathogen interaction will depend on the sensitivity of the pathogen to the concentration of ROS present (Levine et al. 1994). The amount of extracellular H_2O_2 produced depends on several factors including the nature of the elicitor, the plant species, and age or developmental stages of the plant cells (Legendre et al. 1993; Małolepsza 2005; Nurnberger et al. 1994). Micromolar concentrations of H_2O_2 inhibited spore germination of a number of fungal pathogens *in vitro* (Peng and Kuc 1992). Thus, a concentration of 0.1 mM H_2O_2 completely inhibited the growth of cultured bacteria *Pectobacterium carotovorum* subsp. *carotovorum* (formerly *Erwinia carotovora* pv. *carotovora*), and resulted in >95% inhibition of *Phytophthora infestans* growth (Wu et al. 1995). Shetty et al. (2007) demonstrated by *in vitro* experiments that 5 mM H_2O_2 inhibited the development of inoculum from 4 day-old *S. tritici* cultures whereas a concentration of about 50 mM was required to inhibit inoculum from 16 day-old cultures. This reflects the ability of the pathogen to tolerate H_2O_2 during the different stages of its life-cycle. Shetty et al. (2007) also demonstrated that in the wheat–*S. tritici* interaction, infiltration of 4 mM H_2O_2 into a susceptible cultivar made it more resistant, symptoms appearing 6 days later than in control plants, whereas infiltration of catalase resulted in symptoms appearing 4 days earlier. It is currently not known whether the effect of H_2O_2 was direct, i.e., by toxicity of ROS, or indirect by affecting signal transduction or defence gene expression.

It is difficult to determine which H_2O_2 concentrations actually inhibit pathogens in planta since the necessary manipulation of the host tissue may itself trigger the production of ROS and/or antioxidants. However, ROS are also toxic to the plant. Thus, soybean suspension-cultured cells remained viable with up to 4 mM H_2O_2 , whereas slightly higher levels (6–8 mM) resulted in extensive cell death (Levine et

al. 1994). Therefore, ROS accumulation is tightly regulated by the plant to avoid high concentrations, which could damage the plant tissue (Torres et al. 2006).

Involvement of ROS in signal transduction and gene expression


ROS are involved in different signalling pathways for defence mechanisms, such as triggering of the HR, accumulation of phytoalexins and a number of other defence-response genes (cf. Fig. 1). It has been suggested that ROS are sensed by plants via three mechanisms (Mittler et al. 2004): unidentified receptor proteins; redox-sensitive transcription factors such as natriuretic peptide receptor 1 (NPR1) or heat-shock transcription factors (HSFs); and direct inhibition of phosphatase (Apel and Hirt 2004; Mittler et al. 2004; Neill et al. 2002). ROS signalling is the subject of intense studies, but the role of ROS in signalling is poorly understood. Here, we will give an overview of the initial events involved in signalling in plant–pathogen interactions.

Protein phosphorylation, changes in ion fluxes and the oxidative burst, leading to either HR or defence gene expression, or both, are important events taking place after pathogen infection (Chandra et al. 1996; Jabs et al. 1997; Lamb and Dixon 1997; Sasabe et al. 2000). The earliest reactions of plant cells include changes in plasma membrane permeability, which leads to Ca^{2+} and proton influx and K^+ and Cl^- efflux (McDowell and Dangl 2000). Ion fluxes subsequently induce extracellular production of ROS catalysed by

enzymes that act as secondary messengers for the HR and defence gene expression (Lamb and Dixon 1997). Calcium has been shown to be important in signalling. Heteromeric guanosine triphosphate (GTP)-binding proteins and protein phosphorylation/dephosphorylation events are probably involved in transferring the signals from the receptor to calcium channels that activate downstream processes (Legendre et al. 1992). Furthermore, elevation of the cytosolic calcium concentration has been shown to occur during most biotic and abiotic stresses (Price et al. 1996). For example, oxidative stress increased the cytosolic calcium concentration in tobacco (Price et al. 1994) and H_2O_2 induced calcium influx-mediated stomatal closure in *Commelina communis* and *Arabidopsis thaliana* (McAinsh et al. 1996; Pei et al. 2000). In further support of the involvement of calcium in signalling, lanthanide ions (calcium channel blocker) inhibited bacterial elicitor-induced ROS production in tobacco (Baker et al. 1993). Moreover, Urquhart et al. (2007) showed that transient expression of the chimeric cyclic nucleotide-gated ion channel gene ATCNGC11/12 in *Nicotiana benthamiana* gave rise to cell death with characteristics of the HR. Furthermore, it was shown that this gene could function as a Ca^{2+} -conducting channel and that calcium ions were important for the observed cell death. Recently, Ashtamker et al. (2007) showed that nuclei isolated from tobacco were capable of producing H_2O_2 . This was dependent on calcium, suggesting that nuclei can be a source of ROS production.

Different models for the action of calcium in the regulation of ROS have been proposed. One model

Fig. 1 Putative sources and functions of reactive oxygen species (ROS) in host–pathogen interactions of biotrophic and necrotrophic organisms



suggests that an elicitor interacts with a receptor coupled with a G-protein, which leads to Ca^{2+} influx that activates a Ca^{2+} -dependent protein kinase and ultimately NADPH oxidase (Blumwald et al. 1998). Another model, based on studies of innate immunity in *Arabidopsis*, suggests that pathogens or PAMPs are recognised by (unknown) receptors which trigger an ion (calcium) channel, leading to increases in cytosolic Ca^{2+} and subsequent NO generation (Ali et al. 2007). NO generation, together with other required factors such as an avirulent pathogen and an oxidative burst, could lead to the HR and potentially, diffusion of NO to neighbouring cells could act as a signal that thereby activates further calcium channels.

Activation of the oxidative burst is governed by phosphorylation/dephosphorylation (Lamb and Dixon 1997). Thus, a protein phosphorylation cascade that has been shown to be activated by H_2O_2 is a MAPK cascade, which has an important role in signal transduction (Zwerger and Hirt 2001). H_2O_2 has been shown to activate MAPK in *Arabidopsis* suspension cultures (Desikan et al. 1999). Furthermore, Petersen et al. (2000) showed that mutation of the MAPK gene *MPK4* in *Arabidopsis* altered plant defence activation.

H_2O_2 mediates the transcription of specific genes, though the exact mechanism is as yet unknown. Neill et al. (2002) suggested that it could be due to oxidation of cysteine residues of transcription factors. Activation of MAPKs is a common reaction of plant cells in defence-related signal transduction pathways (Neill et al. 2002). Perception of an extracellular signal activates a MAPK, which in turn can facilitate translocation of the signal to the nucleus where it can phosphorylate and activate transcription factors, thereby modulating gene expression (Apel and Hirt 2004; Hirt 1997; Zhou et al. 2004). For example, it has been reported that two tobacco MAPKs, namely salicylic acid-induced protein kinase (SIPK) and wound-induced protein kinase (WIPK), are regulated by a common upstream MAPK, which is involved in signalling for PCD (Zhang and Klessig 1998; Zhang et al. 2000). Ren et al. (2006) showed that there was another MAPK, Ntf4, with a similar function to SIPK and WIPK, which, when expressed in transgenic tobacco plants, accelerated the PCD when treated with the elicitor cryptogen from *Phytophthora cryptogea*. This indicates a role in signalling for PCD. Recently, Liu et al. (2007) showed that the combined activation of SIPK, Ntf4 and WIPK induced an HR-like PCD.

Activation of signal transduction could lead to increased ROS accumulation and activation of defence genes coding for PR-proteins, enzymes involved in the generation of phytoalexins, enzymes involved in oxidative stress protection, lignification and other defence responses (Alvarez et al. 1998; Apel and Hirt 2004; Lamb and Dixon 1997). Further evidence for a role of ROS in signalling has come from the fact that addition of low doses of ROS inducers stimulates the induction of detoxification mechanisms, such as SOD and glutathione-S-transferase, and activation of other defence mechanisms in neighbouring cells (Levine et al. 1994). Mittler et al. (2004) suggested that NADPH oxidase could be involved in ROS signalling by creating a loop where a small enhancement of ROS production and amplification of the ROS signals occurs in specific cellular locations. Pharmacological and genetic studies (Dat et al. 2003) support the existence of positive amplification loops involving NADPH oxidases in ROS signalling. These loops might be activated by low levels of ROS and result in enhanced production and amplification of the ROS signals. It has been reported that a small GTP-binding protein, Rac, regulates ROS production in rice, most likely through an NADPH oxidase, and induces cell death in rice cells with biochemical and morphological features similar to apoptosis in mammalian cells (Kawasaki et al. 1999). Together, MAPK and calcium-dependent protein kinases seem to play central roles in the regulation of pathogen-responsive NADPH oxidases at the transcriptional and post-transcriptional levels, respectively (Kobayashi et al. 2007).

It has been suggested that the HR is triggered only by balanced production of NO and ROS (Delledonne et al. 1998, 2002)—see also below. More specifically, dismutation of O_2^- to H_2O_2 is required to activate cell death, which depends on synergistic interactions between NO, H_2O_2 and SA (Delledonne et al. 1998; Mur et al. 2006). Scavenging of O_2^- by surplus NO (or vice versa) disturbs the NO/ H_2O_2 ratio, resulting in reduced cell death (Mur et al. 2006). Little is known about signalling pathways downstream of NO/ H_2O_2 . Nevertheless, it has been shown that NO signalling during both PCD and defence responses requires cyclic GMP and cyclic ADP ribose, two molecules that can serve as secondary messengers for NO signalling in mammals (Van Breusegem and Dat 2006).

SA has been shown to be an important signalling molecule involved in defence responses to pathogen attack in many plant–pathogen interactions. Thus, Enyedi et al. (1992) showed that SA levels increased dramatically in tobacco cells surrounding infection sites when infected with *Tobacco mosaic virus*. Torres et al. (2006) suggested that ROS acted synergistically in a signal amplification loop with SA to drive the HR and the establishment of systemic defences. SA accumulation can also down-regulate those ROS-scavenging systems that, in turn, can contribute to increased overall ROS levels following pathogen recognition (Klessig et al. 2000; Shah 2003). In addition to SA, ET and JA are also involved in signalling (Thatcher et al. 2005). The activation of a redox-signalling pathway possessing a MAPK module has also been reported in response to infection by avirulent pathogens in *Arabidopsis* (Suzuki 2002). This signalling network functions independently of the plant hormones ET, SA and JA (Thatcher et al. 2005). Additionally, when some mutants, which develop spontaneous lesions mimicking HR cell death, are placed in a *NahG* background to degrade SA (Shah 2003), lesion formation is suppressed but can be restored by SA treatment (Lorrain et al. 2003). However, this is not the case for all lesion-mimic mutants; some show intensified lesions in plants defective in JA signalling, while others have delayed lesion formation in plants defective in ET signalling (Lorrain et al. 2003). These differences in lesion formation can be due to synergistic or antagonistic effects between SA, JA and ET signalling pathways (Lorrain et al. 2003), but also indicates that mutants displaying the same phenotype could be mutated in widely different genes.

Involvement of ROS in oxidative cross-linking of cell walls

Barriers operating at the cell periphery to prevent invasion represent the first line of defence against pathogens that penetrate plant cells directly (Schulze-Lefert 2004). These barriers can, for example, depend on the nature and thickness of the epicuticular wax layer and cuticle or the composition and physical properties of the cell wall. Alternatively, they may occur by reinforcement of the cell wall, e.g., by deposition of callose-rich papillae and lignin at attempted penetration sites (Heitefuss 1997). ROS

production has been associated with the formation of physical defensive barriers (Hückelhoven and Kogel 2003; Lamb and Dixon 1997)—see also Fig. 2b. Association of H₂O₂ with lignification during plant development has been shown in several systems (Olson and Varner 1993; Repka 2002). Thus, H₂O₂ accumulation resulted in lignification in wounded *Zinnia* stem sections (Olson and Varner 1993). Furthermore, Thordal-Christensen et al. (1997) showed that the H₂O₂ production in barley infected with *B. graminis* f.sp. *hordei* led to cell wall cross-linking. Collins et al. (2003) showed that H₂O₂ was associated with vesicles containing cell wall components which were in transit to CWA, suggesting that H₂O₂ may play a role upstream of CWA or that compounds of CWA are oxidatively cross-linked on the way to the site of deposition. These findings have been confirmed by An et al. (2006), who showed that multivesicular bodies, intravacuolar vesicle aggregates and paramural bodies, which might participate in the secretion of building blocks for CWA, are associated with H₂O₂ accumulation. Another conspicuous role of the CWA, besides arresting fungal penetration, is blockage of all plasmodesmata between intact cells and those undergoing the HR, thereby containing the hypersensitive cell death. Also, Iwano et al. (2002) showed that in suspension-cultured rice cells infected with *Acidovorax avenae* (formerly *Pseudomonas avenae*), callose synthesis occurred at the H₂O₂ generation site.

Studies of pearl millet infected with *Sclerospora graminicola* have indicated that cell wall protein cross-linking is induced by enhanced H₂O₂ production at the time of pathogen attack (Kumudini and Shetty 2002). Possibly, hydroxyproline-rich glycoproteins (HRGPs) accumulate and contribute to disease resistance involving cross-linking between HRGP monomers to form a network which provide anchorage for lignification. This might also lead to obstruction of haustorial formation and nutrient shortage, which may be particularly unfavourable for biotrophic pathogens that use specific organs, e.g., haustoria for feeding (Bradley et al. 1992; Shailashree et al. 2004). Likewise, studies on the interaction between wheat and *B. graminis* f.sp. *tritici* showed that H₂O₂ plays important roles in defence, by driving among others the cross-linking to strengthen the cell wall (in effective papillae), and in association with HR (Li et al. 2005).




Fig. 2 Accumulation of H_2O_2 as seen by DAB-staining (Thordal-Christensen et al. 1997) in the barley–*B. graminis* f. sp. *hordei* interaction (**a**, **b**) and the wheat–*S. tritici* interaction (**c**, **d**). **a** shows barley isolate P-01 inoculated with isolate c15. A cell is undergoing HR as a response to penetration and is completely stained with DAB. **b** shows barley isolate P-02 inoculated with isolate c15, 2 days after inoculation. Note red–brown staining in the papillae and that the papilla in the cell containing an haustorium is not stained with DAB whereas the

other papillae are stained. **c** shows wheat cv. Stakado inoculated with isolate IPO323 of *S. tritici* (incompatible interaction) at 5 days after inoculation. H_2O_2 is accumulating in the apoplast of the substomatal cavity of a stoma penetrated by the pathogen. **d** shows wheat cv. Sevin inoculated with isolate IPO323 of *S. tritici* (compatible interaction) at 15 days after inoculation. H_2O_2 is accumulating throughout the tissue in which fungal sporulation occurs

Involvement of ROS in the hypersensitive response (HR)

The HR is a rapid host response occurring in a host cell, which is infected by a pathogen (Lam et al. 2001; Lam 2004). The cells die shortly after penetration Fig. 2a, often together with some of the surrounding cells (Greenberg 1997; Van Breusegem and Dat 2006). The HR occurs in order to restrict pathogen growth and is highly effective against biotrophic pathogens, since, with the death of host cells, the nutrient supply is removed (Greenberg and Yao 2004; Mellersh et al. 2002; Thordal-Christensen et al. 1997). In addition, toxic substances like ROS and phytoalexins produced in these cells apparently help to kill the pathogen (Lamb and Dixon 1997; Li et al. 2006; Moerschbacher and Reiserer 1997). ROS may originate primarily from chloroplasts, mitochondria and peroxisomes (Amirsadeghi et al. 2007; Lam 2004; Op den Camp et al. 2003; Van Breusegem and Dat 2006). The HR is often not effective against necrotrophic pathogens because these usually kill host cells to feed on them (Govrin and Levine 2000; Mayer et al.

2001). Thus, for true necrotrophic pathogens, such as *Botryotinia fuckeliana* (formerly *Botrytis cinerea*), it has been suggested that plant cell death is beneficial for infection, leading to enhanced colonisation (Govrin and Levine 2000; Greenberg and Yao 2004). However, the general nature of this conclusion has been questioned even for *B. fuckeliana* (Unger et al. 2005). In addition, there is a group of pathogens, often considered to be necrotrophic, which are in fact inhibited to some extent by HR, e.g., *Pyrenophora teres* (anamorph *Drechslera teres*; Jørgensen et al. 1998) and *Magnaporthe grisea* (Iwai et al. 2007). Collectively, these findings raise the question whether these pathogens are, in fact, necrotrophic, or should be considered as hemibiotrophic or whether some necrotrophic pathogens may also be inhibited by HR under some circumstances. In our view, the term ‘necrotrophic’ includes a diverse group of pathogens with quite different modes of pathogenicity.

The HR is a type of active PCD (Greenberg and Yao 2004; Lam 2004; Li et al. 2006; Sasabe et al. 2000; Van Breusegem and Dat 2006), which is often characterised by discrete cellular lesions and preceded

by an oxidative burst (Baker and Orlandi 1995; Dat et al. 2003; Levine et al. 1994; Sasabe et al. 2000). The process of HR may involve several steps including chromatin condensation, DNA cleavage and membrane blebbing, eventually leading to membrane disruption and release of cell contents (Dat et al. 2003; Hoeberichts and Woltering 2003; Lam 2004; Li et al. 2006; Sasabe et al. 2000). The cell death process thus shares some features with mammalian apoptosis (Greenberg and Yao 2004; Hoeberichts et al. 2003; Hoeberichts and Woltering 2003; Lam 2004). Further similarities include a group of proteins, termed metacaspases, in the genome of *Arabidopsis* with homology to the specific cysteine proteases, termed caspases, which play a key role in execution of mammalian apoptosis (Hoeberichts et al. 2003).

Involvement of ROS in HR has been studied by several different tools, including infiltration of anti-oxidants (Li et al. 2006) and ROS inhibitors or scavengers (Levine et al. 1994; Li et al. 2006; Sasabe et al. 2000). Furthermore, specific lines of *Arabidopsis* and other plants mutated in their ability to accumulate ROS or express antioxidants (such as SOD, catalase and ascorbate peroxidase) and subsequently activate HR have been studied (Dat et al. 2003; Hoeberichts and Woltering 2003; Jabs et al. 1996; Lorrain et al. 2003; Mateo et al. 2004; Mittler et al. 1999; Montillet et al. 2005; Op den Camp et al. 2003; Torres et al. 2005; Van Breusegem and Dat 2006). Also ROS accumulation has been studied following treatment with pathogen elicitors of HR (Greenberg and Yao 2004; Levine et al. 1994; Montillet et al. 2005; Sasabe et al. 2000). The use of such a diversity of approaches and systems strengthens and substantiates our knowledge on the role of ROS in HR, since broad background information is obtained. However, it also reveals that this process is highly complex, and not yet understood in detail (Van Breusegem and Dat 2006). For example, several studies have shown a correlation between accumulation of ROS (H_2O_2 , $^1\text{O}_2$, O_2^-), NO and HR (Dat et al. 2003; Floryszak-Wieczorek et al. 2007; Jabs et al. 1996; Levine et al. 1994; Mittler et al. 1999; Montillet et al. 2005; Op den Camp et al. 2003). In this respect, NADPH oxidase has been found to be an important generator of ROS (Lamb and Dixon 1997; Li et al. 2006; Torres et al. 2002). On the other hand, lack of correlation has also been reported in some cases (Dorey et al. 1999; Glazner et

al. 1996; Repka 2002; Torres et al. 2005). This discrepancy in results illustrate that, although there is overwhelming evidence that ROS accumulation plays a central role for the HR, we do not yet understand the process or processes in detail and different pathways may operate in different systems or under different conditions. Elucidation of the causal relation between ROS and HR is further complicated by the fact that, for example, plant hormones such as SA, JA, ET and abscisic acid also influence the elicitation and expression of HR (Hoeberichts and Woltering 2003; Torres et al. 2005; Van Breusegem and Dat 2006).

Different models have been proposed to explain how ROS (H_2O_2 , O_2^- , $^1\text{O}_2^-$) and NO may elicit and regulate HR (e.g., Delledonne et al. 2002; Torres et al. 2005; Van Breusegem and Dat 2006). Thus, when plants are subjected to stress and ROS accumulate at levels insufficient to kill the cell (as opposed to necrosis which is passive, accidental cell death), signalling events lead to PCD represented by HR. Initiation of HR may further lead to activation of other defence responses and systemic acquired resistance (Greenberg 1997; Van Breusegem and Dat 2006).

The exact role and mechanism of ROS in elicitation of the HR remains unclear. For example, it has been suggested from studies of soybean suspension-cultured cells that elicitation of HR requires tightly balanced production of H_2O_2 and NO where NO reacts with H_2O_2 (generated from O_2^- by SOD) and elicits the HR (Delledonne et al. 2001), although it has been questioned whether this is a general phenomenon (Greenberg and Yao 2004). In wheat infected by the hemibiotrophic pathogen *S. tritici*, there was a strong accumulation of H_2O_2 in a resistant cultivar, coinciding with the restriction of pathogen growth and expression of defence genes, but there is no classical HR in the host (Shetty et al. 2003, 2007). Likewise, Sasabe et al. (2000) also found that elicitor treatment of tobacco cell suspension cultures resulted in an oxidative burst, but not in cell death or defence gene activation. This indicates that signalling pathways leading to the oxidative burst, cell death and defence gene activation may branch at an early stage. Also Montillet et al. (2005) found that an HR could be elicited in tobacco by different pathways in light and darkness. There are also reports where elicitors and pathogens have been shown to trigger a strong oxidative burst without causing an HR but activate

other defence mechanisms involved with the oxidative burst (Glazner et al. 1996; Jabs et al. 1997; Repka 2002). For example, Glazner et al. (1996) showed that ROS accumulation in tobacco leaves and cultured cells in response to an incompatible strain of *Pseudomonas syringae* pv. *syringae* was not sufficient to cause HR. Likewise, in parsley suspension-cultured cells inoculated with fungal elicitor, ROS production and activation of defence genes was observed, but no HR, indicating that ROS accumulation could play other roles, such as to act as a direct antimicrobial agent, for induction of defence gene expression and phytoalexin accumulation in the absence of the HR (Jabs et al. 1997).

Clearly, the role of ROS for triggering and executing the HR is complicated and influenced by many factors. Already Levine et al. (1994) suggested that ROS at interaction sites may have different roles in the elicitation or prevention of cell death depending on their concentration, sub-cellular localisation and the duration of their production.

Involvement of ROS in successful pathogenesis

Biotrophic pathogens obtain their nutrition from living host cells (Oliver and Ipcho 2004), and H_2O_2 has been reported as an effective factor in stopping growth of biotrophic pathogens such as *B. graminis* f. sp. *hordei* (Mellersh et al. 2002; Thordal-Christensen et al. 1997; Trujillo et al. 2006). For example, Trujillo et al. (2006) found in the barley–*Blumeria* interaction that H_2O_2 is produced in non-penetrated CWA (see Fig. 2b). However, they also showed that superoxide (O_2^-) was produced locally at the site of penetration and appeared to enhance infection, thus suggesting that ROS can also be important for the pathogenesis of biotrophic pathogens. Biotrophic pathogens may suppress the host defence responses during infection (Ferreira et al. 2007). For example, the fungal metabolite mannitol, which can suppress ROS-related defence mechanisms by scavenging ROS, was found in apoplastic fluids of *Vicia faba* leaves infected with *Uromyces viciae-fabae* (formerly *Uromyces fabae*; Link et al. 2005; Voegelé et al. 2005). On the other hand, HR and ROS such as H_2O_2 have been reported to benefit infection by necrotrophic pathogens, which may even be able to produce ROS themselves or stimulate the host to do so (Govrin and Levine 2000; Von Gönner and Schlösser 1992). For example, Van

der Vlugt-Bergmans et al. (1997) studied *B. fuckeliana* infection in *V. faba* and reported that the fungus released H_2O_2 which could destroy host membrane lipids, thereby facilitating penetration. *B. fuckeliana* was also reported to enhance ROS production to aid its tissue colonization by triggering changes in the host (tomato) peroxisomal antioxidant system, leading to a collapse of regulatory mechanisms (Kuźniak and Skłodowska 2005). In vitro studies by Gil-ad and Mayer (1999) showed that *B. fuckeliana* spores could germinate at a concentration of about 180 mM H_2O_2 and that the mycelium could reduce the H_2O_2 present in the culture medium, thus clearly demonstrating that the pathogen could cope with this high H_2O_2 concentration. In contrast, Unger et al. (2005) performed studies in bean leaves and suspension-cultured cells, which indicated that a non-aggressive isolate of *B. fuckeliana* was in fact inhibited by ROS (O_2^- and partly H_2O_2) and HR. On the other hand, an aggressive isolate induced HR-like necrosis and was able to complete its life cycle. The aggressive isolate produced high amounts of a suppressor of O_2^- , i.e., 2-methyl succinate. When this suppressor was added to the non-aggressive isolate, enhanced tissue necrosis occurred. These results demonstrate a situation equivalent to biotrophic pathogens suppressing the host defence responses and indicate that ROS may also inhibit the necrotrophic pathogen *B. fuckeliana* in some cases (c.f. Małolepsza and Urbanek 2002), contrary to previous conclusions regarding this organism.

The so-called hemibiotrophic pathogens are a diverse group of organisms with an initial biotrophic phase where infection is established, followed by a necrotrophic phase where the pathogen completes its life-cycle (Oliver and Ipcho 2004). A correlation between pathogen growth at the late stages of their life-cycle and large quantities of H_2O_2 has also been reported in such host-pathogen systems. Thus, Able (2003) reported that in barley infected by *Rhynchosporium secalis* and *P. teres*, there was a large accumulation of H_2O_2 in compatible interactions in the later stage of infection, and it was concluded that H_2O_2 was necessary for successful infection as for *B. fuckeliana* (see above), however, based only on correlative evidence. Recently, doubts have been raised whether this is a valid conclusion. Shetty et al. (2003) observed a similar correlation in wheat

infected by the hemibiotrophic pathogen *S. tritici*. During the biotrophic phase of the interaction, H₂O₂ accumulation occurred as a defence response only in an incompatible interaction (Fig. 2c). On the other hand, in a compatible interaction, large amounts of H₂O₂ accumulated after extensive tissue colonization just before the symptoms appeared and the pathogen sporulated (Fig. 2d). However, Shetty et al. (2007) found that, even though sporulation of *S. tritici* in wheat coincided with a massive accumulation of H₂O₂, removal of this H₂O₂ by infiltration of catalase resulted in increased pathogen growth, indicating both that it can survive and tolerate the presence of H₂O₂, but also that this H₂O₂ inhibits the pathogen.

The ability to colonise and proliferate in an environment with high concentrations of ROS shows that the pathogens have efficient systems enabling them to protect themselves against the harsh environment in the host. Thus, for the necrotrophic pathogen *B. fuckeliana*, Van der Vlugt-Bergmans et al. (1997) found that it could protect itself by expressing genes encoding catalase which could scavenge H₂O₂. In accordance with this, Goodwin et al. (2001) cloned a catalase gene from the hemibiotrophic pathogen *Colletotrichum gloeosporioides* f.sp. *malvae* (pathogen of *Malva pusilla*). Catalase genes have also been reported from *S. tritici* (Levy et al. 1992), but their expression was not studied. However, it can be predicted that catalases will be activated during the necrotrophic phase of *S. tritici* growth to help protect the pathogen from the deleterious effect of H₂O₂. In agreement with this, Shetty et al. (2003) found high levels of catalase activity in a susceptible host during pycnidial formation, but were unable to determine whether this was of host or pathogen origin.

Other types of hemibiotrophic pathogens might benefit from ROS accumulation at some stage. Thus, Kumar et al. (2001) reported that barley with *mlo* resistance against *B. graminis* f.sp. *hordei* was very susceptible to *Bipolaris sorokiniana*. Toxins from *B. sorokiniana* killed the host cells, leading to massive H₂O₂ accumulation, and it was hypothesised that the *mlo*-resistant plants were very susceptible because cell death occurred more easily compared to other barley genotypes. After cell death, the pathogen could grow unhindered in the dead tissue and was not inhibited by H₂O₂ accumulation (cf. Fig. 1). However, in the initial stages of penetration, it was concluded that *B.*

sorokiniana was inhibited by H₂O₂ accumulation just beneath appressoria from which penetration was attempted, i.e., before cells died.

Most conclusions regarding the role of ROS in successful pathogenesis are based solely on correlative data and come from rather few pathosystems. *B. fuckeliana* is most often used as a representative necrotrophic pathogen. The influence of ROS on this pathogen is fairly well studied even though conflicting results have been reported, but generalisations to other pathogens should be made with caution. Thus, as pointed out by Shetty et al. (2007), even if there is a correlation between ROS accumulation and pathogen colonisation, the fact that the pathogen can tolerate the presence of large amounts of ROS does not necessarily mean that it benefits from ROS. Furthermore, there is disagreement as to which categories different pathogens belong (see, e.g., Oliver and Ipcho 2004). For example, *R. secalis* was reported to be necrotrophic (Able 2003), whereas previous research has shown this pathogen to have a long symptomless phase (Jørgensen et al. 1993), suggesting that it should be considered as hemibiotrophic. Therefore, caution should also be taken when concluding about a definite role of ROS for specific types or even species of pathogens before thorough studies have been conducted.

Discussion

Although our understanding of the oxidative burst in plant–pathogen interactions has advanced considerably since the first reports, there are still several unanswered questions. Thus, Fig. 1 shows an overview of our current knowledge of the different roles of ROS in host–pathogen interactions, but also indicates some of the areas where there are unanswered questions and gaps in our knowledge.

The rapid production of ROS in the apoplast in response to pathogens has been proposed to orchestrate the establishment of different defensive barriers against pathogens (Torres et al. 2006). Our understanding of sources and roles of ROS has been greatly enhanced by the identification of defence-associated mutants in the model plant *A. thaliana* (Lorrain et al. 2003). These mutants have not only allowed the identification of important signalling intermediates but also allowed the dissection of ROS-mediated

signalling pathways. However, the influence of other factors such as environment, plant hormones and activation of different signalling pathways (Op den Camp et al. 2003; Montillet et al. 2005; Sasabe et al. 2000; Torres et al. 2005) plays an important role for the accumulation of ROS, and this needs to be taken into consideration and studied in detail. Likewise, external factors such as different types of pathogens and elicitors may vary in their ability to trigger ROS production (cf. Fig. 1), and are therefore possible reasons for conflicting results. This emphasises the need for studies of several different host–pathogen systems in order to clarify if and when different pathways are activated in different situations. It is therefore essential to study several different hosts infected by taxonomically different pathogens which represent different life-style strategies before making general conclusions and thus avoid over-simplification. There are profound differences between monocots and dicots as well as in the biology of biotrophic, hemi-biotrophic and necrotrophic pathogens. Caution should therefore be exercised before stating that processes occur in a similar way in totally different systems.

It is also important to adopt different approaches to increase the robustness of conclusions as illustrated by the example of involvement of ROS in HR. Besides a genetic approach, using mutants, gene silencing, gene knock-outs and/or over-expression, careful physiological and biochemical characterisation of different host–pathogen interactions and defence responses activated should be carried out followed by studies of the role of proteins encoded by ROS genes in the different systems. This approach provides insight into their precise function in defence, cell death, and/or pathogen development, through determination of their sub-cellular localisation and biochemical function. In particular, in relation to the evaluation of the role of ROS in successful pathogenesis, it is important to try to inhibit the cell death machinery selectively and simultaneously to monitor other defence and pathogenesis-related events. Using this approach, it should be possible to determine whether cell death can be uncoupled from other defence responses and if so, the specific contribution to resistance or susceptibility in the interaction in question. Of particular interest in this context is to determine which role ROS plays in HR/necrosis against necrotrophic pathogens (cf. Fig. 1). Thus, do these pathogens all benefit from

ROS accumulation or are some of them actually inhibited to some extent by ROS or other substances in the dying cells? Another important question regarding necrotrophic pathogens relates to potential toxins produced (cf. Fig. 1). Thus, will those toxins which kill host cells always cause the release of ROS, which in turn causes an HR (cf. *B. sorokiniana*)? It is also important to study the interplay between ROS and SA/NO, in order to gain further insights into the regulation of resistance, as these are important defence response regulators that interact with ROS signalling in response to pathogens (Mur et al. 2006). Thus, ROS may be part of many signalling pathways and provide a crucial link in the cross-talk to different responses (Apel and Hirt 2004).

The flux of information between different cell compartments needs to be elucidated to further understand the regulatory capabilities of ROS. Previously, genetic engineering for improved disease resistance has mainly targeted genes involved in the recognition of the pathogen or in the over-expression of defence molecules like phytoalexins (Jalali et al. 2006). An interesting alternative approach would be to target key molecules like ROS that act at points of convergence of different signalling pathways. Engineering plants with such genes using a pathogen-inducible promoter would enable expression of different downstream genes simultaneously in the host, ensuring that plants develop an array of effective responses, which will ultimately secure a sustainable and durable resistance against a range of plant pathogens.

References

- Able, A. J. (2003). Role of reactive oxygen species in the response of barley to necrotrophic pathogens. *Protoplasma*, 221, 137–143.
- Ali, R., Ma, W., Lemtiri-Chlieh, F., Tsaltas, D., Leng, Q., von Bodman, S., et al. (2007). Death don't have no mercy and neither does calcium: *Arabidopsis* CYCLIC NUCLEOTIDE GATED CHANNEL2 and innate immunity. *The Plant Cell*, 19, 1081–1095.
- Allan, A. C., & Fluhr, R. (1997). Two distinct sources of elicited reactive oxygen species in tobacco epidermal cells. *The Plant Cell*, 9, 1559–1572.
- Alvarez, M. E., Pennell, R. I., Meijer, P.-J., Ishikawa, A., Dixon, R. A., & Lamb, C. (1998). Reactive oxygen intermediates mediate a systemic signal network in the establishment of plant immunity. *Cell*, 92, 773–784.

- Amirsadeghi, S., Robson, C. A., & Vanlerberghe, G. C. (2007). The role of the mitochondrion in plant responses to biotic stress. *Physiologia Plantarum*, *129*, 253–266.
- An, Q., Ehlers, K., Kogel, K.-H., van Bel, A. J. E., & Hükelhoven, R. (2006). Multivesicular compartments proliferate in susceptible and resistant *MLA12*-barley leaves in response to infection by the biotrophic powdery mildew fungus. *New Phytologist*, *172*, 563–576.
- Apel, K., & Hirt, H. (2004). Reactive oxygen species: Metabolism, oxidative stress, and signal transduction. *Annual Review of Plant Biology*, *55*, 373–399.
- Apostol, I., Heinstein, P. F., & Low, P. S. (1989). Rapid induction of an oxidative burst during elicitation of cultured plant cells. *Plant Physiology*, *90*, 109–116.
- Ashtamker, C., Kiss, V., Sagi, M., Davydov, O., & Fluhr, R. (2007). Diverse subcellular locations of cryptogein-induced reactive oxygen species production in tobacco Bright Yellow-2 cells. *Plant Physiology*, *143*, 1817–1826.
- Auh, C. K., & Murphy, T. M. (1995). Plasma-membrane redox enzyme is involved in the synthesis of O₂⁻ and H₂O₂ by *Phytophthora* elicitor-stimulated rose cells. *Plant Physiology*, *107*, 1241–1247.
- Babitha, M. P., Prakash, H. S., & Shetty, H. S. (2004). Purification and properties of lipoxygenase induced in downy mildew resistant pearl millet seedlings due to infection with *Sclerospora graminicola*. *Plant Science*, *166*, 31–39.
- Baker, C. J., & Orlandi, E. W. (1995). Active oxygen in plant pathogenesis. *Annual Review of Phytopathology*, *33*, 299–321.
- Baker, C. J., Orlandi, E. W., & Mock, N. M. (1993). Harpin, an elicitor of the hypersensitive response in tobacco caused by *Erwinia amylovora*, elicits active oxygen production in suspension cells. *Plant Physiology*, *102*, 1341–1344.
- Bedard, K., Lardy, B., & Krause, K.-H. (2007). NOX family NADPH oxidases: Not just in mammals. *Biochimie*, *89*, 1107–1112.
- Bindschedler, L. V., Dewdney, J., Blee, K. A., Stone, J. M., Asai, T., Plotnikov, J., et al. (2006). Peroxidase-dependent apoplastic oxidative burst in *Arabidopsis* required for pathogen resistance. *The Plant Journal*, *47*, 851–863.
- Blumwald, E., Aharon, G. S., & Lam, B. C.-H. (1998). Early signal transduction pathways in plant–pathogen interactions. *Trends in Plant Science*, *3*, 342–346.
- Bolwell, G. P., Bindschedler, L. V., Blee, K. A., Butt, V. S., Davies, D. R., Gardner, S. L., et al. (2002). The apoplastic oxidative burst in response to biotic stress in plants: A three-component system. *Journal of Experimental Botany*, *53*, 1367–1376.
- Bradley, D. J., Kjellbom, P., & Lamb, C. J. (1992). Elicitor- and wound-induced oxidative cross-linking of a proline-rich plant cell wall structural protein: A novel, rapid plant defense response. *Cell*, *70*, 21–30.
- Carter, C., Healy, R., O'Tool, N. M., Naqvi, S. M. S., Ren, G., Park, S., et al. (2007). Tobacco nectaries express a novel NADPH oxidase implicated in the defense of floral reproductive tissues against microorganisms. *Plant Physiology*, *143*, 389–399.
- Chandra, S., Martin, G. B., & Low, P. S. (1996). The Pto kinase mediates a signaling pathway leading to the oxidative burst in tomato. *Proceedings of the National Academy of Sciences of the United States of America*, *93*, 13393–13397.
- Chisholm, S. T., Coaker, G., Day, B., & Staskawicz, B. J. (2006). Host–microbe interactions: Shaping the evolution of the plant immune response. *Cell*, *124*, 803–814.
- Collins, N. C., Thordal-Christensen, H., Lipka, V., Bau, S., Kombrink, E., Qiu, J. L., et al. (2003). SNARE-protein-mediated disease resistance at the plant cell wall. *Nature*, *425*, 973–977.
- Cona, A., Rea, G., Angelini, R., Federico, R., & Tavaladorak, P. (2006). Functions of amine oxidases in plant development and defence. *Trends in Plant Science*, *11*, 80–89.
- Custers, J. H. H. V., Harrison, S. J., Sela-Buurlage, M. B., van Deventer, E., Lageweg, W., Howe, P. W., et al. (2004). Isolation and characterisation of a class of carbohydrate oxidases from higher plants, with a role in active defence. *Plant Journal*, *39*, 147–160.
- Dat, J. F., Pellinen, R., Beeckman, T., Van de Cotte, B., Langebartels, C., Kangasjärvi, J., et al. (2003). Changes in hydrogen peroxide homeostasis trigger an active cell death process in tobacco. *The Plant Journal*, *33*, 621–632.
- Deepak, S. A., Ishii, H., & Park, P. (2006). Acibenzolar-S-methyl primes cell wall strengthening genes and reactive oxygen species forming/scavenging enzymes in cucumber after fungal pathogen attack. *Physiological and Molecular Plant Pathology*, *69*, 52–61.
- Delledonne, M., Murgia, I., Ederle, D., Sbicego, P. F., Biondani, A., Polverari, A., et al. (2002). Reactive oxygen intermediates modulate nitric oxide signaling in the plant hypersensitive disease-resistance response. *Plant Physiology and Biochemistry*, *40*, 605–610.
- Delledonne, M., Xia, Y., Dixon, R. A., & Lamb, C. (1998). Nitric oxide functions as a signal in plant disease resistance. *Nature*, *394*, 585–588.
- Delledonne, M., Zeier, J., Marocco, A., & Lamb, C. (2001). Signal interactions between nitric oxide and reactive oxygen intermediates in the plant hypersensitive disease resistance response. *Proceedings of the National Academy of Sciences of the United States of America*, *98*, 13454–13459.
- Desikan, R., Clarke, A., Hancock, J. T., & Neill, S. J. (1999). H₂O₂ activates a MAP kinase-like enzymes in *Arabidopsis thaliana* suspension cultures. *Journal of Experimental Botany*, *50*, 1863–1866.
- Dorey, S., Kopp, M., Geoffroy, P., Fritig, B., & Kauffmann, S. (1999). Hydrogen peroxide from the oxidative burst is neither necessary nor sufficient for hypersensitive cell death induction, phenylalanine ammonia lyase stimulation, salicylic acid accumulation or scopoletin consumption in cultured tobacco cells treated with elicitor. *Plant Physiology*, *121*, 163–171.
- Enyedi, A. J., Yalpani, N., Silverman, P., & Raskin, I. (1992). Localization, conjugation, and function of salicylic acid in tobacco during the hypersensitive reaction to tobacco mosaic virus. *Proceedings of the National Academy of Sciences of the United States of America*, *89*, 2480–2484.
- Ferreira, R. B., Monteiro, S., Freitas, R., Santos, C. N., Chen, Z., Batista, L. M., et al. (2007). The role of plant defence proteins in fungal pathogenesis. *Molecular Plant Pathology*, *8*, 677–700.
- Floryszak-Wieczorek, J., Arasimowicz, M., Milczarek, G., Jelen, H., & Jackowiak, H. (2007). Only an early nitric

- oxide burst and the following wave of secondary nitric oxide generation enhanced effective defence responses of pelargonium to a necrotrophic pathogen. *New Phytologist*, *175*, 718–730.
- Gil-ad, N. L., & Mayer, A. M. (1999). Evidence for rapid breakdown of hydrogen peroxide by *Botrytis cinerea*. *FEMS Microbiology Letters*, *176*, 455–461.
- Glazner, J. A., Orlandi, E. W., & Baker, C. J. (1996). The active oxygen response of cell suspensions to incompatible bacteria is not sufficient to cause hypersensitive cell death. *Plant Physiology*, *110*, 759–763.
- Goodwin, P. H., Li, J., & Jin, S. (2001). A catalase gene of *Colletotrichum gloeosporioides* f. sp. *malvae* is highly expressed during the necrotrophic phase of infection of round-leaved mallow, *Malva pusilla*. *FEMS Microbiology Letters*, *202*, 103–107.
- Govrin, E. M., & Levine, A. (2000). The hypersensitive response facilitates plant infection by the necrotrophic pathogen *Botrytis cinerea*. *Current Biology*, *10*, 751–757.
- Grant, J. J., Yun, B.-W., & Loake, G. J. (2000). Oxidative burst and cognate redox signalling reported by luciferase imaging: Identification of a signal network that functions independently of ethylene, SA and Me-JA but is dependent on MAPKK activity. *The Plant Journal*, *24*, 569–582.
- Greenberg, J. T. (1997). Programmed cell death in plant–pathogen interactions. *Annual Review of Plant Physiology and Plant Molecular Biology*, *48*, 525–545.
- Greenberg, J. T., & Yao, N. (2004). The role and regulation of programmed cell death in plant–pathogen interactions. *Cellular Microbiology*, *6*, 201–211.
- Heitefuss, R. (1997). Cell wall modification in relation to resistance. In H. Hartleb, R. Heitefuss, & H.-H. Hoppe (Eds.), *Resistance of crop plants against fungi* (pp. 100–125). Jena: Gustav Fischer.
- Hirt, H. (1997). Multiple roles of MAP kinases in plant signal transduction. *Trends in Plant Science*, *2*, 11–15.
- Hoeberichts, F. A., ten Have, A., & Woltering, E. J. (2003). A tomato metacaspase gene is upregulated during programmed cell death in *Botrytis cinerea*-infected leaves. *Planta*, *217*, 517–522.
- Hoeberichts, F. A., & Woltering, E. J. (2003). Multiple mediators of plant programmed cell death: Interplay of conserved cell death mechanisms and plant-specific regulators. *BioEssays*, *25*, 47–57.
- Hu, X., Bidney, D. L., Yalpani, N., Duvick, J. P., & Crasta, O. (2003). Overexpression of a gene encoding hydrogen peroxide-generating oxalate oxidase evokes defense responses in sunflower. *Plant Physiology*, *133*, 170–181.
- Hückelhoven, R., & Kogel, K.-H. (2003). Reactive oxygen intermediates in plant microbe interactions: Who is who in powdery mildew resistance? *Planta*, *216*, 891–902.
- Iwai, T., Seo, S., Mitsuhara, I., & Ohashi, Y. (2007). Probenazole-induced accumulation of salicylic acid confers resistance to *Magnaporthe grisea* in adult rice plants. *Plant and Cell Physiology*, *48*, 915–924.
- Iwano, M., Che, F.-S., Goto, K., Tanaka, N., Takayama, S., & Isogai, A. (2002). Electron microscopic analysis of the H₂O₂ accumulation preceding hypersensitive cell death induced by an incompatible strain of *Pseudomonas avenae* in cultured rice cells. *Molecular Plant Pathology*, *3*, 1–8.
- Jabs, T., Dietrich, R. A., & Dangl, J. L. (1996). Initiation of runaway cell death in an *Arabidopsis* mutant by extracellular superoxide. *Science*, *273*, 1853–1856.
- Jabs, T., Tschöpe, M., Colling, C., Hahlbrock, K., & Scheel, D. (1997). Elicitor-stimulated ion fluxes and O₂⁻ from the oxidative burst are essential components in triggering defense gene activation and phytoalexin synthesis in parsley. *Proceedings of the National Academy of Sciences of the United States of America*, *94*, 4800–4805.
- Jalali, B. L., Bhargava, S., & Kamble, A. (2006). Signal transduction and transcriptional regulation of plant defence responses. *Journal of Phytopathology*, *154*, 65–74.
- Jørgensen, H. J. L., de Neergaard, E., & Smedegaard-Petersen, V. (1993). Histological examination of the interaction between *Rhynchosporium secalis* and susceptible and resistant cultivars of barley. *Physiological and Molecular Plant Pathology*, *42*, 345–358.
- Jørgensen, H. J. L., Lübeck, P. S., Thordal-Christensen, H., de Neergaard, E., & Smedegaard-Petersen, V. (1998). Mechanisms of induced resistance in barley against *Drechslera teres*. *Phytopathology*, *88*, 698–707.
- Kariola, T., Brader, G., Li, J., & Palva, E. T. (2005). Chlorophyllose 1, a damage control enzyme, affects the balance between defense pathways in plants. *The Plant Cell*, *17*, 282–294.
- Kawasaki, T., Henmi, K., Ono, E., Hatakeyama, S., Iwano, M., Satoh, H., et al. (1999). The small GTP-binding protein Rac is a regulator of cell death in plants. *Proceedings of the National Academy of Sciences of the United States of America*, *96*, 10922–10926.
- Klessig, D. F., Durner, J., Noad, R., Navarre, D. A., Wendehenne, D., Kumar, D., et al. (2000). Nitric oxide and salicylic acid signaling in plant defense. *Proceedings of the National Academy of Sciences of the United States of America*, *97*, 8849–8855.
- Kobayashi, M., Ohura, I., Kawakita, K., Yokota, N., Fujiwara, M., Shimamoto, K., et al. (2007). Calcium-dependent protein kinases regulate the production of reactive oxygen species by potato NADPH oxidase. *The Plant Cell*, *19*, 1065–1080.
- Kumar, J., Hückelhoven, R., Beckhove, U., Nagarajan, S., & Kogel, K.-H. (2001). A compromised Mlo pathway affects the response of barley to the necrotrophic fungus *Bipolaris sorokiniana* (teleomorph: *Cochliobolus sativus*) and its toxins. *Phytopathology*, *91*, 127–133.
- Kumudini, B. S., & Shetty, H. S. (2002). Association of lignification and callose deposition with host cultivar resistance and induced systemic resistance of pearl millet to *Sclerospora graminicola*. *Australasian Plant Pathology*, *32*, 157–164.
- Kuźniak, E., & Skłodowska, M. (2005). Fungal pathogen-induced changes in the antioxidant systems of leaf peroxisomes from infected tomato plants. *Planta*, *222*, 192–200.
- Lam, E. (2004). Controlled cell death, plant survival and development. *Nature Reviews in Molecular Cell Biology*, *5*, 305–315.
- Lam, E., Kato, N., & Lawton, M. (2001). Programmed cell death, mitochondria and the plant hypersensitive response. *Nature*, *411*, 848–853.

- Lamb, C., & Dixon, R. A. (1997). The oxidative burst in plant disease resistance. *Annual Review of Plant Physiology and Plant Molecular Biology*, *48*, 251–275.
- Legendre, L., Heinstejn, P. F., & Low, P. S. (1992). Evidence for the participation of GTP-binding proteins in the elicitation of rapid oxidative burst in cultured soybean cells. *Journal of Biological Chemistry*, *267*, 20140–20147.
- Legendre, L., Rueter, S., Heinstejn, P. S., & Low, P. S. (1993). Characterisation of the oligogalacturonide-induced oxidative burst in cultured soybean (*Glycine max*) cells. *Plant Physiology*, *102*, 233–240.
- Levine, A., Tenhaken, R., Dixon, R., & Lamb, C. (1994). H₂O₂ from the oxidative burst orchestrates the plant hypersensitive disease resistance response. *Cell*, *79*, 583–593.
- Levy, E., Eyal, Z., & Hochman, A. (1992). Purification and characterization of a catalase–peroxidase from the fungus *Septoria tritici*. *Archives of Biochemistry and Biophysics*, *296*, 321–327.
- Li, A. L., Wang, M. L., Zhou, R. H., Kong, X. Y., Huo, N. X., Wang, W. S., et al. (2005). Comparative analysis of early H₂O₂ accumulation in compatible and incompatible wheat–powdery mildew interactions. *Plant Pathology*, *54*, 308–316.
- Li, J., Zhang, Z.-G., Ji, R., Wang, Y.-C., & Zheng, X.-B. (2006). Hydrogen peroxide regulates elicitor PB90-induced cell death and defense in non-heading Chinese cabbage. *Physiological and Molecular Plant Pathology*, *67*, 220–230.
- Link, T., Lohaus, G., Heiser, I., Mendgen, K., Hahn, M., & Voegelé, R. T. (2005). Characterization of a novel NADP⁺-dependent D-arabitol dehydrogenase from the plant pathogen *Uromyces fabae*. *Biochemical Journal*, *389*, 289–295.
- Liu, G., Greenshields, D. L., Sammynaiken, R., Hirji, R. N., Selvaraj, G., & Wei, Y. (2007). Targeted alterations in iron homeostasis underlie plant defense responses. *Journal of Cell Science*, *120*, 596–605.
- Lorrain, S., Vailliau, F., Balagué, C., & Roby, D. (2003). Lesion mimic mutants: Keys for deciphering cell death and defense pathways in plants? *Trends in Plant Science*, *8*, 263–271.
- Malolepsza, U. (2005). Spatial and temporal variation of reactive oxygen species and antioxidant enzymes in *o*-hydroxyethylrutin-treated tomato leaves inoculated with *Botrytis cinerea*. *Plant Pathology*, *54*, 317–324.
- Malolepsza, U., & Urbanek, H. (2002). *o*-Hydroxyethylrutin-mediated enhancement of tomato resistance to *Botrytis cinerea* depends on a burst of reactive oxygen species. *Journal of Phytopathology*, *150*, 616–624.
- Mateo, A., Mühlenbock, P., Rustérucchi, C., Chang, C. C.-C., Miszalski, Z., Karpinska, B., et al. (2004). *LESION SIMULATING DISEASE 1* is required for acclimation to conditions that promote excess excitation energy. *Plant Physiology*, *136*, 2818–2830.
- Mayer, A. M., Staples, R. C., & Gil-ad, N. L. (2001). Mechanisms of survival of necrotrophic fungal plant pathogens in hosts expressing the hypersensitive response. *Phytochemistry*, *58*, 33–41.
- McAinsh, M. R., Clayton, H., Mansfield, T. A., & Hetherington, A. M. (1996). Changes in stomatal behavior and guard cell cytosolic free calcium in response to oxidative stress. *Plant Physiology*, *111*, 1031–1042.
- McDowell, J. M., & Dangl, J. L. (2000). Signal transduction in the plant immune response. *Trends in Biochemical Science*, *25*, 79–82.
- Mellersh, D. G., Foulds, I. V., Higgins, V. J., & Heath, M. C. (2002). H₂O₂ plays different roles in determining penetration failure in three diverse plant–fungal interactions. *The Plant Journal*, *29*, 257–268.
- Mittler, R., Herr, E. H., Orvar, B. L., van Camp, W., Willekens, H., Inzé, D., et al. (1999). Transgenic tobacco plants with reduced capability to detoxify reactive oxygen intermediates are hyperresponsive to pathogen infection. *Proceedings of the National Academy of Sciences of the United States of America*, *96*, 14165–14170.
- Mittler, R., Vanderauwera, S., Gollery, M., & Van Breusegem, F. (2004). The reactive oxygen gene network in plants. *Trends in Plant Science*, *9*, 490–498.
- Moerschbacher, B. M., & Reiser, H.-J. (1997). The hypersensitive resistance reaction. In H. Hartleb, R. Heitefuss, & H.-H. Hoppe (Eds.), *Resistance of crop plants against fungi* (pp. 126–158). Jena: Gustav Fischer.
- Montillet, J.-L., Chamnongpol, S., Rustérucchi, C., Dat, J., Van de Cotte, B., Agnel, J.-P., et al. (2005). Fatty acid hydroperoxides and H₂O₂ in the execution of hypersensitive cell death in tobacco leaves. *Plant Physiology*, *138*, 1516–1526.
- Mur, L. A. J., Carver, T. L. W., & Prats, E. (2006). NO way to live; the various roles of nitric oxide in plant–pathogen interactions. *Journal of Experimental Botany*, *57*, 489–505.
- Neill, S. J., Desikan, R., Clarke, A., Hurst, R. D., & Hancock, J. T. (2002). Hydrogen peroxide and nitric oxide as signalling molecules in plants. *Journal of Experimental Botany*, *53*, 1237–1247.
- Numberger, T. M., Nennsteil, O., Jabs, T., Sacks, W. R., Hahlbrock, K., & Scheel, D. (1994). High affinity binding of a fungal oligopeptide elicitor to parsley plasma membranes triggers multiple defense responses. *Cell*, *78*, 449–460.
- Oliver, R. P., & Ipcho, S. V. S. (2004). *Arabidopsis* pathology breathes new life into the necrotrophs-vs.-biotrophs classification of fungal pathogens. *Molecular Plant Pathology*, *4*, 347–352.
- Olson, P. D., & Varner, J. E. (1993). Hydrogen peroxide and lignification. *The Plant Journal*, *4*, 887–892.
- Op den Camp, R. G. L., Przybyla, D., Ochsenshein, C., Laloi, C., Kim, C., Danon, A., et al. (2003). Rapid induction of distinct stress responses after the release of singlet oxygen in *Arabidopsis*. *The Plant Cell*, *15*, 2320–2332.
- Pei, Z.-M., Murata, Y., Benning, G., Thomine, S., Klüsener, B., Allen, G. J., et al. (2000). Calcium channels activated by hydrogen peroxide mediate abscisic acid signalling in guard cells. *Nature*, *406*, 731–734.
- Peng, M., & Kuc, J. (1992). Peroxidase-generated hydrogen peroxide as a source of antifungal activity in vitro and on tobacco leaf disks. *Phytopathology*, *82*, 696–699.
- Petersen, M., Brodersen, P., Naested, H., Andreasson, E., Lindhart, U., Johansen, B., et al. (2000). *Arabidopsis*

- MAP kinase 4 negatively regulates systemic acquired resistance. *Cell*, *103*, 1111–1120.
- Price, A., Knight, M., Knight, H., Cuin, T., Tomos, D., & Ashenden, T. (1996). Cytosolic calcium and oxidative plant stress. *Biochemical Society Transactions*, *24*, 479–483.
- Price, A. H., Taylor, A., Ripley, S. J., Griffiths, A., Trewavas, A. J., & Knight, M. R. (1994). Oxidative signals in tobacco increase cytosolic calcium. *The Plant Cell*, *6*, 1301–1310.
- Ren, D., Yang, K.-Y., Li, G.-J., Liu, Y., & Zhang, S. (2006). Activation of Ntf4, a tobacco mitogen-activated protein kinase, during plant defence response and its involvement in hypersensitive response-like cell death. *Plant Physiology*, *141*, 1482–1493.
- Repka, V. (2002). Hydrogen peroxide generated via the octadecanoid pathway is neither necessary nor sufficient for methyl jasmonate-induced hypersensitive cell death in woody plants. *Biologia Plantarum*, *45*, 105–115.
- Sasabe, M., Takeuchi, K., Kamoun, S., Ichinose, Y., Govers, F., Toyoda, K., et al. (2000). Independent pathways leading to apoptotic cell death, oxidative burst and defense gene expression in response to elicitor in tobacco cell suspension culture. *European Journal of Biochemistry*, *267*, 5005–5013.
- Schulze-Lefert, P. (2004). Knocking on the heaven's wall: Pathogenesis of and resistance to biotrophic fungi at the cell wall. *Current Opinion in Plant Biology*, *7*, 377–383.
- Shah, J. (2003). The salicylic acid loop in plant defense. *Current Opinion in Plant Biology*, *6*, 365–371.
- Shailashree, S., Kini, K. R., Deepak, S., Kumudini, B. S., & Shetty, H. S. (2004). Accumulation of hydroxyproline-rich glycoproteins in pearl millet seedlings in response to *Sclerospora graminicola* infection. *Plant Science*, *167*, 1227–1234.
- Shetty, N. P., Kristensen, B. K., Newman, M.-A., Møller, K., Gregersen, P. L., & Jørgensen, H. J. L. (2003). Association of hydrogen peroxide with restriction of *Septoria tritici* in resistant wheat. *Physiological and Molecular Plant Pathology*, *62*, 333–346.
- Shetty, N. P., Mehrabi, R., Lütken, H., Haldrup, A., Kema, G. H. J., Collinge, D. B., et al. (2007). Role of hydrogen peroxide during the interaction between the hemibiotrophic fungal pathogen *Septoria tritici* and wheat. *New Phytologist*, *174*, 637–647.
- Suzuki, K. (2002). Map kinase cascade in elicitor signal transduction. *Journal of Plant Research*, *115*, 237–244.
- Thatcher, L. F., Anderson, J. P., & Singh, K. B. (2005). Plant defence responses: What have we learnt from *Arabidopsis*? *Functional Plant Biology*, *32*, 1–19.
- Thordal-Christensen, H., Zhang, Z., Wei, Y., & Collinge, D. B. (1997). Subcellular localization of H₂O₂ in plants. H₂O₂ accumulation in papillae and hypersensitive response during the barley–powdery mildew interaction. *The Plant Journal*, *11*, 1187–1194.
- Torres, M. A., Dangl, J. L., & Jones, J. D. G. (2002). *Arabidopsis* gp91^{phox} homologues, *AtrbohD* and *AtrbohF* are required for accumulation of reactive oxygen intermediates in the plant defense response. *Proceedings of the National Academy of Sciences of the United States of America*, *99*, 517–522.
- Torres, M. A., Jones, J. D. G., & Dangl, J. L. (2005). Pathogen-induced, NADPH oxidase-derived reactive oxygen intermediates suppress spread of cell death in *Arabidopsis thaliana*. *Nature Genetics*, *37*, 1130–1134.
- Torres, M. A., Jones, J. D. G., & Dangl, J. L. (2006). Reactive oxygen species signaling in response to pathogens. *Plant Physiology*, *141*, 373–378.
- Trujillo, M., Altschmeid, L., Schweizer, P., Kogel, K.-H., & Hüchelhoven, R. (2006). *Respiratory Burst Oxidase Homologue A* of barley contributes to penetration by the powdery mildew fungus *Blumeria graminis* f. sp. *hordei*. *Journal of Experimental Botany*, *57*, 3781–3791.
- Unger, C., Kleta, S., Jandl, G., & v. Tiedemann, A. (2005). Suppression of the defence-related oxidative burst in bean leaf tissue and bean suspension cells by the necrotrophic pathogen *Botrytis cinerea*. *Journal of Phytopathology*, *153*, 15–26.
- Urquhart, W., Gunawardena, A. H. L. A. N., Moeder, W., Ali, R., Berkowitz, G. A., & Yoshioka, K. (2007). The chimeric cyclic nucleotide-gated ion channel ATCNGC11/12 constitutively induces programmed cell death in a Ca²⁺ dependent manner. *Plant Molecular Biology*, *65*, 747–761.
- Van Breusegem, F., & Dat, J. F. (2006). Reactive oxygen species in plant cell death. *Plant Physiology*, *141*, 384–390.
- Van der Vlugt-Bergmans, C. J. B., Wagemakers, C. A. M., Dees, D. C. T., & Van Kan, J. A. L. (1997). Catalase A from *Botrytis cinerea* is not expressed during infection on tomato leaves. *Physiological and Molecular Plant Pathology*, *50*, 1–15.
- Voegelé, R. T., Hahn, M., Lohaus, G., Link, T., Heiser, I., & Mendgen, K. (2005). Possible roles for mannitol and mannitol dehydrogenase in the biotrophic plant pathogen *Uromyces fabae*. *Plant Physiology*, *137*, 190–198.
- Von Gönner, M., & Schlösser, E. (1992). Effect of radical scavengers on pathogenesis in the host–parasite-system *Avena sativa*–*Drechslera avenae*. *Zeitschrift für Pflanzenkrankheiten und Pflanzenschutz*, *99*, 617–625.
- Walters, D. R. (2003). Polyamines and plant disease. *Phytochemistry*, *64*, 97–107.
- Wu, G. S., Short, B. J., Lawrence, E. B., Levine, E. B., Fitzsimmons, K. C., & Shah, D. M. (1995). Disease resistance conferred by expression of a gene encoding H₂O₂-generating glucose oxidase in transgenic potato plants. *The Plant Cell*, *7*, 1357–1368.
- Zhang, S. Q., & Klessig, D. F. (1998). Resistance gene N-mediated de novo synthesis and activation of a tobacco mitogen-activated protein kinase by *Tobacco mosaic virus*. *Proceedings of the National Academy of Sciences of the United States of America*, *95*, 7433–7438.
- Zhang, S., Liu, Y., & Klessig, D. F. (2000). Multiple levels of tobacco WIPK activation during the induction of cell death by fungal elicitors. *The Plant Journal*, *23*, 339–347.
- Zhou, F., Menke, F. L. H., Yoshioka, K., Moder, W., Shirano, Y., & Klessig, D. F. (2004). High humidity suppresses *ssi4*-mediated cell death and disease resistance upstream of MAP kinase activation, H₂O₂ production and defense gene expression. *The Plant Journal*, *39*, 920–932.
- Zimmermann, G., Baumlein, H., Mock, H. P., Himmelbach, A., & Schweizer, P. (2006). The multigene family encoding germin-like proteins of barley. Regulation and function in basal host resistance. *Plant Physiology*, *142*, 181–192.
- Zwerger, K., & Hirt, H. (2001). Recent advances in plant MAP kinase signalling. *Biological Chemistry*, *382*, 1123–1131.