

## Pleistocene oceanographic changes indicated by deep sea benthic foraminifera in the northern Indian Ocean

AJAI K RAI\* and M S SRINIVASAN

\*Department of Earth and Planetary Sciences, Allahabad University, Allahabad 211 002, India

Department of Geology, Banaras Hindu University, Varanasi 221 005, India

MS received 18 July 1994; revised 5 October 1994

**Abstract.** An attempt has been made to understand the Pleistocene bottom water history in response to the paleoclimatic changes in the northern Indian Ocean employing quantitative analyses of deep sea benthic foraminifera at the DSDP sites 219 and 238. Among the 150 benthic foraminifera recorded a few species show dominance with changing percent frequencies during most of the sequence. The dominant benthic foraminiferal assemblages suggest that most of the Pleistocene bottom waters at site 219 and Early Pleistocene bottom waters at site 238 are of North Indian Deep Water (NIDW) origin. However, Late Pleistocene assemblage at site 238 appears to be closely associated with a water mass intermediate between North Indian Deep Water (NIDW) and Antarctic Bottom Water (AABW).

*Uvigerina proboscidea* is the most dominant benthic foraminiferal species present during the Pleistocene at both the sites. A marked increase in the relative abundance of *U. proboscidea* along with less diverse and equitable fauna during Early Pleistocene suggests a relative cooling, an intensified oceanic circulation and upwelling of nutrient rich bottom waters resulting in high surface productivity. At the same time, low sediment accumulation rate during Early Pleistocene reveals increased winnowing of the sediments possibly due to more corrosive and cold bottom waters. The Late Pleistocene in general, is marked by relatively warm and stable bottom waters as reflected by low abundance of *U. proboscidea* and more diverse and equitable benthic fauna.

The lower depth range for the occurrence of *Bulimina aculeata* in the Indian Ocean is around 2300 m, similar to that of many other areas. *B. aculeata* also shows marked increase in its abundance near the Pliocene/Pleistocene boundary while a sudden decrease in the relative abundance of *Stilostomella lepidula* occurs close to the Early/Late Pleistocene boundary.

**Keywords.** Pleistocene; benthic foraminifera; northern Indian Ocean; oceanography.

### 1. Introduction

The deep sea circulation influences the thermal structure of individual oceans, the distribution of deep sea biota and the state of preservation of deep sea sediments. It has now been established that the earth's climatic history is closely linked with the changes in the deep water circulation of the oceans (Weyl 1968; Newell 1974).

The study of benthic foraminifera is very useful in interpreting the changes in deep sea environment. They are the largest biomass present at the lower bathyal (> 1000 m) and abyssal depths in the modern oceans (Hessler and Jumars 1974) and are the dominant carbonate tests to be preserved in the deep sea sediments. Many benthic species have separate stratigraphic ranges, and their evolutionary and migratory patterns provide significant biostratigraphic and paleoecologic information about the deep sea environments (Douglas and Woodruff 1981; Kurihara and Kennett 1988).

Investigations on the benthic foraminiferal assemblages in the core tops show that different assemblages are associated with major bottom and deep water masses (Streeter 1973; Schnitker 1974; Lohmann 1978; Corliss 1979a, b; Ingle *et al* 1980; Douglas and Woodruff 1981; Kurihara and Kennett 1986; Sen Gupta 1988; Gupta and Srinivasan 1990a, b; Gupta 1992, 1994). These associations help in reconstructing the past distribution of the water masses and derive their physical and chemical characteristics.

In the present investigation DSDP sites 219 and 238 in the northern Indian Ocean were selected for quantitative benthic foraminiferal analysis to infer the Pleistocene bottom water oceanographic history of this region.

## 2. General setting of the area


The DSDP site 219 (Lat. 9°01'75"N; Long. 72°52'67"E) was drilled on the crest of the Chagos-Laccadive Ridge, a north-south trending structure extending southward from the eastern margin of the Arabian Sea at a water depth of 1764 m. Site 238 (Lat. 11°09'21"S; Long. 70°31'56"E) is situated east of the Central Indian Ridge near the southern end of the Chagos-Laccadive Ridge at a greater water depth of 2832 m. Locations of these two sites and prominent physiographic features of the Indian Ocean are shown in figure 1.

Antarctic Bottom Water (AABW) mostly confined to the deep basins (> 4,000 m) is the dominant bottom water mass in the Indian Ocean (Kolla *et al* 1976; Tchernia 1980). The Weddell Sea and the Ross Sea – Adelie coast are supposed to be two different sources for AABW in the Indian Ocean (Kolla *et al* 1976; Kolla and Kidd 1982). The Western Indian Ocean bottom waters are mainly of Weddell Sea origin whereas the bottom water from the Ross Sea is mostly confined to the Eastern Indian Ocean (Kolla and Kidd 1982). Kolla *et al* (1976) and Johnson and Damuth (1979) observed that the sea floor of the western and southern regions are more influenced by AABW than the northern regions of the Indian Ocean. Thus, at present the bottom waters at sites 219 and 238 are predominantly influenced by AABW although, the vigour of AABW appears to be somewhat subdued in this region compared to the southern part of the Indian Ocean. The bottom water potential temperatures in the Arabian basin and in the central Indian basin range from 1.00 to 1.20°C and 0.96 to 1.20°C respectively (Kolla *et al* 1976). The Calcium Carbonate Compensation Depth (CCD) in the Indian Ocean is at about 5,000 m (Kolla *et al* 1976). Both the examined sites lie above CCD and under Equatorial Water (EQW) and are ideal for calcium carbonate deposition.

The presence of uninterrupted Pliocene–Pleistocene sequences at both the sites enabled us to study the changes in the benthic foraminiferal distribution during the Pleistocene and also across the Pliocene/Pleistocene transition. The Pleistocene sections at sites 219 and 238 are dominated by foraminifera-rich nannofossil ooze.

## 3. Pliocene/Pleistocene boundary

The availability of a large number of Late Cenozoic deep sea cores of continuous sequences from tropical to polar regions as a result of Deep Sea Drilling Project



**Figure 1.** Location of DSDP sites (solid circles) examined and bottom water circulation in the Indian Ocean (modified after Kolla *et al.* 1976).

(DSDP) provided a unique opportunity in delineating the Pliocene/Pleistocene boundary from widely separated regions of the world. Detailed integrated micropaleontological and paleomagnetic studies of the deep sea cores proved useful in identifying various physical and bio-events and the duration of the Pleistocene. Based on recent studies the last appearance of *Globigerinoides fistulosus* which occurs at the top of Olduvai Normal Paleomagnetic Event at 1.6 Ma appears to be an excellent marker for delineating the Pliocene/Pleistocene boundary in the Tropical Indo-Pacific deep sea sequences (Srinivasan and Singh 1991; Srinivasan and Sinha 1991, 1992). Following the above criteria Pliocene/Pleistocene boundary is well defined at the examined DSDP sites 219 and 238.

#### 4. Method of study

The present investigation is based on a total of 19 samples from 33.0 m of core at site 219 and 22 samples from 34.5 m of core at site 238 at a regular interval of about 1.5 m. The samples (10 cc in volume) were treated with 15% hydrogen peroxide solution and water (1:3 ratio) and wet sieved over 63  $\mu\text{m}$  and 149  $\mu\text{m}$  tyler sieves. All the benthic foraminifera over 149  $\mu\text{m}$  size fraction (dry sieved) were picked from each sample. At both the sites benthic foraminifera are well preserved and commonly do not constitute more than 5% of the total foraminiferal population. The benthic foraminifera were mounted on microfaunal slides, identified and counted.

#### 5. Quantitative analyses of benthic foraminifera

Although much work has been carried out on generating qualitative benthic foraminiferal data, there are only few published records on the quantitative aspect of the Indian Ocean deep sea benthic foraminifera (Corliss 1979a, b, 1983, 1985; Gupta and Srinivasan 1990a, b, 1992a, b; Srinivasan and Gupta 1990; Gupta 1991, 1992, 1994; Rai and Srinivasan 1992, 1993). Most of the modern benthic foraminifera whose distribution pattern in the Indian Ocean is now well established also occur in the Pleistocene. Total of 137 and 107 benthic foraminiferal species were identified at sites 219 and 238 respectively (Appendix 1); of these only a few species dominate the assemblage and show changes in their relative abundances during the Pleistocene.

The relative percentages of dominant benthic foraminiferal groups and species (seven species at 219 and six at site 238) showing significant changes are plotted in figures 2-5. The Species Diversity,  $H(S)$  and Equitability,  $E'$  were also calculated using Shannon-Wiener Diversity Index (Rai and Srinivasan 1992) and are shown in figures 3 and 5.

The calcareous benthic foraminifera constitute the most dominant part of the Pleistocene assemblages at both the examined sites. Arenaceous foraminifera constitute about 5% of the total benthic foraminiferal population at both the sites whereas relative abundances of porcellaneous taxa range from 5 to 10% at site 219 and 5 to 15% at site 238 during the Pleistocene (figures 3 and 5). The dominant species at these two sites include *Uvigerina proboscidea*, *Cibicides wuellerstorfi*, *Globocassidulina subglobosa*, *Bulimina aculeata* and *Oridorsalis umbonatus*. Such a dominant assemblage has been recorded from the core top samples of the Indian




Figure 2. Percent distribution of important benthic foraminiferal species during the Pleistocene at the DSDP site 219. (*U. p. Uvigerina proboscidea*; *G. s. Globocassidulina subglobosa*; *B. a. Bulimina aculeata*; *S. l. Stilostomella lepidula*; *E. e. Epistomina exigua*; *O. u. Oridorsalis umbonatus*).





Figure 3. Percent distribution of important benthic foraminiferal groups and plots of Species Diversity, H(S); Equitability, E' and Number of Species, S during the Pleistocene at the DSDP site 219.



**Figure 4.** Percent distribution of important benthic foraminiferal species during the Pleistocene at the DSDP site 238. (*Bi. m. Biloculina murrina*; See figure 2 for rest of caption).




Figure 5. Percent distribution of important benthic foraminiferal groups and plots of Species Diversity,  $H(S)$ ; Equitability,  $E'$  and Number of Species,  $S$  during the Pleistocene at the DSDP site 238.




Figure 6. Distribution of *Bulimina aculeata* and *Stilostomella lepidula* during Pliocene-Pleistocene at seven DSDP sites in the Indian Ocean (Water Depth of each site is given in parenthesis) (Data of sites 214, 216 and 217 from Gupta 1987; and sites 220 and 237 from Rai 1992).

Ocean by Gupta (1994) in a water depth ranging 1500–2500 m, representing a high productivity region.

Most of the Pleistocene sequence at site 219 and Early Pleistocene at site 238 are dominated by *U. proboscidea* and *Globocassidulina subglobosa* assemblage (figures 2 and 4). Late Pleistocene in the latter site is marked by the abundance of benthic foraminiferal assemblage consisting of *C. wuellerstorfi*, *O. umbonatus*, *Epistominella exigua* and *Pullenia* spp. (mainly *P. bulloides*) (figures 4 and 5).

The most dominant species prevailing at both the sections is *U. proboscidea* which occurs in frequencies between 5 to 35% and shows remarkable changes through the Pleistocene. The earliest Pleistocene (c. 1.6 to c. 1.0 Ma) is marked by the highest relative abundance of *U. proboscidea* (> 35%) at both these sites (figures 2 and 4) which corresponds to the interval of less diverse and less equitable benthic fauna (figures 3 and 5). A distinct peak of abundance (> 25% at site 219 and > 15% at site 238) of *U. proboscidea* is also marked at about 0.4 Ma with corresponding low values of H(S) and E' (figures 2–5). *U. proboscidea* shows declining trend during the Early/Late Pleistocene transition at both these sites (figures 2 and 4). In general, Early Pleistocene is characterized by comparatively high and changing abundance of *U. proboscidea* than the Late Pleistocene at both the examined sites (figures 2 and 4).

At site 219, *Stilostomella lepidula* abundantly occurs during the Early Pleistocene whereas Late Pleistocene is marked by rare occurrence of this taxon (figures 2 and 6). *Bulimina aculeata* occurs abundantly (commonly > 5%) during the Pleistocene at site 219 and declines suddenly at the Pliocene/Pleistocene boundary (figures 5 and 6). At site 238 *B. aculeata* and *S. lepidula* show very rare and sporadic occurrences. The relative abundances of *O. umbonatus* and *U. proboscidea* show mutually opposite trends reflecting different ecological preferences (figures 2 and 4).

## 6. Discussion

In the two DSDP sites studied here the assemblages are characterized by several benthic foraminiferal species. However, only a few species in the assemblages show marked changes in their abundances through time. To interpret these variations in terms of oceanographic changes we need to understand their response to the deep water characteristics.

The characteristic North Indian Deep Water (NIDW) assemblage (*U. proboscidea* and *G. subglobosa*) dominates most of the Pleistocene sequence at site 219 and the Early Pleistocene interval at site 238. Whereas, the Late Pleistocene interval at site 238 is dominated by an assemblage consisting mainly of *C. wuellerstorfi*, *Oridorsalis umbonatus*, *Epistominella exigua*, *Pullenia* spp. alongwith occasional occurrence of *Nuttallides umbonifera*. Studies reveal that *Nuttallides umbonifera* is a cosmopolitan deep water species found associated with AABW (Streeter 1973; Schnitker 1974; Anderson 1975; Bremer and Lohmann 1982; Kurihara and Kennett 1986; Gupta 1992, 1994).

*Uvigerina* spp. (mainly *U. proboscidea*) is the most dominant element present during the Pleistocene and changes in its relative abundance appear to be closely linked with some climatically induced paleoceanographic changes (Gupta and Srinivasan 1992a). Recently, Boyle (1990) argued that factors unrelated to the deep water properties, such as surface productivity or the depositional environment may play a

significant role in controlling the distribution patterns of *Uvigerina* spp. This was supported by Gupta and Srinivasan (1992a), who pointed out that the peaks of abundance of *U. proboscidea* during the Late Neogene reflect intervals of increased upwelling leading to high surface productivity and perhaps high rates of biogenic sediment accumulation.

The Early Pleistocene interval (c. 1.6 to c. 1.0 Ma) at both the sites is marked by high relative abundance of *U. proboscidea* with corresponding low sediment accumulation. Gupta and Srinivasan (1990a, b) also recorded a similar change during the Early Pleistocene at site 214 (Ninetyeast Ridge). This reflects the high rate of surface productivity and increased winnowing of deep sea sediments due to dissolution by corrosive bottom waters during this interval. The Early Pleistocene is characterized by widespread deep sea hiatuses in global oceans (Ledbetter and Ciesielski 1986; Barker *et al* 1988). In general the presence of a hiatus reflects climatic cooling affecting the intensity of oceanic circulation (Kennett 1977; Barron and Keller 1982; Ciesielski *et al* 1982; Osborn *et al* 1983). The widespread deep sea hiatuses during the Early Pleistocene correspond to the times of enhanced Antarctic glaciation leading to low sea level stands during which an increase in AABW activity causes more erosion and dissolution of deep sea sediments (Moore *et al* 1978; Barron and Keller 1982). Keller and Barron (1987) suggested that increased polar cooling would cause high surface productivity due to upwelling and hence high rates of carbonate sedimentation. The low values of H(S) and E' during this interval also reflect the instability in the deep sea environments. The role of species diversity in the environmental studies have been discussed by several workers. The integrated study of the species diversity pattern and stable isotopic records in the Indian Ocean revealed that the lower values of H(S) and E' occurred during the intervals of higher  $\delta^{18}\text{O}$  and lower  $\delta^{13}\text{C}$  values reflecting global cooling and low sea level stands (Rai and Srinivasan 1992). The dominance of temperate planktonic foraminiferal assemblages in the northern Indian Ocean during the interval also reveals cooling of surface water due to climatic deterioration (Singh and Srinivasan 1993).

The low abundance of *U. proboscidea* and more diverse and equitable assemblage at both the sites during Late Pleistocene mark the onset of relative stability of the deep sea environments in the northern Indian Ocean. The intensity of deep and shallow hiatuses is low due to reduced velocity of AABW during the Late Pleistocene (Ledbetter and Ciesielski 1986). A short interval of high percent frequency of *U. proboscidea* at about 0.4 Ma represents the unstable deep sea environment of the northern Indian Ocean which corresponds to the high  $\delta^{18}\text{O}$  value in the North Atlantic (Shackleton and Hall 1984). This coherence in the faunal and isotopic data appears to be due to global climatic cooling leading to an increase in Northern and Southern Hemisphere ice sheets.

The quantitative benthic foraminiferal analyses at sites 219 and 238 as well as site 214 (Gupta and Srinivasan 1990a, b) reveal that in general, the Early Pleistocene was relatively cooler than the Late Pleistocene and marked by more intensified bottom water circulation. However, several short intervals of marked faunal changes represent the oscillations in the deep sea environments due to enhanced bottom water activity representing the intervals of frigid and unstable conditions. Because of the paucity of core samples at required sampling intervals it is not possible at this stage to identify various Pleistocene glacial and interglacial episodes.

The distribution pattern of *B. aculeata* during the Pleistocene at various DSDP

**Table 1.** Major faunal, isotopic and paleoceanographic events in the northern Indian Ocean and their relation with global climatic events (modified after Gupta and Srinivasan 1990a).

Age (Ma)	Epoch	Faunal events	Isotopic events	Paleoceanographic events	Global climatic events
		Changing abundance of cold and warm fauna		Cold/warm cycles	Major glacial/interglacial episodes
Late Pleistocene		Less diverse benthic assemblage increase in <i>U. proboscidea</i>	$\delta^{18}\text{O}$ increase	Distinct bottom water cooling	Increase in polar ice sheets.
0.6		Increase in benthic foraminiferal diversity and – low abundance of <i>U. proboscidea</i>		Stable and warm bottom water	Low intensity of shallow and deep sea hiatuses.
		Abrupt decline in <i>S. lepidula</i>			
	Early Pleistocene	Major increase in <i>U. proboscidea</i> abundance and lowest diversity of benthic foraminifera	$\delta^{18}\text{O}$ increase	Significant bottom water cooling, increased upwelling and high organic carbon flux	Enhanced glaciation, sea level low stands and widespread deep sea hiatuses
1.6		Sudden increase in <i>B. aculeata</i>	$\delta^{13}\text{C}$ decrease		

sites in the Indian Ocean reveals that it does not occur at water depths greater than 2300 m (figure 6). In other areas also the lower depth range for the occurrence of *B. aculeata* is around 2300 m (Kaiho and Nishimura 1992) suggesting that the distribution of this species is mainly controlled by the water depth. It is interesting to note that *B. aculeata* shows sudden increase in abundance close to Pliocene/Pleistocene boundary while abrupt decline in relative abundance of *S. lepidula* occurs near Early/Late Pleistocene boundary (c. 0.6 Ma).

## 7. Conclusions

A detailed quantitative study of the Pleistocene benthic foraminifera at the DSDP sites 219 and 238 as well as available data from site 214 (Gupta and Srinivasan 1990a, b) in the northern Indian Ocean has led to the following inferences (table 1).

1. The dominance of *U. proboscidea* and *G. subglobosa* assemblage during most of the Pleistocene at site 219 and beginning of the Pleistocene at site 238 suggests the influence of North Indian Deep Water (NIDW). However, during the later part of Pleistocene the nature of watermass at site 238 appears to be intermediate between NIDW and AABW as evidenced by the dominance of *C. wuellerstorfi*, *O. umbonatus*, *E. exigua* and *Pullenia* spp. assemblage along with occasional occurrences of *Nuttallides umboinifera*.
2. The peaks of abundance of *U. proboscidea* along with low values of H(S) and E' at both the sites during c. 1.6 to c. 1.0 Ma (Early Pleistocene) and at c. 0.4 Ma (Late Pleistocene) reflects marked cooling, intensified oceanic circulation and upwelling of the nutrient rich bottom waters giving rise to high surface productivity.
3. In general, the northern Indian Ocean bottom waters were relatively cooler during the Early Pleistocene than the Late Pleistocene as evidenced by the high abundance of *U. proboscidea* along with less diverse and equitable benthic fauna. The high dissolution due to cold and corrosive bottom waters during the Early Pleistocene possibly resulted into the low sediment accumulation.
4. The modern distribution pattern of *B. aculeata* in the Indian Ocean DSDP sites reveals that the lower depth range for this taxon is about 2300 m.

## Acknowledgements

The Deep Sea Drilling Project (DSDP) is thankfully acknowledged for providing the core samples. This research was supported by the grant of SAP (UGC) to the Department of Geology, Banaras Hindu University, Varanasi.

**Appendix 1.** Benthic foraminifera recorded from the examined sites.

Benthic foraminifera	Found in DSDP sites
<i>Textularia halkyardi</i> Lalicker	219, 238
<i>T. lythostrota</i> (Schwager)	219, 238
<i>T. porrecta</i> (Brady)	238
<i>Siphonotextularia rolshauseni</i> Phleger and Parker	219, 238
<i>S. solita</i> (Schwager)	219, 238
<i>Gaudryina solida</i> Schwager	219
<i>Dorothia brevis</i> Cushman and Stainforth	219
<i>Eggerella bradyi</i> (Cushman)	219, 238
<i>Karreriella baccata</i> (Schwager)	219
<i>K. bradyi</i> (Cushman)	219, 238
<i>K. subrotundata</i> (Schwager)	219, 238
<i>Martinottiella communis</i> (d'Orbigny)	219, 238
<i>Spiroloculina antillarum</i> d'Orbigny	219
<i>Quinqueloculina</i> cf. <i>pygmaea</i> Reuss	219, 238
<i>Q. seminulum</i> Linne'	219
<i>Q. venusta</i> Karrer	219, 238
<i>Q. weaveri</i> Rau	219, 238
<i>Pyrgo depressa</i> (d'Orbigny)	219, 238
<i>Biloculina comata</i> Brady	219
<i>B. lucernula</i> Schwager	219, 238
<i>B. murrhina</i> Schwager	219, 238
<i>Triloculina tricarinata</i> d'Orbigny	219, 238
<i>Nummoloculina irregularis</i> (d'Orbigny)	219
<i>Sigmoilopsis asperula</i> (Karrer)	219
<i>S. schlumbergeri</i> (Silvestri)	219, 238
<i>Nodosaria</i> aff. <i>lamnulifera</i> Boomgart	219
<i>N. raphanus</i> Schwager	219
<i>N. spirostriolata</i> Cushman	219
<i>Astacolus crepidulus</i> (Fichtel and Moll)	219, 238
<i>A. increscens</i> (Reuss)	219
<i>A. insolitus</i> (Schwager)	238
<i>A. reniformis</i> (d'Orbigny)	238
<i>Chrysagonium equisetiformis</i> (Schwager)	219, 238
<i>Dentalina costai</i> (Schwager)	219, 238
<i>D. intertenuata</i> (Schwager)	219
<i>D. intorta</i> (Dervieux)	238
<i>D. neugeboreni</i> (Schwager)	219, 238
<i>D. stimulea</i> (Schwager)	219, 238
<i>D. subsoluta</i> (Cushman)	219, 238
<i>Lagena</i> spp.	219, 238
<i>Robulus</i> aff. <i>gibbus</i> (d'Orbigny)	219, 238
<i>R. nicobarensis</i> (Schwager)	219
<i>Marginulina obesa</i> Cushman	219, 238

Appendix 1. *Continued*

Benthic foraminifera	Found in DSDP sites
<i>Orthomorphina</i> aff. <i>antillea</i> (Cushman)	238
<i>O. challengeriana</i> (Thalmann)	219
<i>O. koina</i> (Schwager)	219
<i>O. modesta</i> (Bermudez)	219
<i>O. perversa</i> (Schwager)	219, 238
<i>Guttulina austriaca</i> d'Orbigny	219
<i>Pyrulina cylindroides</i> (Roemer)	219, 238
<i>P. extensa</i> (Cushman)	219, 238
<i>P. fistulosa</i> (Thalmann)	219, 238
<i>P. labiata</i> Schwager	219, 238
<i>Fissurina</i> spp.	219, 238
<i>Oolina</i> spp.	219, 238
<i>Ceratobulimina pacifica</i> Cushman and Harris	219
<i>Hoeglundina elegans</i> (d'Orbigny)	219
<i>Buliminella andamanica</i> (Srinivasan and Singh)	219
<i>Sphaeroidina bulloides</i> d'Orbigny	219, 238
<i>Bolivinita pseudoplicata</i> (Heron-Allen and Earland)	219, 238
<i>Bolivina pseudopunctata</i> Hoeglund	219, 238
<i>Brizalina pusilla</i> (Schwager)	219
<i>Favocassidulina australis</i> Eade	219, 238
<i>F. decorata</i> (Sidebottom)	238
<i>F. favus</i> (Brady)	219, 238
<i>Stilostomella consobrina</i> (d'Orbigny)	219
<i>S. fistuca</i> (Schwager)	219
<i>S. insecta</i> (Schwager)	219, 238
<i>S. lepidula</i> (Schwager)	219, 238
<i>Bulimina aculeata</i> d'Orbigny	219, 238
<i>B. alazanensis</i> Cushman	219, 238
<i>B. jarvisi</i> Cushman and Parker	238
<i>B. striata</i> d'Orbigny	219
<i>B. subacuminata</i> Cushman and Stewart	219
<i>Protoglobobulimina affinis</i> (d'Orbigny)	238
<i>P. pacifica</i> Cushman	219
<i>Praeglobobulimina spinescens</i> (Brady)	219
<i>Uvigerina hispida</i> (Schwager)	219, 238
<i>U. cf. longa</i> (Cushman and Bermudez)	238
<i>U. peregrina</i> (Cushman)	219, 238
<i>U. porrecta</i> (Brady)	219
<i>U. proboscidea</i> (Schwager)	219
<i>U. schencki</i> (Asano)	219
<i>Noviuva interrupta-costata</i> (LeRoy)	219, 238
<i>Rectuvigerina royoi</i> Bermudez and Fuenmayor	219
<i>Discorbis subvilardeboanus</i> (Rhezak)	219, 238

**Appendix 1. Continued**

Benthic foraminifera	Found in DSDP sites
<i>Discopulvinulina bertheloti</i> (d'Orbigny)	219, 238
<i>Epistominella exigua</i> (Brady)	219, 238
<i>Gavelinopsis lobatulus</i> (Parr)	219
<i>Laticarinina pauperata</i> (Parker and Jones)	219, 238
<i>Planulina ariminensis</i> d'Orbigny	238
<i>Heronallenella boltovskoyi</i> Gupta and Sen Gupta	219, 238
<i>Nuttallides umbonifera</i> (Cushman)	219, 238
<i>Cibicides bengalensis</i> Srinivasan and Sharma	219, 238
<i>C. bradyi</i> (Trauth)	219, 238
<i>C. cicatricosus</i> (Schwager)	219
<i>C. kullenbergi</i> Parker	219, 238
<i>C. lucidus</i> (Reuss)	219
<i>C. robertsonianus</i> (Brady)	219, 238
<i>C. aff. soendaensis</i> Le Roy	219
<i>C. telisaensis</i> Le Roy	219, 238
<i>C. wuellerstorfi</i> (Schwager)	219, 238
<i>Pleurostomella acuminata</i> Cushman	238
<i>P. alternans</i> Schwager	219, 238
<i>P. bolivinoides</i> Schubert	238
<i>P. brevis</i> Schwager	219, 238
<i>P. obtusa</i> Berthelin	219, 238
<i>P. sapperi</i> Schubert	219
<i>Ellipsoidella</i> sp.	219
<i>Fursenkoina bradyi</i> (Cushman)	219
<i>F. bramletti</i> (Galloway and Morrey)	219
<i>F. schreibersiana</i> (Czjzek)	238
<i>F. texturata</i> (Brady)	219, 238
<i>Francesita advena</i> (Cushman)	219, 238
<i>Cassidulina carinata</i> (Silvestri)	219, 238
<i>C. laevigata</i> d'Orbigny	219, 238
<i>Ehrenbergina carinata</i> Eade	219, 238
<i>E. praebicornis</i> n. sp.	219, 238
<i>Evolovocassidulina bradyi</i> Norman	219, 238
<i>E. seranensis</i> Germeraad	219
<i>Globocassidulina elegans</i> (Sidebottom)	219, 238
<i>G. oblonga</i> (Reuss)	219, 238
<i>G. murrhina</i> (Schwager)	219
<i>G. subglobosa</i> (Brady)	219, 238
<i>G. tumida</i> (Heron-Allen and Earland)	219, 238
<i>Chilostomella oolina</i> Schwager	219, 238
<i>Allomorphina pacifica</i> Hofker	219, 238
<i>Quadrrimorphina laevigata</i> (Phleger and Parker)	219, 238
<i>Astrononion stelligerum</i> (d'Orbigny)	219, 238

Appendix 1. *Continued*

Benthic foraminifera	Found in DSDP sites
<i>A. umbilicatum</i> Uchio	219, 238
<i>Nonionella auris</i> (d'Orbigny)	219, 238
<i>N. clavata</i> Cushman	219
<i>N. japonicum</i> (Asano)	219, 238
<i>Pullenia bulloides</i> (d'Orbigny)	219, 238
<i>P. osloensis</i> Feyling-Hanssen	219, 238
<i>P. quadriloba</i> Reuss	219, 238
<i>P. quinqueloba</i> (Reuss)	219, 238
<i>P. salisburyi</i> Stewart and Stewart	219, 238
<i>Oridorsalis prominula</i> (Stache)	219, 238
<i>O. umbonatus</i> (Reuss)	219, 238
<i>Osangularia culter</i> (Parker and Jones)	219, 238
<i>Gyroidinoides broeckhianus</i> (Karrer)	219, 238
<i>G. cibaoensis</i> (Bermudez)	219, 238
<i>G. nitidula</i> (Schwager)	219, 238
<i>G. polius</i> (Phleger and Parker)	219, 238
<i>Anomalina globulosa</i> Chapman and Parr	219, 238
<i>Anomalinoides glabratus</i> (Cushman)	219
<i>Melonis barleeanum</i> (Williamson)	219, 238
<i>M. nicobarensis</i> (Cushman)	219, 238
<i>M. pompilioides</i> (Fichtel and Moll)	219, 238

## References

Anderson J B 1975 Ecology and distribution of foraminifera in the Weddell Sea of Antarctica; *Micro-paleontology* **21** 69–96

Barker P F, Kennett J P and Scientific party 1988 Weddell Sea Paleoceanography: Preliminary results of ODP Leg 113; *Palaeogeogr., Palaeoclimatol., Palaeoecol.* **67** 75–102

Barron J A and Keller G 1982 Widespread Miocene deep-sea hiatuses: coincidence with periods of global cooling; *Geology* **10** 577–581

Boyle E A 1990 Quaternary deep water paleoceanography; *Science* **249** 863–873

Bremer M L and Lohmann G P 1982 Evidence for primary control of the distribution of certain Atlantic Ocean benthonic foraminifera by degree of carbonate saturation; *Deep Sea Res.* **29** 987–989

Ciesielski P F, Ledbetter M T and Ellwood B B 1982 The development of Antarctic glaciation and the Neogene paleoenvironment of the Maurice Ewing Bank; *Mar. Geol.* **46** 1–51

Corliss B H 1979a Recent deep-sea benthic foraminiferal distribution in the southeast Indian Ocean: inferred bottom-water routes and ecological implication; *Mar. Geol.* **31** 115–138

Corliss B H 1979b Quaternary Antarctic bottom water history: Deep sea benthonic foraminiferal evidence from the southeast Indian Ocean; *Quat. Res.* **12** 271–289

Corliss B H 1983 Distribution of Holocene deep-sea benthonic foraminifera in the southwest Indian Ocean; *Deep Sea Res.* **30** 95–117

Corliss B H 1985 Microhabitats of benthic foraminifera within deep sea sediments; *Nature (London)* **314** 435–438

Douglas R C and Woodruff F 1981 Deep-sea benthic foraminifera. The Oceanic Lithosphere, The Sea. In: (ed.) C Emiliani (New York: John Wiley) 7 pp. 1233–1327

Gupta A K 1987 Neogene deep water benthic foraminifera from the Northern Indian Ocean (DSDP Leg 22). Ph. D. Thesis, Banaras Hindu University, Varanasi (Unpublished)

Gupta A K 1991 Pleistocene benthic foraminifera and deep sea environment at Northern Indian Ocean DSDP Site 219; *J. Geol. Soc. India* **37** 388–406

Gupta A K 1992 Benthic foraminifera and deep-sea water masses in the tropical Indian Ocean; *J. Geol. Soc. India* **39** 475–486

Gupta A K 1994 Holocene deep-sea benthic foraminifera and water masses in the Indian Ocean and Red Sea; *J. Geol. Soc. India* **43** 691–703

Gupta A K and Srinivasan M S 1990a Response of northern Indian Ocean deep-sea benthic foraminiferal to global climates during Pliocene-Pleistocene; *Mar. Micropaleontology* **16** 77–91

Gupta A K and Srinivasan M S 1990b Quaternary bottom water circulation and benthic foraminiferal changes in the northern Indian Ocean; *Indian J. Geol.* **62** 102–116

Gupta A K and Srinivasan M S 1992a *Uvigerina proboscidea* abundances and paleoceanography of the Northern Indian Ocean DSDP Site 214 during the Late Neogene; *Mar. Micropaleontology* **19** 355–367

Gupta A K and Srinivasan M S 1992b Deep-sea benthic foraminiferal changes and terminal Miocene event at tropical Indian Ocean Site 214; *J. Geol. Soc. India* **40** 262–278

Hessler R P and Jumars P 1974 Abyssal community analysis from replicate box cores in the central north Pacific; *Deep Sea Res.* **21** 185–209

Ingle J C Jr, Keller G and Kolpack R L 1980 Benthic foraminiferal biosfacies, sediments and water masses of the Southern Peru-Chile Trench area, Southeastern Pacific Ocean; *Micropaleontology* **26** 113–150

Johnson D A and Damuth J E 1979 Deep thermohaline flow and current-controlled sedimentation in the Amirante Passage: Western Indian Ocean; *Mar. Geol.* **33** 1–44

Kaiho K and Nishimura A 1992 Distribution of Holocene benthic foraminifera in the Izu-Bonin Arc; *Proc. ODP, Scientific Results* **126** 311–320

Keller G and Barron J A 1987 Paleodepth distribution of Neogene deep-sea hiatuses; *Paleoceanography* **2** 697–713

Kennett J P 1977 Cenozoic evolution of Antarctic glaciation, the circum-Antarctic Ocean and their impact on global paleoceanography; *J. Geophys. Res.* **82** 3843–3860

Kolla V and Kidd R B 1982 Sedimentation and sedimentary processes in the Indian Ocean. In: *The ocean basins and margins*. (eds.) A E Nairn and F G Stehli (New York: Plenum Press) **6** 1–50

Kolla V, Sullivan L, Streeter S S and Langseth M G 1976 Spreading of Antarctic Bottom Water and its effect on the floor of the Indian Ocean inferred from bottom-water potential temperature, turbidity and sea-floor photography; *Mar. Geol.* **21** 171–189

Kurihara K and Kennett J P 1986 Neogene benthic foraminifers: Distribution in depth traverse, Southwest Pacific. In: *Initial Reports of the Deep Sea Drilling Project* (ed.) J P Kennett and C C Von der Borch *et al* (Washington, DC: US Government Printing Office) **90** 1037–1077

Kurihara K and Kennett J P 1988 Bathymetric migration of deep-sea benthic foraminifera in the southwest Pacific during the Neogene; *J. Foraminifera Res.* **18** 75–83

Ledbetter M T and Ciesielski P F 1986 Post Miocene disconformities and paleoceanography in the Atlantic sector of the Southern Ocean; *Palaeogeogr. Palaeoclimatol. Palaeoecol.* **52** 185–214

Lohmann G P 1978 Abyssal benthonic foraminifera as hydrographic indicators in the western South Atlantic Ocean; *J. Foraminiferal Res.* **8** 6–34

Moore T C Jr, Van Andel T H, Sancetta C and Pisias N G 1978 Cenozoic hiatuses in marine sediments; *Micropaleontology* **24** 113–138

Newell R E 1974 Change in the poleward energy flux by the atmosphere and ocean as a possible cause for ice ages; *Quat. Res.* **4** 117–127

Osborn N I, Ciesielski P F and Ledbetter M T 1983 Disconformities and paleoceanography in the southeast Indian Ocean during the past 5.4 million years; *Geol. Soc. Am. Bull.* **94** 1345–1358

Rai A K 1992 Neogene deep sea benthic foraminifera of DSDP sites 219, 220 (Arabian Sea) and 237 and 238 (Central Indian Ocean); Ph. D. Thesis, Banaras Hindu University, Varanasi (Unpublished)

Rai A K and Srinivasan M S 1992 Neogene deep sea benthic foraminiferal diversity in the Indian Ocean: Paleoceanographic implications; *Proc. Indian Acad. Sci. (Earth Planet. Sci.)* **101** 299–316

Rai A K and Srinivasan M S 1993 Benthic foraminiferal changes across the Miocene/Pliocene boundary in the Northern Indian Ocean; Paleoceanographic implications; *Proc. Indian Natn. Sci. Acad.* **A59** 507–518

Schnitker D 1974 West Atlantic abyssal circulation during the past 120,000 years; *Nature (London)* **248** 385–387

Sen Gupta B K 1988 Water mass relation of the benthic foraminifer *Cibicides wuellerstorfi* in the Eastern Caribbean Sea; *Bulletin de l'Institut de Géologie du Bassin d'Aquitaine, Bordeaux* **44** 23-32

Shackleton N J and Hall M A 1984 Oxygen and carbon isotope stratigraphy of Deep Sea Drilling Project Hole 552a: Plio-Pleistocene glacial history. In: *Initial Reports of Deep Sea Drilling Project* (eds) D G Roberts, *et al* US Govt. Printing Office pp. 599-609

Singh A D and Srinivasan M S 1993 Quaternary climatic changes indicated by planktonic foraminifera of northern Indian Ocean; *Curr. Sci.* **64** 908-915

Srinivasan M S and Gupta A K 1990 Miocene benthic foraminifera and paleoceanography of DSDP sites along Ninetyeast Ridge, northern Indian Ocean; *J. Paleont. Soc. India* **35** 61-72

Srinivasan M S and Singh A D 1991 The Pliocene/Pleistocene boundary in the tropical Indian Ocean deep sea cores; *J. Sci. Res. BHU*, **B41** 199-222

Srinivasan M S and Sinha D K 1991 Improved correlation of the Late Neogene planktonic foraminiferal datums in the Equatorial to cool subtropical DSDP sites, Southwest Pacific: Application of the graphic correlation method; *Geol. Soc. India Mem.* **20** 55-93

Srinivasan M S and Sinha D K 1992 Late Neogene planktonic foraminiferal events of the Southwest Pacific and Indian Ocean: A comparison. In: *Pacific Neogene: Environment evolution and events*. (eds.) R Tsuchi and J C Ingle Jr (Japan University of Tokyo Press) pp. 203-220

Streeter S S 1973 Bottom water and benthonic foraminifera in the North Atlantic-glacial-interglacial contrasts; *Quat. Res.* **3** 131-141

Tchernia P 1980 *Descriptive Regional Oceanography*. (New York: Pergamon Press) pp. 1-253

Weyl P K 1968 The role of the oceans in climatic change: A theory of the ice ages; *Meteorological Monographs* **8** 37-62