

Exact ground and excited states of a t-J ladder doped with two holes

Indrani Bose and Saurabh Gayen*

Department of Physics

Bose Institute

93/1, A.P.C.Road

Calcutta-700 009, India.

* Physics Department

Scottish Church College

Calcutta-700006

Abstract

A two chain ladder model is considered described by the strong coupling $t - t' - J - J'$ Hamiltonian. For the case of two holes moving in a background of antiferromagnetically interacting spins, exact, analytical results are derived for the ground state energy and low-lying excitation spectrum. The ground state is a bound state of two holes with total spin $S=0$. The charge excitation is gapless and the spin excitation has a gap. The corresponding wavefunctions are also exactly determined. The bound hole pair is found to have symmetry of the d-wave type. In the limit of strong rung coupling, the model maps onto an effective hard core boson model which exhibits dominant superconducting pairing correlations.

I. Introduction

In the last few years, ladder systems have been studied extensively [1, 2, 3]. Interacting electron systems in one dimension (1d) are more or less well understood. There are several rigorous results available for such systems [3]. Powerful techniques like the Bethe Ansatz [4] and bosonization [5] have yielded much useful information about such systems. After the discovery of high- T_c cuprate superconductivity, 2d interacting electron systems acquired new significance due to the fact that the dominant electronic and magnetic properties of the cuprate systems are associated with the CuO_2 plane [6, 7].

There are, however, very few rigorous results available for 2d systems. Ladders, consisting of n-chains coupled by rungs, interpolate between 1d and 2d and their study is expected to be useful for a proper understanding of interacting many body systems. The possibility of deriving rigorous results is also more. A number of ladder systems have been discovered recently exhibiting a variety of interesting phenomena[1, 2, 3]. Physical insight obtained from the study of ladders is also expected to be relevant for high- T_c cuprate systems. The cuprates, in the spin-disordered phase, are doped spin liquids. Below optimal doping levels and well above the superconducting transition temperature T_c , there are experimental signatures of a spin gap(SG)[7] opening up. The 'gap' has been ascribed to pre-formed Cooper pairs of holes which lack the long-range phase coherence of the superconducting state. The Cooper pairs become phase coherent only below T_c giving rise to superconductivity. Dagotto et al[8] were the first to show that a two-chain ladder has a spin liquid ground state and a SG in the excitation spectrum. On doping the system with holes, binding of holes in pairs is possible, giving rise to dominant superconducting(SC) pairing correlations. A few years later, a hole doped two-chain ladder system $Sr_{0.4}Ca_{13.6}Cu_{24}O_{41.84}$ was discovered which exhibits superconductivity under pressure[9].

The relationship between 'pseudo' spin-gap, pre-formed hole pairs and superconductivity is not well-understood in the case of cuprate systems. For the ladder system, the SG is a real gap and the binding of holes leading to SC pairing correlations can be explicitly demonstrated. Resistivity measurements of the ladder compound $(Sr, Ca)_{14}Cu_{24}O_{41}$ show unusual temperature dependence as in the case of cuprates [10]highlighting further similarities between the two systems. Bose and Gayen [11, 12, 13, 14]have constructed a two-chain t-J type ladder model for which several exact, analytical results can be derived in the undoped as well as doped cases. For two holes, the possibility of binding of holes was suggested but the bound state spectrum was not derived. In Section II of this paper, we give a detailed derivation of the low-lying spin and charge excitation spectrum of the ladder model in the two-hole sector. We show that the ground state consists of a bound pair of holes. The spin excitation spectrum has a gap and the charge excitation is gapless. The two-hole wave functions are also computed. The two-hole bound state wave function is shown to have modified d-wave symmetry. All these results are exact and analytic in nature. The dominance of SC pairing correlations in the ladder model is shown in an approximate, analytical

manner.

II. Exact two-hole excitation spectrum

The two-chain ladder model consists of two chains, each described by a t - J Hamiltonian, coupled by t' - J' interactions between them (Fig.1). The model is described by the $t - t' - J - J'$ Hamiltonian:

$$\begin{aligned}
 H &= - \sum_{i,j,\sigma} t_{ij} (1 - n_{i-\sigma}) C_{i,\sigma}^\dagger C_{j,\sigma} (1 - n_{j-\sigma}) + H.C. + \sum_{\langle ij \rangle} J_{ij} \vec{S}_i \cdot \vec{S}_j \\
 &= H_t + H_{t'} + H_J + H_{J'}
 \end{aligned} \tag{1}$$

The constraint that no site can be doubly occupied is implied in the model. The hopping integral t_{ij} has the value t for nearest-neighbour(NN) hopping within a chain and also for diagonal transfer between chains (solid lines in Fig.1). The corresponding spin-spin interactions J_{ij} are of strength J . The spins have magnitude $1/2$. The hopping integral across vertical links (broken lines) connecting two chains has the strength t' . The corresponding spin-spin interaction strength J_{ij} is J' . We assume t and t' to be positive. In the conventional two-chain spin-ladder, the diagonal interaction and hopping terms are absent. The inclusion of the diagonal terms of the same strength as the intra-chain ones enables one to reduce the difficult N-body problem to an easily solvable few body problem. The conventional spin ladder model, in the absence of diagonal terms, constitutes a many-body problem for which no simplification occurs. The only exact results, which are available, are numerical results based on exact diagonalization of finite ladders [1, 2, 15].

In the half-filled limit, i.e, in the absence of holes, the $t - t' - J - J'$ -Hamiltonian in (1) reduces to $H_J + H_{J'}$. The exact ground state ψ_g (for $J' \geq 2J$) consists of singlets along the rungs of the ladder [11]. The ground state energy $E_g = -(3J'/4)N$, where N is the number of rungs. An exact excited state can be constructed by replacing a singlet by a triplet. Creation of a triplet costs an amount of energy $J'/4$ so that the spin gap $\Delta_{SG} = J'$. The excitation is localised and has no dynamics. Let us now consider the case of a single hole doped into the ladder. In the presence of holes a single rung can exist in nine possible states: (i) empty, (ii) two bonding states, (iii) two anti-bonding states, (iv) one singlet state and (v) three triplet states.

These states are shown below.

$$\begin{aligned}
(i) & \begin{pmatrix} o \\ o \end{pmatrix} & (ii) & \frac{1}{\sqrt{2}} \begin{pmatrix} \uparrow o \\ o \uparrow \end{pmatrix}, \quad \frac{1}{\sqrt{2}} \begin{pmatrix} \downarrow o \\ o \downarrow \end{pmatrix} \\
(iii) & \frac{1}{\sqrt{2}} \begin{pmatrix} \downarrow o \\ o \downarrow \end{pmatrix}, \quad \frac{1}{\sqrt{2}} \begin{pmatrix} \uparrow o \\ o \uparrow \end{pmatrix} \\
(iv) & \frac{1}{\sqrt{2}} \begin{pmatrix} \uparrow \\ \downarrow \end{pmatrix} - \frac{1}{\sqrt{2}} \begin{pmatrix} \downarrow \\ \uparrow \end{pmatrix} & (v) & \uparrow, \downarrow, \quad \frac{1}{\sqrt{2}} \begin{pmatrix} \uparrow \\ \downarrow \end{pmatrix} + \frac{1}{\sqrt{2}} \begin{pmatrix} \downarrow \\ \uparrow \end{pmatrix}
\end{aligned}$$

A single hole hops in a background of antiferromagnetically interacting spins. This, in general, is a difficult many body problem because as the hole hops it gives rise to spin excitations in the system. The inclusion of diagonal hopping terms in our model leads to a cancellation of all the terms containing spin excitations, resulting in a perfect, coherent motion of the hole. We illustrate this through an explicit example. Consider a single hole in a bonding state, located in the m -th rung. All other rungs are in singlet spin configurations. A pictorial representation of the state is

$$| \dots \frac{1}{\sqrt{2}} \begin{pmatrix} \uparrow o \\ o \uparrow \end{pmatrix}_m |_{(m+1)} \dots \quad (2)$$

$$\frac{1}{\sqrt{2}} \begin{pmatrix} \uparrow o \\ o \uparrow \end{pmatrix}_m |_{(m+1)} \equiv \frac{1}{2} \left(\begin{pmatrix} \uparrow \uparrow \\ o \downarrow \end{pmatrix} - \begin{pmatrix} \uparrow \downarrow \\ o \uparrow \end{pmatrix} + \begin{pmatrix} o \uparrow \\ \uparrow \downarrow \end{pmatrix} - \begin{pmatrix} o \downarrow \\ \uparrow \uparrow \end{pmatrix} \right) \quad (3)$$

The state is an exact eigenstate of the J, J', t' part of the $t - t' - J - J'$ Hamiltonian. Let us now apply H_t on the state. Since background electrons are fermions, their ordering is important and one has to keep track of signs during interchanges. The ordering of fermions follows the convention

$$\begin{array}{cccc}
1 & 3 & 5 & \cdot \cdot \\
2 & 4 & 6 & \cdot \cdot
\end{array}$$

On operating with H_t on the state shown in (3), one gets

$$\begin{aligned}
H_t \begin{pmatrix} \uparrow \uparrow \\ o \downarrow \end{pmatrix} &= t \begin{pmatrix} \uparrow \uparrow \\ \downarrow o \end{pmatrix} - t \begin{pmatrix} \uparrow o \\ \uparrow \downarrow \end{pmatrix} \\
-H_t \begin{pmatrix} \uparrow \downarrow \\ o \uparrow \end{pmatrix} &= -t \begin{pmatrix} \uparrow \downarrow \\ \uparrow o \end{pmatrix} + t \begin{pmatrix} \uparrow o \\ \downarrow \uparrow \end{pmatrix} \\
H_t \begin{pmatrix} o \uparrow \\ \uparrow \downarrow \end{pmatrix} &= t \begin{pmatrix} \uparrow o \\ \uparrow \downarrow \end{pmatrix} - t \begin{pmatrix} \downarrow \uparrow \\ o \end{pmatrix} \\
-H_t \begin{pmatrix} o \downarrow \\ \uparrow \uparrow \end{pmatrix} &= -t \begin{pmatrix} \downarrow o \\ \uparrow \uparrow \end{pmatrix} + t \begin{pmatrix} \uparrow \downarrow \\ \uparrow o \end{pmatrix}
\end{aligned}$$

The states in the second column are obtained due to diagonal hopping of the hole. There is a cancellation of the terms containing parallel spin pairs and the final state is given by

$$|_m \frac{1}{\sqrt{2}} (\uparrow_o + \uparrow_{\uparrow})_{m+1} \quad (4)$$

One finds that the hole accompanied by a free spin-1/2 moves coherently by one lattice unit(compare with Eqn.(3)). The eigenvalue problem now becomes very easy to solve. Let

$$\Psi(m) = ||| \cdots \frac{1}{\sqrt{2}} (\uparrow_o + \uparrow_{\uparrow})_m |_{(m+1)} \cdots \quad (5)$$

$$\Psi = \frac{1}{\sqrt{N}} \sum_{m=1}^N e^{ikm} \Psi(m) \quad (6)$$

Ψ is an exact eigenstate of the $t - t' - J - J'$ Hamiltonian with the energy eigenvalue

$$E_1 = 2t \cos(k) - t' + 3J'/4 \quad (7)$$

The energy is measured with respect to that of the ground state energy in the undoped state. Refs.[11, 12] give a detailed discussion of the single hole spectrum for both bonding and antibonding hole states. For conventional spin ladders, Troyer et al [15] have found numerical evidence of quasi-particle (QP) excitations carrying charge +e and spin-1/2. The charge and spin may be located on different rungs. In the exact eigenstate of Eqn.(6), the positively charged hole and the spin-1/2 are always located on the same rung. We refer to the composite object as hole -QP.

Let us now consider the case of two holes. The two holes can be introduced on the same rung or on separate rungs. Other rungs are in the singlet spin configurations. If the holes are located on two separate rungs, there are two free spins which can combine to give either a triplet or a singlet. In the triplet sector, the two hole-QPs can scatter against each other giving rise to a continuum of scattering states with energy

$$E_{cont} = 4t \cos(K/2) \cos q - 2t' + 3J'/2 \quad (8)$$

$K(= k_1 + k_2)$ and $q(= (k_1 - k_2)/2)$ are the centre of mass momentum and the relative momentum wave vectors. The two-hole ground state belongs to

the singlet sector. The exact eigenvalue equations have already been derived in Ref.[13] but a full analysis of these equations has so far not been done. Define the wave functions

$$\begin{aligned} \phi(m_1, m_2) = & \frac{1}{2\sqrt{2}} [|\cdots | (\uparrow + \downarrow)_{m_1} | \cdots (\uparrow + \downarrow)_{m_2} | \cdots | \\ & - |\cdots | (\uparrow + \downarrow) | \cdots (\uparrow + \downarrow) | \cdots |] \end{aligned} \quad (9)$$

and

$$\phi(m, m) = || \cdots \downarrow_m \cdots ||. \quad (10)$$

Define also the Fourier transforms

$$\phi(m, m+r) = \frac{1}{\sqrt{N}} \sum_K \exp[iK(m+r/2)] \phi_K(r) \quad (11)$$

for $0 \leq r \leq N/2 - 1$ and

$$\phi(m, m+N/2) = \sqrt{\frac{2}{N}} \sum_K \exp[iK(m+N/2)] \phi_K(N/2) \quad (12)$$

The two holes are separated by a distance r . From the periodic boundary condition and for $r \neq N/2$, the allowed values of K are $K = (2\pi/N)\lambda$, with $\lambda = 0, 1, 2, \dots, N-1$. For $r=N/2$, the allowed values of K are odd multiples of $2\pi/N$. An eigenfunction in the momentum space is given by

$$\Psi_e^K = \sum_{r=0}^{N/2-1} a(r) \phi_K^r \quad (13)$$

where K is an even multiple of $2\pi/N$. When K is an odd multiple of $2\pi/N$, the eigenfunction is Ψ_0^K and the sum in Eq.(13) runs from 0 to $N/2$. The exact eigenvalue equations for both the cases are given in Ref.[13]. When K is an even multiple of $2\pi/N$, the amplitudes $a(r)$ have the form

$$a(r) = \sin[q(N/2 - r)] \text{ for } 1 \leq r \leq N/2 - 1 \quad (14)$$

The energy eigenvalues are obtained by simultaneously solving the equations

$$\epsilon = 2T \cos q \quad (15)$$

$$\epsilon + \frac{3J}{4} = \frac{4T^2}{\epsilon + \frac{3J'}{4} - 2t'} + \frac{T \sin[q(N/2 - 2)]}{\sin[q(N/2 - 1)]} \quad (16)$$

where $\epsilon = E - 3J'/2 + 2t'$ and, as before, energy E is measured w.r.t. that of the ground state in the undoped case. The energies for real values of q correspond to free hole states. Energies for bound and antibound states are obtained by making q complex. When T is +ve, making the changes $q \rightarrow iq$ and $q \rightarrow \pi + iq$, one gets the energies for antibound and bound states, respectively. When T is negative, the reverse is true. Similar results are obtained when K is an odd multiple of $2\pi/N$.

We now study the eigenvalue problem in the limit $N \rightarrow \infty$. The continuum of hole excited states, for real q , is given by Eqn.(15). For complex q bound and antibound states are obtained. Let us now replace q by $\pi + iq$ in Eqns.(15,16). Since N is large, Eqn.(16) reduces to

$$\epsilon + \frac{3J}{4} = \frac{4T^2}{\epsilon + \frac{3J'}{4} - 2t'} - T e^{-q} \quad (17)$$

From a simultaneous solution of Eqn.(15) (with q replaced by $\pi + iq$) and Eqn.(17), one gets the following cubic equation in e^q ,

$$e^{3q} - e^{2q} \left[\frac{3J + 3J'}{4T} - \frac{2t'}{T} \right] + e^q \left[\frac{3J}{4T^2} (-2t' + 3J'/4) - 3 \right] - \left(\frac{3J}{4T} \right) = 0 \quad (18)$$

The exact, analytic solutions of a cubic equation are given in Ref.[16]. For a physical solution, e^q is greater than or equal to 1. There are at most two physical solutions of the cubic equation in (18). Once a solution for e^q is obtained, the energy eigenvalue is obtained from Eqn.(15) (with q replaced by $\pi + iq$). For positive values of T , one gets the solution for a bound state of two holes and for T -ve, a solution for the antibound state is obtained. The other values of the excitation branches are obtained by symmetry. Fig.2 shows the exact energy spectrum for the bound state, continuum of scattering states and antibound states of two holes for $J = t = t' = 1$ and $J' = 2J$. Fig.3 shows the same for the parameter values $J/t = 0.25$, $t = t' = 1$ and $J' = 2J$. The bound state of holes is obtained irrespective of the value of J/t being less than or greater than 1. Dagotto et al [8] were the first to show the binding of two holes in a two-chain ladder system. Their finding was based on exact diagonalization of finite-sized ladder systems. Later, Troyer et al [15] also found evidence for the binding of holes in finite ladder systems. In

the case of our model, we have shown exactly and analytically the binding of two holes for $N \rightarrow \infty$. For finite systems also, one can solve the eigenvalue problem exactly.

The two-hole ground state is the bound state of two holes with centre of mass momentum wave vector $K=0$. The exact bound state wave function is given by (13) with $K=0$ and q replaced by $\pi + iq$ in (14). In the limit $N \rightarrow \infty$, one obtains

$$\frac{a(n)}{a(0)} = (-1)^{(n-1)} e^{-(n-1)q} \frac{a(1)}{a(0)} \quad (19)$$

This result shows explicitly that the bound state wave function has an exponential decay as the separation between the two holes increases. With the knowledge of the eigenvalue ϵ , the ratio $\frac{a(1)}{a(0)}$ can be computed from the exact eigenvalue equations derived in Ref.[13]. Fig.4 shows a plot of $|\frac{a(r)}{a(0)}|^2$ versus r for the ground state wave function with parameter values $J = t = t' = 1.0$ and $J' = 2J$ (dotted line), $J' = 10J$ (solid line). When J' is much larger than J , the holes prefer to be on the same rung to minimise the loss in exchange interaction energy. The hole delocalization energy along the rung is, however, lost. When J' and J are comparable, $|\frac{a(r)}{a(0)}|^2$ has maximum value when holes are separated by approximately one lattice constant. The exchange energy loss is less when two holes are on NN rungs than when they are further apart. Being on separate rungs, the holes gain in delocalization energy. The bound state is also more extended. These results are in agreement with the numerical results of Troyer et al [15].

The low energy modes of a ladder system are characterised by their spin. Singlet and triplet excitations correspond to charge and spin modes respectively. The two hole ground state is in the singlet sector and, as already discussed, corresponds to a bound state of two holes for $K=0$. Since a hole bound state branch exists in the singlet sector, excitations with energy infinitesimally close to the ground state energy are possible. These excitations are the charge excitations since the total spin is still zero and the charge excitation spectrum is gapless.

There are two distinct types of spin excitations. The first is the magnon ($S=1$) excitation of the undoped ladder with energy J' measured with respect to the ground state energy. The spin triplet excitation appears on doping the ladder. For a pair of holes, the lowest triplet excitation energy is $-4t -$

$2t' + 3J'/2$ from Eqn.(8). The lowest triplet excitation energy depends on the values of t , t' and J' . The spin gap energy Δ_{SG} is the difference in energies of the lowest triplet excitation and the ground state (two hole bound state in the singlet sector) energy. Fig.5 shows Δ_{SG} versus J/t for $t = t' = 1.0$ and $J' = 2J$. Thus, the two-chain ladder model has the feature that the charge excitation is gapless but the spin excitation has a gap. The same result holds true for the conventional spin ladder [2, 15]. In the notation CxSy [17](x gapless charge and y gapless spin excitations), the t-J type ladder model exists in the C1S0(Luther-Emery) phase.

The experimental evidence of hole based superconductivity [9] in a ladder system provides the motivation to look for superconducting pairing correlations in our ladder model. We have already shown the existence of the two-hole bound state. Define the pairing operator

$$\Delta_{ij} = c_{i\downarrow} c_{j\uparrow} - c_{i\uparrow} c_{j\downarrow} \quad (20)$$

and consider the quantity

$$\tilde{\Delta}_{ij} = \langle 2|\Delta_{ij}|0\rangle \quad (21)$$

$|0\rangle$ and $|2\rangle$ are the ground states in the case of zero and two holes respectively. For our ladder model, both of those ground states are exactly known and one can verify that $\tilde{\Delta}_{i i+\hat{x}}$ and $\tilde{\Delta}_{i i+\hat{y}}$ have opposite signs, \hat{x} and \hat{y} denote unit vectors in the x(along chain) and y(along rung) directions. This is a signature of d-wave pairing and shows that the bound state of two holes has symmetry of the d-wave type. In the case of cuprate superconductors, there is much experimental evidence that the pairing wave function has d-wave symmetry [18].

In the large J' limit, the ladder model can be mapped onto an effective boson model [15]. The physical picture is that of bound hole pairs existing along rungs and moving in a background of rung spin singlets. The hole pairs can be considered as hard core bosons. The pair hopping matrix element to second order in perturbation theory is

$$t_b = \frac{2t^2}{\frac{3J'}{4} - 2t'} \quad (22)$$

There is also an interaction V_b between two hole pairs on NN rungs. To

second order in perturbation theory,

$$V_b = \frac{4t^2}{\frac{3J'}{4} - 2t'} \quad (23)$$

Both $t_b, V_b \ll J'$ and one can map the ladder model onto an effective hard-core boson model on a chain with NN interaction:

$$H_{eff} = -t_b \sum_i (b_i^\dagger b_{i+1} + H.C.) + V_b \sum_i n_i n_{i+1} \quad (24)$$

b_i^\dagger is the hard core boson creation operator, creating a hole pair on the rung i and $n_i = b_i^\dagger b_i$ is the corresponding number operator. There is a well known mapping between the effective boson model and the quantum XXZ spin model in a magnetic field [19], the Hamiltonian of which is given by

$$H_{xxz} = \sum_i [J_z S_i^z S_{i+1}^z + J_{xy} (S_i^x S_{i+1}^x + S_i^y S_{i+1}^y)] - h \sum_i S_i^z \quad (25)$$

The operator transformations connecting H_{eff} and H_{xxz} are

$$b_j = S_j^\dagger, b_j^\dagger = S_j^-, n_j = 1/2 - S_j^z \quad (26)$$

There is a one-to-one correspondence between the phases of the spin model and those of the boson model. The disordered paramagnetic phase corresponds to the metallic phase for charged bosons. The AFM *Ne'el*-type order in the z direction (when $J_z > J_{xy}$) describes the ordering of bosons on the lattice. For charged bosons, one obtains an insulating charge-ordered phase. The transition from the paramagnetic to the AFM phase represents a metal-insulator transition. The AFM XY order ($J_{xy} > J_z$) is characterised by a two-component order parameter and in the bosonic language corresponds to the off-diagonal long range order of a superfluid condensate. For charged bosons, this is the SC phase.

For the XXZ chain, the asymptotic forms of the correlation functions have been obtained by Luther and Peschel using bosonization theory [20]. For $|\frac{J_z}{J_{xy}}| \leq 1$, the expressions for the correlation functions in the limit of large x and zero magnetic field are:

$$\langle S^z(x, t) S^z \rangle \sim \cos(2k_F x) x^{(-1/\theta)} \quad (27)$$

$$\langle S^\dagger(x,t)S^- \rangle + \langle S^-(x,t)S^\dagger \rangle \sim x^{-\theta} \quad (28)$$

where the exponent

$$\theta = \frac{1}{2} - \pi^{-1} \arcsin(J_z/J_{xy}) \quad (29)$$

For the equivalent bosonic model, the correlation functions corresponding to (27) and (28) are the charge density wave(CDW) correlation $\langle n_r n_0 \rangle$ and the superconducting(SC) correlation $\langle b_r^\dagger b_0 \rangle$. The SC correlations are dominant if $\theta < 1$. For our ladder model, $J_z = V_b$ and $J_{xy} = -2t_b = -V_b$, i.e., for large r both the CDW and SC pairing correlations exist. The transformed Hamiltonian(Eqn.(25)), however, contains a magnetic field term. In the presence of the magnetic field $h(h = V_b)$, the spin chain with $|J_z/J_{xy}| = 1$ is in a spin-flop phase [21] which is equivalent to the SC phase in the bosonic theory. Thus for our ladder model, the SC pairing correlations are dominant for large J' .

III. Conclusion

We have considered a two-chain t-J ladder model for which several exact, analytical results can be derived for the case of two holes. Inclusion of the diagonal exchange and hopping terms enables us to reduce the original N-body (N-2 spins and two holes) problem to an effective two-body problem which is easily solved. The ground state is a bound state of two holes with centre of mass momentum wave vector $K=0$ and total spin $S=0$. The bound state wave function has modified d-wave symmetry. The charge excitation is gapless whereas the spin excitation has a gap. All the results derived by us are in agreement with the numerical results for the conventional two-chain spin-ladder. In the strong coupling limit, our results are the only exact, analytical results for the lightly doped two-chain t-J ladder. For more than two holes, we have not been able, as yet, to calculate the ground state and low lying excitation spectrum exactly and analytically.

Recently, in a remarkable paper [22], Lin, Balents and Fisher have studied weakly interacting electrons hopping on a two-chain ladder. Using bosonization and perturbative renormalisation-group(RG) analysis, they have shown that at half-filling the model scales onto the Gross-Neveu(GN) model. The

GN model happens to be integrable and has $SO(8)$ symmetry. For repulsive interactions, the two-chain ladder exhibits a Mott insulating phase at half-filling with d-wave pairing correlations. The exact energies of all the low-lying excited states can be calculated because of the integrability of the GN model. Lin et al further studied the effects of doping a small density of holes into the d-Mott spin liquid phase at half-filling. Again, for a pair of holes, the ladder system exists in a SG phase with hole binding in the ground state and gapless charge excitations. Scalapino, Zhang and Hanke [23] have considered the strong coupling limit of a two-chain ladder model with local interactions designed to exhibit exact $SO(5)$ symmetry. This model too has a SG phase with hole pairs in the ground state. Numerical calculations on the t-J [15] and Hubbard ladders [24] also show the existence of such a phase. Thus, the SG phase with bound hole pairs appears to be a universal feature of the two-chain ladder system irrespective of the strength of the coupling. This phase also exhibits superconducting pairing correlations. For ladder systems the existence of a SG is favourable for the binding of holes. As mentioned in the Introduction, the existence of a ‘pseudo-SG’ in the cuprates is conjectured to be associated with pre-formed Cooper pairs of holes. This conjecture is supported by our rigorous demonstration that the ground state in the SG phase consists of a bound pair of holes.

Figure Captions

Fig. 1 The spin ladder model described by the $t - t' - J - J'$ Hamiltonian (Eqn.1).

Fig. 2 Exact energy spectrum (ϵ vs K) for the bound state, continuum of scattering states and anti-bound states of two holes ($J = t = t' = 1, J' = 2J$).

Fig. 3 Exact energy spectrum (ϵ vs K) for the bound state, continuum of scattering states and anti-bound states of two holes ($J = 0.25, J' = 2J, t = t' = 1$).


Fig.4 A plot of $|\frac{a(r)}{a(0)}|^2$ vs. r for the ground state wave function of two holes (Eqn.(13)) ($J' = 2J$ (dotted line), $J' = 10J$ (solid line)).


Fig. 5 The spin gap Δ_{SG} vs. J/t ($t = t' = 1.0, J' = 2J$).


References


- [1] Dagotto E and Rice T M 1996 Science 271 618 and references therein; see also 1996 Physics Today 49 17
- [2] Rice T M 1997 Z.Phys. B 103 165 and references therein
- [3] Bose I 1998 Communications in Mathematical and Theoretical Physics 1 219
- [4] Bethe H 1931 Z.Phys.71 205 (English translation in the book The Many-body Problem: An Encyclopedia of Exactly Solved Models in One Dimension Ed. by Mattis D C 1993 World Scientific)
- [5] Voit J 1995 Rep. Prog. Phys. 58 977
- [6] Dagotto E 1994 Rev. Mod. Phys. 66 763
- [7] Kampf A P 1994 Physics Reports 249 219
- [8] Dagotto E, Riera J, Scalapino D J 1992 Phys.Rev.B 45 5744
- [9] Maekawa S 1996 Science 273 1515
- [10] Balakriev F F, Betts J B, Boebinger G S, Motoyama N, Eisaki H and Uchida S cod-mat/9808284
- [11] Bose I and Gayen S 1993 Phys.Rev.B 48 10653
- [12] Bose I and Gayen S 1994 J.Phys.: Condens.Matter 6 L405
- [13] Gayen S and Bose I 1995 J.Phys.: Condens.Matter 7 5871
- [14] Bose I and Gayen S 1996 Physica B 223 and 224 628
- [15] Troyer M, Tsunetsugu H and Rice T M 1996 Phys.Rev.B.53 251
- [16] Murray J D Mathematical Biology, p.705 (Springer-Verlag(1989))
- [17] Balents L and Fisher M P A 1996 Phys.Rev.B.53 12133
- [18] Scalapino D J 1995 Phys.Rep. 250 329


- [19] Jongh L J de 1989 Physica C 161 631
- [20] Luther A J and Peschel I 1975 Phys.Rev.B 12 3908
- [21] Johnson J D 1981 J.Appl.Phys.52 1991
- [22] Lin H H, Balents L and Fisher M P A cond-mat/9801285
- [23] Scalapino D J, Zhang S C and Hanke W cond-mat/9711117
- [24] Noack R M, White S R and Scalapino D J 1996 Physica C 270 281;











■