

**A BEURLING ALGEBRA IS SEMISIMPLE:  
AN ELEMENTARY PROOF**

S.J. BHATT AND H.V. DEDANIA

The Beurling algebra  $L^1(G, \omega)$  on a locally compact Abelian group  $G$  with a measurable weight  $\omega$  is shown to be semisimple. This gives an elementary proof of a result that is implicit in the work of M.C. White (1991), where the arguments are based on amenable (not necessarily Abelian) groups.

Let  $G$  be a locally compact Abelian group with Haar measure  $\lambda$ . A *weight* on  $G$  is a measurable function  $\omega : G \rightarrow (0, \infty)$  such that  $\omega(s+t) \leq \omega(s)\omega(t)$  ( $s, t \in G$ ). Then the *Beurling algebra*  $L^1(G, \omega)$  consists of all complex-valued measurable functions  $f$  on  $G$  such that  $f\omega \in L^1(G)$ . It is a commutative Banach algebra with convolution product and with the norm  $\|f\|_\omega := \int_G |f(s)|\omega(s)d\lambda(s)$ . The authors faced the problem of the semisimplicity of  $L^1(G, \omega)$  in the investigation of the unique uniform norm property in Banach algebras ([1]). It is shown in [5] that if  $G$  is amenable, then there exists a continuous, positive,  $\omega$ -bounded character on  $G$ . Then Lemma 2 (below) quickly implies that  $L^1(G, \omega)$  is semisimple for an Abelian  $G$ . Since the theory of amenable groups is not (yet) a standard part of Harmonic Analysis, and certainly not a part of Abelian Harmonic Analysis, we present an elementary proof of this basic result within the context of Abelian groups.

**THEOREM 1.** *The Beurling algebra  $L^1(G, \omega)$  is semisimple.*

**LEMMA 2.**  *$L^1(G, \omega)$  is either semisimple or radical.*

**PROOF:** Assume that  $L^1(G, \omega)$  is not radical. So its Gelfand space  $\Delta(L^1(G, \omega))$  is non-empty. Let  $\varphi \in \Delta(L^1(G, \omega))$ . Then there exists a function  $\alpha \in L^\infty(G, 1/\omega)$ , the Banach space dual of  $L^1(G, \omega)$ , such that

$$\varphi(f) = \int_G f(s)\alpha(s)d\lambda(s)$$

for all  $f \in L^1(G, \omega)$ . By the standard argument in the case of  $L^1(G)$ , one can show that  $\alpha$  is a continuous function,  $0 < |\alpha(s)| \leq \omega(s)$  ( $s \in G$ ) and  $\alpha(s+t) = \alpha(s)\alpha(t)$  ( $s, t \in G$ ).

---

Received 8th January, 2002

The authors are thankful to Professor H.G. Dales, Leeds (UK), for drawing their attention towards the paper [5].

---

Copyright Clearance Centre, Inc. Serial-fee code: 0004-9727/02 \$A2.00+0.00.

For each  $\theta \in \widehat{G}$ , define  $\alpha_\theta$  by

$$\alpha_\theta(g) = \int_G g(s)\alpha(s)\theta(s)d\lambda(s), \quad g \in L^1(G, \omega).$$

Then  $\alpha_\theta \in \Delta(L^1(G, \omega))$ . Now let  $f \in \text{rad } L^1(G, \omega)$ , the radical of  $L^1(G, \omega)$ . Then  $\alpha_\theta(f) = \widehat{f}(\alpha_\theta) = \widehat{f}\widehat{\alpha}(\theta) = 0$  ( $\theta \in \widehat{G}$ ). Since  $f \in L^1(G, \omega)$ , we have  $f\alpha \in L^1(G)$ . Since  $L^1(G)$  is semisimple and  $\widehat{f}\widehat{\alpha}(\theta) = 0$  ( $\theta \in \widehat{G}$ ), we have  $f\alpha \equiv 0$  almost everhwhere on  $G$ . But  $\alpha(s) \neq 0$  for any  $s \in G$ ; and hence  $f \equiv 0$  almost everywhere on  $G$ . This proves that  $L^1(G, \omega)$  is semisimple.  $\square$

**LEMMA 3.** *Let  $G_1$  be a locally compact Abelian group such that  $L^1(G_1, \omega)$  is semisimple for every weight  $\omega$  on  $G_1$ . Let  $G_2$  be a locally compact Abelian group such that  $L^1(G_2, \omega)$  is semisimple for every weight  $\omega$  on  $G_2$ . Let  $G = G_1 \oplus G_2$  be the direct sum. Then  $L^1(G, \omega)$  is semisimple for every weight  $\omega$  on  $G$ .*

**PROOF:** Let  $\omega$  be a weight on  $G$ . By Lemma 2, it is enough to prove that  $L^1(G, \omega)$  is not radical. Let  $U_1$  and  $U_2$  be symmetric neighbourhoods of the identities in  $G_1$  and  $G_2$  respectively such that their closures are compact. Define  $f = \chi_{U_1 \times U_2}$ , the characteristic function of  $U_1 \times U_2$ . Then  $f$  is a non-zero element of  $L^1(G, \omega)$ . It is clear that  $f^n = \chi_{U_1^n} \chi_{U_2^n}$  for all  $n \in \mathcal{N}$ . It is enough to show that  $\lim_{n \rightarrow \infty} \|f^n\|_\omega^{1/n} > 0$ . So define

$$\begin{aligned} \omega_1(s) &= \omega(s, 0) \quad (s \in G_1) \quad \text{and} \quad \omega_2(s) = \omega(0, s) \quad (s \in G_2); \\ m &= \inf\{\omega_1(s) : s \in U_1\} \quad \text{and} \quad M = \sup\{\omega_2(s) : s \in U_2\}. \end{aligned}$$

It is clear that  $\omega_i$  is a weight on  $G_i$  ( $i = 1, 2$ ). Then by [2, Proposition 2.1],  $m > 0$  and  $M < \infty$ . Also note that for any  $n \in \mathcal{N}$ ,  $\omega_2(s) \leq M^n$  for all  $s \in U_2 + \dots + U_2$  ( $n$ -times) and

$$\begin{aligned} \|f^n\|_\omega &= \int_G |f^n(s, t)| \omega(s, t) d\lambda_1(s) d\lambda_2(t) \\ &= \int_{G_1} \int_{G_2} |\chi_{U_1^n}(s)| |\chi_{U_2^n}(t)| \omega(s, t) d\lambda_1(s) d\lambda_2(t) \\ &\geq \int_{G_1} \int_{G_2} |\chi_{U_1^n}(s)| |\chi_{U_2^n}(t)| \frac{\omega_1(s)}{\omega_2(-t)} d\lambda_1(s) d\lambda_2(t) \\ &= \int_{G_1} |\chi_{U_1^n}(s)| \omega_1(s) d\lambda_1(s) \int_{G_2} |\chi_{U_2^n}(t)| \frac{1}{\omega_2(-t)} d\lambda_2(t) \\ &\geq \|\chi_{U_1^n}\|_{\omega_1} \frac{1}{M^n} \int_{G_2} |\chi_{U_2^n}(t)| d\lambda_2(t) \\ &= \frac{1}{M^n} \|\chi_{U_1^n}\|_{\omega_1} \|\chi_{U_2^n}\|_1, \end{aligned}$$

where  $\|\cdot\|_1$  denotes the  $L^1$ -norm and  $\lambda_i$  denotes the Haar measure on  $G_i$  for  $i = 1, 2$ . Then  $\lim_{n \rightarrow \infty} \|f^n\|_\omega^{1/n} \geq (1/M) \lim_{n \rightarrow \infty} \|\chi_{U_1^n}\|_{\omega_1}^{1/n} \lim_{n \rightarrow \infty} \|\chi_{U_2^n}\|_1^{1/n} > 0$ . This proves that  $L^1(G, \omega)$  is semisimple.  $\square$

PROOF OF THEOREM 1: Note that if  $G$  is a compact Abelian group, then  $L^1(G, \omega) = L^1(G)$  for any weight  $\omega$  on  $G$ ; so it is semisimple. By [3, p. 113],  $L^1(\mathcal{R}, \omega)$  is semisimple for any weight  $\omega$  on  $\mathcal{R}$ ; so Lemma 3 implies that  $L^1(\mathcal{R}^n, \omega)$  is semisimple for any weight  $\omega$  on  $\mathcal{R}^n$ , where  $n \geq 1$ . Hence, again by Lemma 3,  $L^1(\mathcal{R}^n \oplus H, \omega)$  is semisimple for any weight  $\omega$  on  $\mathcal{R}^n \oplus H$ , where  $n \geq 0$  and  $H$  is a compact Abelian group.

Now let  $G$  be an arbitrary locally compact Abelian group and let  $\omega$  be a weight on  $G$ . By [4, Theorem 2.4.1], there exists an open subgroup  $G_1$  of  $G$  such that  $G_1 = \mathcal{R}^n \oplus H$ , where  $n \geq 0$  and  $H$  is a compact Abelian group. By above argument  $L^1(G_1, \omega|_{G_1})$  is semisimple. But the later is a closed subalgebra of  $L^1(G, \omega)$ . Hence  $L^1(G, \omega)$  is not radical. Thus it is semisimple due to Lemma 2.  $\square$

#### REFERENCES

- [1] S.J. Bhatt and H.V. Dedania, 'Banach algebras with unique uniform norm II', *Studia Math.* **147** (2001), 211–235.
- [2] H.A.M. Dzinotyiweyi, 'Weighted function algebras on groups and semigroups', *Bull. Austral. Math. Soc.* **33** (1986), 307–318.
- [3] I. Gelfand, D. Raikov and G. Shilov, *Commutative normed rings* (Chelsea Publication Company, New York, 1964).
- [4] W. Rudin, *Fourier analysis on groups*, Interscience Tracts in Pure and Applied Mathematics **12** (Interscience Publishers, New York, London, 1962).
- [5] M.C. White, 'Characters on weighted amenable groups', *Bull. London Math. Soc.* **23** (1991), 375–380.

Department of Mathematics  
 Sardar Patel University  
 Vallabh Vidyanagar - 388 120  
 Gujarat  
 India  
 e-mail: [sjb@spu.ernet.in](mailto:sjb@spu.ernet.in)  
[haresh@spu.ernet.in](mailto:haresh@spu.ernet.in)