

# Kinetics and Mechanism of Complex Formation of some Bivalent and Trivalent Metal Ions with Pentaammine-(nitrilotriacetato)cobalt(III) in Aqueous Medium†

Rita Das,<sup>a</sup> Nigamananda Das<sup>b</sup> and Anadi C. Dash<sup>\*c</sup>

<sup>a</sup> Department of Chemistry, M. P. C. College, Baripada-757 001, Orissa, India

<sup>b</sup> Regional Research Laboratory, Bhubaneswar-751 013, Orissa, India

<sup>c</sup> Department of Chemistry, Utkal University, Bhubaneswar-751 004, Orissa, India

The kinetics of reversible complex formation of  $\text{Ni}^{II}$ ,  $\text{Co}^{II}$  and  $\text{Cu}^{II}$  with the pentaammine-(nitrilotriacetato)cobalt(III) ion,  $[\text{Co}(\text{NH}_3)_5(\text{H}_2\text{nta})]^{2+}$  ( $\text{H}_3\text{nta}$  = nitrilotriacetic acid) have been investigated at  $0.0025 \leq [\text{M}^{2+}] \leq 0.04$ ,  $0.004 \leq [\text{H}^+] \leq 0.05 \text{ mol dm}^{-3}$ ,  $10.0 \leq T \leq 40.0 \text{ }^\circ\text{C}$  and  $I = 0.3 \text{ mol dm}^{-3}$ . The rate constants for the formation of the binuclear species are at least  $10^3$  times less than the water exchange rate constants of  $[\text{M}(\text{OH}_2)_6]^{2+}$  under comparable conditions. General base catalysis indicated that proton transfer from the  $\text{NH}^+$  site of the co-ordinated ligand (nta) is involved in the rate determining step. The binuclear species undergo dissociation *via* spontaneous and acid-catalysed paths. The low values of spontaneous dissociation rate constants also support the chelate nature of the binuclear species. It is likely that the nta moiety of  $(\text{NH}_3)_5\text{Co}(\text{nta})$  acts at least as a tridentate ligand and the chelate ring closure/opening *via* N–M<sup>II</sup> bond formation/dissociation is rate limiting. Complex formation with  $\text{Fe}^{III}$  and  $\text{Al}^{III}$  has been investigated at  $15\text{--}35 \text{ }^\circ\text{C}$  ( $I = 1.0 \text{ mol dm}^{-3}$ ) and  $25 \text{ }^\circ\text{C}$  ( $I = 0.3 \text{ mol dm}^{-3}$ ), respectively. General base catalysis was not observed for these trivalent metal ions. The  $[\text{M}(\text{OH}_2)_5(\text{OH})]^{2+}$  species reacted faster than  $[\text{M}(\text{OH}_2)_6]^{3+}$ . The reaction of  $[\text{M}(\text{OH}_2)_6]^{3+}$  may involve an associative interchange mechanism while that for  $[\text{M}(\text{OH}_2)_5(\text{OH})]^{2+}$  involves dissociative interchange.

The aquation of pentaammine(nitrilotriacetato)cobalt(III),  $[\text{Co}(\text{NH}_3)_5(\text{H}_2\text{nta})]^{2+}$  ( $\text{H}_3\text{nta}$  = nitrilotriacetic acid), is reported<sup>1</sup> to be catalysed by metal ions *via* formation of reactive binuclear complexes. The kinetics and mechanism of formation/dissociation of binuclear species involving bivalent and trivalent metal ions and  $[\text{Co}(\text{NH}_3)_5(\text{H}_2\text{nta})]^{2+}$  have not been reported. As a sequel to our current research interest on the study of the mechanism of formation/dissociation of binuclear complexes of metal ions with pentaammine-(carboxylato)cobalt(III) substrates<sup>2–9</sup> we undertook the present study. The complex  $[\text{Co}(\text{NH}_3)_5(\text{H}_2\text{nta})]^{2+}$  can exist in different stages of protonation under varying pH conditions resulting in species of varying charge with multiple co-ordinating sites. Thus the role of varying charge of the cobalt(III) substrate on the kinetics and mechanism of binuclear complex formation with some bivalent and trivalent metal ions could be examined. Further, the complexation kinetics of  $\text{Cu}^{II}$  is generally fast and requires the use of a relaxation technique to be followed. Interestingly the pentaammine(nitrilotriacetato)-cobalt(III) cation proved suitable for the study of its complexation with  $\text{Cu}^{II}$  by a conventional stopped-flow technique.

## Experimental

**Materials and Methods.**—Pentaammine(nitrilotriacetato)-cobalt(III) perchlorate,  $[\text{Co}(\text{NH}_3)_5(\text{H}_2\text{nta})][\text{ClO}_4]_2$  ( $\text{H}_3\text{nta}$  =  $\text{HN}^+(\text{CH}_2\text{CO}_2^-)(\text{CH}_2\text{CO}_2\text{H})_2$ ) was prepared as described earlier.<sup>1</sup> The purity of the sample was checked by analysis of


cobalt, and the UV/VIS spectral data of the complex agreed well with those reported earlier.<sup>1</sup> The FT-IR spectra of  $[\text{Co}(\text{NH}_3)_5(\text{H}_2\text{nta})]^{2+}$  displayed bands at  $1750$  and  $1632 \text{ cm}^{-1}$  attributed to free ( $\nu_{\text{Co},\text{H}}$ ) and cobalt(III)-bound carboxylate ( $\nu_{\text{Co},\text{M}}$ ) of the nta moiety, respectively.<sup>10</sup> The latter band was broad presumably owing to the NH-deformation mode.

The stock solutions of  $\text{Ni}^{II}$ ,  $\text{Co}^{II}$ ,  $\text{Cu}^{II}$ ,  $\text{Al}^{III}$  and  $\text{Fe}^{III}$  perchlorates were prepared and estimated as described earlier.<sup>11</sup> 2,6-Dimethylpyridine (Fluka, AG) was used whenever required. All other chemicals used for kinetic studies were of AR grade. Freshly prepared doubly distilled water was used in all experiments. The dissociation constants of the unbound carboxyl and protonated amine functions were determined potentiometrically at  $15\text{--}35 \text{ }^\circ\text{C}$ ,  $I = 0.3$  and  $1.0 \text{ mol dm}^{-3}$ . The pH measurements were made with an Elico digital model LI-120 pH-meter, using glass–Ag–AgCl,  $\text{NaCl}$  ( $2.0 \text{ mol dm}^{-3}$ ) combined electrode model CL 51.

**Kinetic Measurements.**—The kinetics of reversible formation of binuclear complexes were studied under pseudo-first order conditions ( $5 < [\text{M}^{2+/(3+)}]/[\text{complex}]_T < 50$ ) at suitable wavelengths (see footnotes to Tables 1, 5 and 6) using a fully automated Hi-Tech (UK) SF-51 stopped-flow spectrophotometer. For acid-catalysed dissociation the preformed binuclear complex was acidified rapidly in the stopped-flow assembly as described earlier.<sup>7</sup> In every case absorbance–time data for any run fitted well to a single exponential equation applicable to first-order kinetics. At least seven replicate measurements were made for each run to compute  $k_{\text{obs}}$  and its standard deviation.

**Dissociation Constants.**—The potentiometric titration using the method of Irving and Rossotti<sup>12</sup> yielded the acid dissociation constants [equation (1)] of the complex [ $\text{p}K_1 = 2.13$  (2.05), 2.09 (1.96) and 2.03 (1.96);  $\text{p}K_2 = 8.20$  (8.15),

† Supplementary data available (No. Sup 57104, 11 pp.):  $k_{\text{obs}}$  values at different temperatures. See Instructions for Authors, *J. Chem. Soc., Dalton Trans.*, 1995, Issue 1, pp. xxv–xxx.



7.91 (7.80) and 7.63 (7.55) at  $I = 0.3$  (1.0) mol dm<sup>-3</sup> and 15, 25 and 35 °C, respectively] in satisfactory agreement with those reported earlier [ $\text{p}K_1 = 1.82 \pm 0.07$  ( $1.74 \pm 0.2$ );  $\text{p}K_2 = 7.96 \pm 0.01$  ( $8.3 \pm 0.2$ ) at 25 °C (6 °C) and  $I = 1.0$  mol dm<sup>-3</sup>].<sup>1</sup>

## Results and Discussion

*Reversible Formation of Binuclear Complexes Between Bivalent Metal Ions and  $[\text{Co}(\text{NH}_3)_5(\text{H}_2\text{nta})]^{2+}$ .*—The interaction of the bivalent metal ions with  $[\text{Co}(\text{NH}_3)_5(\text{H}_2\text{nta})]^{2+}$  is indicated by the instantaneous increase in absorbance in the range 260–320 nm (see Fig. 1). Our observations are in agreement with those of Cannon and Gardiner.<sup>1</sup> The pseudo-first-order rate constants [see equation (2)] were collected at varying  $[\text{H}^+]$  (0.004–0.05 mol dm<sup>-3</sup>),  $[\text{M}^{2+}]$  (0.005–0.04 for  $\text{Co}^{\text{II}}$  or  $\text{Ni}^{\text{II}}$  and 0.0025–0.020 for  $\text{Cu}^{\text{II}}$ ) and are available as supplementary material (sup no. 57104). The plots of  $k_{\text{obs}}$  versus  $[\text{M}^{2+}]$  are linear with positive intercepts on the rate axis; the slopes, however, increase with decreasing  $[\text{H}^+]$ . This may be attributed to the reduced reactivity of  $[\text{Co}(\text{NH}_3)_5(\text{H}_2\text{nta})]^{2+}$  relative to that of  $[\text{Co}(\text{NH}_3)_5(\text{Hnta})]^+$ . Plausible reaction pathways are given in Scheme 1. Accordingly  $k_{\text{obs}}$  takes the form given in equation (2), where the overall

$$k_{\text{obs}} = k_f[\text{M}^{2+}] + k_r \quad (2)$$

formation rate constant ( $k_f$ ) and the dissociation rate constant ( $k_r$ ) of the binuclear species,  $[(\text{NH}_3)_5\text{Co}(\text{nta})\text{M}]^{2+}$ , are given by equations (3) and (4), respectively. The  $k_f$  and  $k_r$  values are

$$k_f = \{k_1([\text{H}^+/K_1) + k_2\}/(1 + [\text{H}^+/K_1]) \quad (3)$$

$$k_r = k_{-1}[\text{H}^+]^2 + k_{-2}[\text{H}^+] \quad (4)$$

obtained from the gradient and intercept of the least-squares best line plot of  $k_{\text{obs}}$  versus  $[\text{M}^{2+}]$  at constant  $[\text{H}^+]$ . The  $[\text{H}^+]$  dependence of  $k_f$  enabled the calculation of  $k_1$  and  $k_2$  using equation (3) (see Table 1).

The  $[\text{M}(\text{OH}_2)_6]^{2+}$  ( $\text{M}^{2+} = \text{Ni}^{\text{II}}$ ,  $\text{Co}^{\text{II}}$  or  $\text{Cu}^{\text{II}}$ ) ions are believed to undergo ligand substitution *via* a dissociative interchange mechanism ( $\text{I}_d$ ).<sup>14–17</sup> The complexation of these metal ions involves prior formation of the outer-sphere ion pair characterised by the equilibrium constant,  $K_{\text{os}}$ . The sequential displacement of three aqua ligands by the multidentate  $[\text{Co}(\text{NH}_3)_5(\text{H}_2\text{nta})]^{2+}$  ( $= \text{H}_2\text{L}^{2+}$ ) may be delineated as in Scheme 2, where the substitution of any one of three aqua ligands may be rate determining. If substitution of the first aqua ligand in  $[\text{M}(\text{OH}_2)_6]^{2+}$  is rate determining then  $k_1'$  ( $= k_1/K_{\text{os}}$ ) or  $k_2'$  ( $= k_2/K_{\text{os}}'$ ) should be comparable to the water exchange-rate constant of the aqua metal ions ( $k_{\text{ex}} = 3.2 \times 10^4$ ,  $3.2 \times 10^6$  and  $4.4 \times 10^9$  s<sup>-1</sup> at 25 °C for  $[\text{Ni}(\text{OH}_2)_6]^{2+}$ ,  $[\text{Co}(\text{OH}_2)_6]^{2+}$  and  $[\text{Cu}(\text{OH}_2)_6]^{2+}$ , respectively).<sup>13,18</sup> Using Fuoss's equation<sup>19</sup> and assuming a distance of closest approach  $a = 5$  Å  $K_{\text{os}}'$  was calculated to be 0.07 dm<sup>3</sup> mol<sup>-1</sup> at 25 °C for a 2+–1+ charge outer-sphere interaction. An even lower value ( $\approx 0.05$  dm<sup>3</sup> mol<sup>-1</sup>) of  $K_{\text{os}}$  is expected for 2+–2+ charge association. At 25 °C the values of  $k_1'$  ( $\approx 60$ ,  $1.2 \times 10^3$  and  $2.3 \times 10^5$  s<sup>-1</sup>) and  $k_2'$  ( $\approx 136$ ,  $3.0 \times 10^3$  and  $7.1 \times 10^5$  s<sup>-1</sup>, for  $\text{Ni}^{\text{II}}$ ,  $\text{Co}^{\text{II}}$  and  $\text{Cu}^{\text{II}}$ , respectively) estimated from  $K_{\text{os}}(K_{\text{os}}')$  are two to four orders of magnitude smaller than the water-exchange rate constants of  $\text{Ni}^{\text{II}}$ ,  $\text{Co}^{\text{II}}$  and  $\text{Cu}^{\text{II}}$ . The rate constants for the formation of  $[(\text{NH}_3)_5\text{Co}(\text{nta})\text{M}]^{2+}$  via






Fig. 1 Spectral evidence for the formation of binuclear complexes between  $[\text{Co}(\text{NH}_3)_5(\text{H}_2\text{nta})]^{2+}$  and  $\text{Cu}^{2+}$  at  $[\text{H}^+] = 1.0 \times 10^{-2}$ ,  $I = 0.3$  mol dm<sup>-3</sup>: (1)  $[\text{Cu}^{2+}] = 1.0 \times 10^{-2}$ , (2)  $[\text{complex}]_T = 5.0 \times 10^{-3}$  mol dm<sup>-3</sup> and (3)  $[\text{Cu}^{2+}] = 1.0 \times 10^{-2}$ ,  $[\text{complex}]_T = 5.0 \times 10^{-3}$  mol dm<sup>-3</sup>



Scheme 1  $\text{R} = (\text{NH}_3)_5\text{Co}^{\text{III}}$ ,  $\text{M} = \text{Ni}^{\text{II}}$ ,  $\text{Co}^{\text{II}}$  or  $\text{Cu}^{\text{II}}$



Scheme 2

the  $k_2$  path are, however, comparable to the rate constants for the formation of  $[\text{M}(\text{nta})]^-$  by the reaction of  $\text{M}^{2+}$  ( $\text{M}^{2+} = \text{Ni}^{\text{II}}$  or  $\text{Cu}^{\text{II}}$ ) with free  $\text{Hnta}^{2-}$  (see Table 2) reported earlier.<sup>20–23</sup> This strongly suggests that the rate-determining step is not the loss of the first water molecule from the inner co-ordination sphere of the aqua metal ion. The relatively low rate constants for the formation of binuclear species, either by


**Table 1** Rate data and activation parameters for the formation of  $[(\text{NH}_3)_5\text{Co}(\text{nta})\text{M}]^{2+}$ <sup>a</sup>

M	T/°C	$k_1/\text{dm}^3 \text{ mol}^{-1} \text{ s}^{-1}$	$k_2/\text{dm}^3 \text{ mol}^{-1} \text{ s}^{-1}$	$\Delta H^\ddagger$ <sup>b</sup> /kJ mol <sup>-1</sup>	$\Delta S^\ddagger/\text{J K}^{-1} \text{ mol}^{-1}$
Ni <sup>II</sup>	20.0	2.9 ± 0.4	5.8 ± 0.3	62 ± 5 (49 ± 3)	-26 ± 15 (-63 ± 8)
	25.0	4.2 ± 0.4	6.8 ± 0.7		
	30.0	6.0 ± 0.2	10.5 ± 0.9		
	35.0	10.6 ± 0.6	17.7 ± 1.3		
	40.0	13.9 ± 0.5	21.7 ± 0.2		
Co <sup>II</sup>	15.0	31 ± 12	75 ± 7	83 ± 8 (52 ± 17)	70 ± 29 (-29 ± 57)
	20.0	46 ± 3	92 ± 4		
	25.0	84 ± 7	150 ± 9		
Cu <sup>II</sup>	10.0	(4.5 ± 0.4) × 10 <sup>3</sup>	(15.2 ± 0.6) × 10 <sup>3</sup>	62 ± 19 (38 ± 3)	51 ± 60 (-30 ± 6)
	15.0	(6.9 ± 1.0) × 10 <sup>3</sup>	(18.8 ± 1.6) × 10 <sup>3</sup>		
	20.0	(10.7 ± 1.0) × 10 <sup>3</sup>	(27.4 ± 1.4) × 10 <sup>3</sup>		
	25.0	(16.3 ± 0.6) × 10 <sup>3</sup>	(35.9 ± 1.2) × 10 <sup>3</sup>		

<sup>a</sup> [Complex]<sub>T</sub> = (0.5–2.0) × 10<sup>-3</sup>, [Ni<sup>2+</sup>] or [Co<sup>2+</sup>] = (0.5–5.0) × 10<sup>-2</sup>, [Cu<sup>2+</sup>] = (0.25–2.0) × 10<sup>-2</sup>, I = 0.3 mol dm<sup>-3</sup>,  $\lambda$  = 260 for Ni<sup>2+</sup> and Co<sup>2+</sup> and 290 nm for Cu<sup>2+</sup>. <sup>b</sup> Values for  $k_2$  ( $k_1$  values in parentheses);  $\Delta H_{\text{ex}}^\ddagger$ /kJ mol<sup>-1</sup> ( $\Delta S_{\text{ex}}^\ddagger/\text{J K}^{-1} \text{ mol}^{-1}$ ) = 57 (32) and 47 (37) for the H<sub>2</sub>O exchange reaction of [Ni(OH<sub>2</sub>)<sub>6</sub>]<sup>2+</sup> and [Co(H<sub>2</sub>O)<sub>6</sub>]<sup>2+</sup> respectively (ref. 13).

**Table 2** Comparison of rate parameters for the formation of bivalent metal complexes with free and partially co-ordinated nitrilotriacetate at 25 °C

Reaction	I/mol dm <sup>-3</sup>	$k_f/\text{dm}^3 \text{ mol}^{-1} \text{ s}^{-1}$	Refs.
Ni <sup>2+</sup> + Hnta <sup>2-</sup>	1.25	7.51	20
	0.20	4.0	21
Ni <sup>2+</sup> + [Co(NH <sub>3</sub> ) <sub>5</sub> (Hnta)] <sup>+</sup>	0.30	6.8 ± 0.7	This work
	—	5.0 × 10 <sup>3</sup>	22
Cu <sup>2+</sup> + Hnta <sup>2-</sup>	0.05	1.12 × 10 <sup>5</sup>	23
	0.30	(36 ± 1) × 10 <sup>3</sup>	This work
Co <sup>2+</sup> + [Co(NH <sub>3</sub> ) <sub>5</sub> (Hnta)] <sup>+</sup>	0.30	150 ± 9	This work

**Scheme 3** M<sup>2+</sup> = Ni<sup>2+</sup>, Co<sup>2+</sup> or Cu<sup>2+</sup>

path  $k_1$  or  $k_2$ , indicate that chelate-ring formation through N (see Scheme 3) is rate limiting. A similar interpretation has been made in studies of a number of metal aminopoly-carboxylic acid complexes.<sup>20–26</sup> If so, then either the proton migrates from the N atom to the solvent before chelate-ring closure, or proton transfer from the NH<sup>+</sup> site and chelate-ring closure are acting in concert. Additional evidence of rate-limiting proton transfer from the NH<sup>+</sup> centre is obtained from buffer catalysis, *i.e.*  $k_{\text{obs}}$  at a fixed pH and [Ni<sup>2+</sup>] is found to increase with increasing 2,6-dimethylpyridine concentration (see Table 3). The activation parameters ( $\Delta H^\ddagger$  and  $\Delta S^\ddagger$ ) are also substantially different from those reported for the water-exchange reaction of [M(OH<sub>2</sub>)<sub>6</sub>]<sup>2+</sup> ions (see Table 1) { $\Delta H^\ddagger$  = 57, 47 and 17 kJ mol<sup>-1</sup>;  $\Delta S^\ddagger$  = 32, 37 and -44 J K<sup>-1</sup> mol<sup>-1</sup> for [Ni(OH<sub>2</sub>)<sub>6</sub>]<sup>2+</sup>, [Co(OH<sub>2</sub>)<sub>6</sub>]<sup>2+</sup> and [Cu(MeOH)<sub>6</sub>]<sup>2+</sup>, respectively}<sup>13</sup>.

Rate data for the acid-catalysed dissociation of binuclear complexes,  $[(\text{NH}_3)_5\text{Co}(\text{nta})\text{M}]^{2+}$  at various temperatures and [H<sup>+</sup>] (I = 0.3 mol dm<sup>-3</sup>) are available as supplementary data. The observed rate constants were corrected for the formation reaction by utilising the values of  $k_1$ ,  $k_2$  and  $K_1$ . The plots of the corrected pseudo-first-order rate constant,  $k_r$  (=  $k_{\text{obs}} -$

**Table 3** Effect of 2,6-dimethylpyridine concentration on the rate of formation of  $[(\text{NH}_3)_5\text{Co}(\text{nta})\text{Ni}]^{2+}$  at 25 °C\*

[C <sub>5</sub> H <sub>3</sub> Me <sub>2</sub> N]/mol dm <sup>-3</sup>	[HClO <sub>4</sub> ]/mol dm <sup>-3</sup>	pH	$k_{\text{obs}}/\text{s}^{-1}$
0.01	0.005	6.72	60 ± 4
0.02	0.010	6.73	70 ± 4
0.03	0.015	6.74	75 ± 7
0.04	0.020	6.72	80 ± 7
0.05	0.025	6.77	85 ± 9
0.06	0.030	6.76	90 ± 8

\* [Complex]<sub>T</sub> = 1.0 × 10<sup>-3</sup>, [Ni<sup>2+</sup>] = 2.0 × 10<sup>-2</sup>, I = 0.3 mol dm<sup>-3</sup>;  $\lambda$  = 290 nm.

$k_r[\text{Ni}^{2+}]$ ) versus [H<sup>+</sup>] are not linear; however, the plots of  $k_r/[\text{H}^+]$  versus [H<sup>+</sup>] yielded good straight lines, consistent with equation (4) (see Fig. 2). The values of  $k_{-1}$  and  $k_{-2}$  obtained from the slopes and intercepts of such plots and their associated activation parameters are collected in Table 4.

A comparison of the dissociation rate constants,  $k_2$ , for  $[(\text{NH}_3)_5\text{Co}(\text{nta})\text{Ni}]^{2+}$  (5.0) and  $[\text{Ni}(\text{nta})]^-$  (0.77 dm<sup>3</sup> mol<sup>-1</sup> s<sup>-1</sup>) at 25 °C, I = 0.3 and 1.0 mol dm<sup>-3</sup><sup>20</sup> showed that the former is only six times kinetically more labile than the latter. Analogous rate comparisons for Co<sup>2+</sup> and Cu<sup>2+</sup> cannot be made due to non-availability of rate data for the acid-catalysed dissociation of  $[\text{Co}(\text{nta})]^-$  and  $[\text{Cu}(\text{nta})]^-$ . However, the small rate difference for Ni<sup>2+</sup> indicates that the M<sup>2+</sup> ions are chelated by the nta moiety in  $[(\text{NH}_3)_5\text{Co}(\text{nta})\text{M}]^{2+}$ , H<sup>+</sup>-promoted Ni–N bond cleavage being the most likely rate-determining step. Indeed, Bydalek and Blomster<sup>20</sup> have suggested that Ni–N bond cleavage is rate determining in the dissociation of  $[\text{Ni}(\text{nta})]^-$ .

**Reversible Formation of Binuclear Complexes of M<sup>3+</sup> Ions with  $[\text{Co}(\text{NH}_3)_5(\text{H}_2\text{nta})]^{2+}$ .**—The observed pseudo-first-order rate constants for the formation of Fe<sup>3+</sup> are available as supplementary material while those of Al<sup>3+</sup> are presented in Table 5. Plots of  $k_{\text{obs}}$  versus [M<sup>3+</sup>] at fixed [H<sup>+</sup>] are linear with positive intercepts and gradients which are identified with the overall dissociation and formation rate constants of the

binuclear species, respectively. Scheme 4 delineates the overall reaction sequence for which  $k_{\text{obs}}$  is given by equation (5) where

$$k_{\text{obs}} = k_f f_1 f_2 [\text{M}^{3+}] + k_r \quad (5)$$




Fig. 2 Plots of  $k_r/[\text{H}^+]$  ( $k_r = k_{\text{obs}} - k_f[\text{Ni}^{2+}]$ ) versus  $[\text{H}^+]$  for the acid-catalysed dissociation of  $[(\text{NH}_3)_5\text{Co}(\text{NTA})\text{Ni}]^{2+}$  at 20 (a), 25 (b), 30 (c), 35 (d) and 40 °C (e)

$k_f, k_r, f_1$  and  $f_2$  are defined as in equations (6)–(9). Values of  $K_h$

$$k_f = k_1 + (k_2 K_h + k_3 K_1)/[\text{H}^+] + k_4 K_h/[\text{H}^+]^2 \quad (6)$$

$$k_r = k_{-4} + (k_{-2} + k_{-3})[\text{H}^+] + k_{-1}[\text{H}^+]^2 \quad (7)$$

$$f_1 = [\text{H}^+]/([\text{H}^+] + K_h) \quad (8)$$

$$f_2 = [\text{H}^+]/([\text{H}^+] + K_1) \quad (9)$$

for  $[\text{Fe}(\text{OH}_2)_6]^{3+}$  [ $(1.2\text{--}2.3) \times 10^{-3}$  mol dm $^{-3}$  at 20–35 °C,  $I = 1.0$  mol dm $^{-3}$ ]<sup>27</sup> and of  $K_1$  (see Experimental section) are such that  $f_1$  and  $f_2$  reduce virtually to unity in the experimental acidity range so that equation (5) reduces to equation (10).

$$k_{\text{obs}} = k_f[\text{Fe}^{3+}] + k_r \quad (10)$$

However, for  $\text{Al}^{3+}$ ,  $f_1 = 1$  and  $f_2$  is significantly less than 1 in the acidity range employed ( $K_h$  of  $[\text{Al}(\text{OH}_2)_6]^{3+} = 1.0 \times 10^{-5}$  mol dm $^{-3}$  at 25 °C,  $I = 0.1$  mol dm $^{-3}$ )<sup>28</sup> and hence equation (5) reduces to equation (11). Linear plots of  $k_{\text{obs}}$  versus  $[\text{Fe}^{3+}]$  or

$$k_{\text{obs}} = k_f[\text{Al}^{3+}]\{[\text{H}^+]/([\text{H}^+] + K_1)\} + k_r \quad (11)$$

$[\text{Al}^{3+}]f_2$  yielded  $k_f$  and  $k_r$  values as the slopes and intercepts, respectively and good linear plots of  $k_f[\text{H}^+]$  against  $[\text{H}^+]^{-1}$  (Fig. 3) were also obtained. Hence  $k_1$  is insignificant, *i.e.*, the dipositive form of the complex,  $[\text{Co}(\text{NH}_3)_5(\text{H}_2\text{NTA})]^{2+}$  does not react with  $[\text{M}(\text{OH}_2)_6]^{3+}$ . A similar observation was made in the complexation reaction of  $\text{Fe}^{3+}$  with pentaammine(oxalato)cobalt(III).<sup>2</sup> The values of  $k_4 K_h K_1$  (and  $k_4$ ) and  $(k_2 K_h + k_3 K_1)$  calculated from the gradients and intercepts of  $k_f[\text{H}^+]$  versus  $[\text{H}^+]^{-1}$  plots are collected in Table 6. The values of  $k_2$

Table 4 Rate and activation parameters for the dissociation of  $[(\text{NH}_3)_5\text{Co}(\text{NTA})\text{M}]^{2+}$ <sup>a</sup>

M	T/°C	$k_{-1}/\text{dm}^6 \text{ mol}^{-2} \text{ s}^{-1}$	$k_{-2}/\text{dm}^3 \text{ mol}^{-1} \text{ s}^{-1}$	$\Delta H^\ddagger$ <sup>b</sup> /kJ mol $^{-1}$	$\Delta S^\ddagger$ <sup>b</sup> /J K $^{-1}$ mol $^{-1}$
Ni <sup>II</sup>	20.0	$3.5 \pm 0.5$	$3.6 \pm 0.1$	$45 \pm 8$ ( $45 \pm 5$ )	$80 \pm 28$ ( $-80 \pm 16$ )
	25.0	$8.0 \pm 0.3$	$5.0 \pm 0.1$		
	30.0	$14.1 \pm 1.0$	$5.6 \pm 0.2$		
	35.0	$15.2 \pm 0.7$	$7.8 \pm 0.1$		
	40.0	$18.8 \pm 0.9$	$12.2 \pm 0.1$		
Co <sup>II</sup>	15.0	$(3.0 \pm 0.2) \times 10^2$	$(3.9 \pm 0.1) \times 10^2$	$79 \pm 10$ ( $55 \pm 6$ )	$76 \pm 36$ ( $-14 \pm 18$ )
	20.0	$(6.0 \pm 0.4) \times 10^2$	$(6.9 \pm 0.1) \times 10^2$		
	25.0	$(9.2 \pm 0.7) \times 10^2$	$(9.4 \pm 0.1) \times 10^2$		
Cu <sup>II</sup>	10.0	$(12.5 \pm 0.1) \times 10^2$	$(2.3 \pm 0.3) \times 10^2$	$40 \pm 2$ ( $18 \pm 8$ )	$-41 \pm 6$ ( $-137 \pm 27$ )
	15.0	$(16.5 \pm 1.4) \times 10^2$	$(3.2 \pm 0.3) \times 10^2$		
	20.0	$(24.0 \pm 0.6) \times 10^2$	$(3.6 \pm 0.1) \times 10^2$		
	25.0	$(29.7 \pm 1.5) \times 10^2$	$(3.7 \pm 0.3) \times 10^2$		

<sup>a</sup> [Complex]<sub>T</sub> =  $(0.5\text{--}1.0) \times 10^{-3}$ ,  $[\text{M}^{2+}] = (0.5\text{--}1.0) \times 10^{-2}$ ,  $I = 0.3$  mol dm $^{-3}$ ;  $\lambda = 260$  nm for  $\text{Ni}^{2+}$  and  $\text{Co}^{2+}$  and 290 nm for  $\text{Cu}^{2+}$ . <sup>b</sup> Values evaluated from dependence of temperature on  $k_{-1}$  ( $k_{-2}$ ).

Table 5 Rate data for the formation of  $[(\text{NH}_3)_5\text{Co}(\text{NTA})\text{Al}]^{3+}$  at 25 °C<sup>a</sup>

$10^2 [\text{Al}^{3+}]$ /mol dm $^{-3}$	$10^3 k_{\text{obs}}/\text{s}^{-1}$ <sup>b</sup>			
	$(2.55 \pm 0.02)^b$	$(2.70 \pm 0.02)$	$(2.85 \pm 0.03)$	$(3.05 \pm 0.02)$
0.50	$12.0 \pm 0.2$	$14.1 \pm 0.8$	$16.0 \pm 0.7$	$17.2 \pm 1.9$
1.00	$13.5 \pm 0.6$	$15.8 \pm 0.8$	$18.5 \pm 0.4$	$21.2 \pm 1.1$
1.50	$14.2 \pm 0.3$	$17.4 \pm 0.5$	$20.7 \pm 1.4$	$25.6 \pm 1.4$
2.00	$15.6 \pm 0.1$	$19.0 \pm 0.8$	$23.1 \pm 0.6$	$27.8 \pm 1.6$
2.50	$17.2 \pm 0.8$	$19.8 \pm 0.6$	$25.6 \pm 0.4$	$31.7 \pm 1.6$
3.00	$17.6 \pm 0.2$	$21.0 \pm 0.6$	$29.0 \pm 1.2$	$35.0 \pm 0.5$
4.00	$19.9 \pm 0.6$	$25.0 \pm 0.6$	—	—
$k_f/\text{dm}^3 \text{ mol}^{-1} \text{ s}^{-1}$	$0.71 \pm 0.02$	$1.15 \pm 0.05$	$2.45 \pm 0.07$	$5.2 \pm 0.1$
$10^3 k_r/\text{s}^{-1}$	$10.9 \pm 0.1$	$12.6 \pm 0.4$	$13.6 \pm 0.3$	$14.0 \pm 0.4$


$k_4 = 85 \pm 10$  dm $^3$  mol $^{-1}$  s $^{-1}$   $10^3 (k_2 K_h + k_3 K_1) = 1.0 \pm 0.3$  s $^{-1}$   $k_2 = 48$  dm $^3$  mol $^{-1}$  s $^{-1}$   $10^2 k_3 = 6.2$  dm $^3$  mol $^{-1}$  s $^{-1}$

<sup>a</sup> [Complex]<sub>T</sub> =  $1.0 \times 10^{-3}$ ,  $I = 0.3$  mol dm $^{-3}$ ,  $\lambda = 270$  nm. <sup>b</sup> Values in parentheses denote pH.

**Table 6** Rate and activation parameters for the formation of  $[(\text{NH}_3)_5\text{Co}(\text{nta})\text{Fe}]^{3+}$ <sup>a</sup>

$T/^\circ\text{C}$	$10^2 [\text{H}^+]/\text{mol dm}^{-3}$	$k_f/\text{dm}^3 \text{mol}^{-1} \text{s}^{-1}$	$10^2 (k_2 K_h + k_3 K_1)/\text{s}^{-1}$	$10^{-3} k_4/ \text{dm}^3 \text{mol}^{-1} \text{s}^{-1}$	$10^{-2} k_2/ \text{dm}^3 \text{mol}^{-1} \text{s}^{-1}$	$k_3/\text{dm}^3 \text{mol}^{-1} \text{s}^{-1}$	$k_r(\text{av.})^b/\text{s}^{-1}$
15.0	5.0	$21.1 \pm 0.5$	$43 \pm 3$	$2.7 \pm 0.3$	2.4	24	$0.06 \pm 0.02$
	10.0	$8.0 \pm 0.5$					
	15.0	$5.3 \pm 0.3$					
	20.0	$3.7 \pm 0.4$					
20.0	5.0	$43 \pm 3$	$85 \pm 10$	$5.4 \pm 1.0$	3.5	42	$0.11 \pm 0.03$
	10.0	$14.5 \pm 1.2$					
	15.0	$8.8 \pm 0.3$					
	20.0	$5.2 \pm 0.6$					
25.0	5.0	$75 \pm 4$	$186 \pm 17$	$6.2 \pm 1.2$	5.7	85	$0.19 \pm 0.08$
	10.0	$30.0 \pm 1.2$					
	15.0	$16.7 \pm 0.7$					
	20.0	$10.9 \pm 0.3$					
35.0	5.0	$223 \pm 8$	$594 \pm 19$	$8.0 \pm 0.4$	10	266	$0.56 \pm 0.21$
	10.0	$88 \pm 5$					
	15.0	$51 \pm 2$					
	20.0	$36 \pm 2$					
$\Delta H^\ddagger/\text{kJ mol}^{-1}$				$34 \pm 7$	$48 \pm 4$	$87 \pm 2$	
$\Delta S^\ddagger/\text{J K}^{-1} \text{mol}^{-1}$				$-61 \pm 22$	$-33 \pm 13$	$82 \pm 6$	


<sup>a</sup> [Complex]<sub>T</sub> =  $1.0 \times 10^{-3}$ ,  $I = 1.0 \text{ mol dm}^{-3}$ ;  $\lambda = 400 \text{ nm}$ . <sup>b</sup> From average values of the intercepts of  $k_{\text{obs}}$  vs.  $[\text{Fe}^{3+}]_T$  plots at different acidities [see equation (10)].



**Scheme 4**  $R = (\text{NH}_3)_5\text{Co}^{3+}$ ,  $\text{nta} = \text{N}(\text{CH}_2\text{CO}_2)_2$ <sup>3-</sup>,  $M^{3+} = \text{Fe}^{3+}$  or  $\text{Al}^{3+}$

and  $k_3$  could not be determined accurately but have been estimated as follows. It is reasonable to assume that  $k_{-2} = k_{-3}$ , since the transition states for these dissociation paths should not differ greatly. Any difference in the distribution of hydrolysis products of the binuclear species is expected to arise, at a given acidity, by very rapid proton transfers governed by  $K_1$  and  $K_h$ . Considering the interrelationships of the various equilibria in Scheme 4 the assumption  $k_2 K_h / k_3 K_1 = k_{-2} / k_{-3}$ , can be made for which  $k_2 K_h = k_3 K_1$ . Thus, the values of  $k_2$  and  $k_3$  are estimated. The various rate parameters for  $\text{Fe}^{3+}$  complexation are collected in Table 6, with the corresponding rate parameters for  $\text{Al}^{3+}$  in Table 5.

The kinetic results for the formation of  $\text{Fe}^{3+}$  complexes with some related carboxylate and aminocarboxylate ligands are summarised in Table 7. It is interesting that the formation rate constants by various pathways in the nta system are comparable (within a factor of 10) with analogous rate



**Fig. 3** Plots of  $k_r[\text{H}^+]$  versus  $[\text{H}^+]^{-1}$  for the formation of  $[(\text{NH}_3)_5\text{Co}(\text{nta})\text{Fe}]^{3+}$  at 15 (a), 20 (b), 25 (c) and 35 °C (d)

constants for other carboxylate systems.<sup>2,6,29-35</sup> Assuming an  $I_d$  mechanism and using  $k = k^* K_{\text{os}}$ <sup>36</sup> where  $K_{\text{os}}$  is the outer-sphere association constant as defined earlier and  $k^*$  is the water-exchange rate constant for the ion pair, the values of  $k^*$  for  $\text{Fe}^{3+}$  or  $\text{FeOH}^{2+}$  complexation with various 0 and -1 charged ligands ( $K_{\text{os}} = 0.15\text{--}2.1 \text{ mol dm}^{-3}$ ) can be calculated; and fall in the ranges  $1.2\text{--}1500 \text{ s}^{-1}$  and  $(0.1\text{--}5.0) \times 10^4 \text{ s}^{-1}$ , respectively.<sup>37,38</sup> The value of  $K_{\text{os}}$  for  $\text{Fe}^{3+}$  or  $\text{FeOH}^{2+}$  with neutral ligands is reported<sup>39</sup> to be  $0.3 \text{ dm}^3 \text{ mol}^{-1}$  at  $25^\circ\text{C}$  ( $I = 1.0 \text{ mol dm}^{-3}$ ). Even lower values of  $K_{\text{os}}$  are expected for  $3+2+$ ,  $2+2+$  and  $2+1+$  charge associations. A value of  $K_{\text{os}} = 0.07 \text{ dm}^3 \text{ mol}^{-1}$  for  $2+1+$  charge association (as estimated for bivalent metal ions,  $25^\circ\text{C}$ ,  $I = 0.3 \text{ mol dm}^{-3}$ ) yielded  $k_4^* = 0.88 \times 10^5 \text{ s}^{-1}$  (at  $25^\circ\text{C}$ ). This is close to the value of the water-exchange rate constant of  $\text{FeOH}^{2+}$  ( $1.4 \times 10^5 \text{ s}^{-1}$ ).<sup>37,40</sup> By an identical calculation, using  $K_{\text{os}} \approx 0.05 \text{ dm}^3 \text{ mol}^{-1}$  for  $3+1+$  and  $2+2+$  associations,  $k_2^*$  and  $k_3^*$  were found to be  $\approx 1.13 \times 10^4 \text{ s}^{-1}$  and  $\approx 1.7 \times 10^3 \text{ s}^{-1}$  ( $25^\circ\text{C}$ ,  $I = 1.0 \text{ mol dm}^{-3}$ ), respectively. Comparison of these values with the water-exchange rate constants of  $[\text{Fe}(\text{OH}_2)_6]^{3+}$  and  $[\text{Fe}(\text{OH}_2)_5\text{OH}]^{2+}$  [ $k_{\text{ex}}(\text{Fe}^{3+}) = 1.6 \times 10^2 \text{ s}^{-1}$ ,

**Table 7** Comparison of rate parameters for the formation of  $\text{Fe}^{\text{III}}$  complexes with some carboxylate ligands at 25 °C; 0.5 ≤ pH ≤ 2.5\*

Reacting species	$I/\text{mol dm}^{-3}$	$k/\text{dm}^3 \text{ mol}^{-1} \text{ s}^{-1}$	Ref.
$\text{Fe}^{3+} + [\text{Co}(\text{NH}_3)_5(\text{C}_2\text{O}_4)]^+$	1.0	870	2
$\text{FeOH}^{2+} + [\text{Co}(\text{NH}_3)_5(\text{C}_2\text{O}_4)]^+$	1.0	$3.7 \times 10^4$	2
$\text{FeOH}^{2+} + [\text{Co}(\text{NH}_3)_5(\text{C}_2\text{O}_4)]^{2+}$	1.0	$4.6 \times 10^3$	2
$\text{FeOH}^{2+} + [\text{Co}(\text{NH}_3)_5(\text{Hmal})]^{2+}$	0.5	$1.26 \times 10^4$	6
$\text{Fe}^{3+} + [\text{Co}(\text{NH}_3)_5(\text{Hmal})]^+$	0.5	71	6
$\text{Fe}^{3+} + [\text{Co}(\text{NH}_3)_5(\text{Hnta})]^+$	1.0	85	This work
$\text{FeOH}^{2+} + [\text{Co}(\text{NH}_3)_5(\text{H}_2\text{nta})]^{2+}$	1.0	$5.7 \times 10^2$	This work
$\text{FeOH}^{2+} + [\text{Co}(\text{NH}_3)_5(\text{Hnta})]^+$	1.0	$6.2 \times 10^3$	This work
$\text{FeOH}^{2+} + \text{H}_2\text{nta}^-$	0.5	$1.0 \times 10^5$	29
		$5.6 \times 10^4$	30
$\text{FeOH}^{2+} + \text{H}_3\text{nta}$	0.5	$1.5 \times 10^4$	29
$\text{FeOH}^{2+} + \text{H}_2\text{idta}$	—	$2.5 \times 10^3$	29
$\text{FeOH}^{2+} + \text{Hida}^-$	—	$8.8 \times 10^3$	29
$\text{FeOH}^{2+} + \text{MeCO}_2\text{H}$	0.5	$2.8 \times 10^3$	31
$\text{Fe}^{3+} + \text{MeCO}_2\text{H}$	0.5	27	31
$\text{FeOH}^{2+} + \text{CH}_2\text{ClCO}_2\text{H}$	1.0	$8.3 \times 10^3$	32
$\text{Fe}^{3+} + \text{HC}_2\text{O}_4^-$	1.0	$8.6 \times 10^2$	33
$\text{FeOH}^{2+} + \text{HC}_2\text{O}_4^-$	1.0	$2.0 \times 10^4$	33
$\text{Fe}^{3+} + \text{H}_2\text{pydca}$	1.0	34	34
$\text{FeOH}^{2+} + \text{H}_2\text{pydca}$	1.0	$1.5 \times 10^4$	34
$\text{Fe}^{3+} + \text{H}_3\text{L}^{1+}$	0.1	180	35
$\text{FeOH}^{2+} + \text{H}_3\text{L}^{1+}$	0.1	$1.27 \times 10^3$	35
$\text{Fe}^{3+} + \text{H}_3\text{L}^{2+}$	0.1	108	35
$\text{FeOH}^{2+} + \text{H}_3\text{L}^{2+}$	0.1	$5.28 \times 10^3$	35
$\text{Fe}^{3+} + \text{H}_3\text{L}^3$	0.1	52	35
$\text{FeOH}^{2+} + \text{H}_4\text{L}^{3+}$	0.1	$1.31 \times 10^3$	35

\*  $\text{H}_2\text{mal}$  = malonic acid,  $\text{H}_2\text{idta}$  = iminodiacetic acid,  $\text{H}_2\text{pydca}$  = pyridine-2,6-dicarboxylic acid,  $\text{H}_2\text{L}^1$  = R-L-Phe,  $\text{H}_2\text{L}^2$  = R-L-Ala,  $\text{H}_3\text{L}^3$  = R-L-Asp [R = 3-hydroxy(5-hydroxymethyl)2-methylpyridine-4-methylene].

**Table 8** Comparison of rate parameters for the formation of  $\text{Al}^{\text{III}}$  complexes with some related carboxylate ligands<sup>a</sup>

Reacting species	$I/\text{mol dm}^{-3}$	$T/\text{°C}$	$k_f/\text{dm}^3 \text{ mol}^{-1} \text{ s}^{-1}$	$k_r/\text{s}^{-1}$	Refs.
$\text{Al}^{3+} + [\text{Co}(\text{NH}_3)_5(\text{C}_2\text{O}_4)]^+$	1.0	30	$2.47 \pm 0.05$	$5.0 \times 10^{-2}$	9
$\text{AlOH}^{2+} + [\text{Co}(\text{NH}_3)_5(\text{C}_2\text{O}_4)]^+$	1.0	30	840	—	9
$\text{Al}^{3+} + [\text{Co}(\text{NH}_3)_5(\text{Hnta})]^+$	0.3	25	$6.2 \times 10^{-2}$	—	This work
$\text{AlOH}^{2+} + [\text{Co}(\text{NH}_3)_5(\text{H}_2\text{nta})]^{2+}$	0.3	25	48	—	This work
$\text{AlOH}^{2+} + [\text{Co}(\text{NH}_3)_5(\text{Hnta})]^+$	0.3	25	$85 \pm 10$	$1.3 \times 10^{-2}$	This work
$\text{Al}^{3+} + [\text{Co}(\text{NH}_3)_5(\text{Hsal})]^{2+}$	0.1	30	$0.16 \pm 0.06$	$5.9 \times 10^{-3}$	41
$\text{AlOH}^{2+} + [\text{Co}(\text{NH}_3)_5(\text{Hsal})]^{2+}$	0.1	30	$67 \pm 15$	—	41
$\text{Al}^{3+} + [\text{Co}(\text{NH}_3)_5(\text{Hnsal})]^{2+}$	0.1	30	$0.10 \pm 0.10$	$3.84 \times 10^{-3}$	41
$\text{AlOH}^{2+} + [\text{Co}(\text{NH}_3)_5(\text{Hnsal})]^{2+}$	0.1	30	$8 \pm 2$	—	41
$\text{Al}^{3+} + \text{Hsal}^-$	0.1	30	0.78	—	42
$\text{AlOH}^{2+} + \text{Hsal}^-$	0.1	25	$1.02 \times 10^3$	—	42

<sup>a</sup>  $\text{Hsal}$  = Salicylic acid,  $\text{Hnsal}$  = 3-nitrosalicylic acid. <sup>b</sup> Spontaneous dissociation rate of  $\text{Al}^{\text{III}}$  complexes.

$k_{\text{ex}}(\text{FeOH}^{2+}) = 1.4 \times 10^5 \text{ s}^{-1}$ , 25 °C<sup>37,40</sup> suggests an associative ( $I_a$ ) mechanism for  $\text{Fe}^{3+}$  ( $k_3$  path) and an  $I_d$  mechanism for  $\text{FeOH}^{2+}$  ( $k_2$  or  $k_4$  paths). This is further supported by the fact that the activation parameters  $\Delta H^\ddagger$  and  $\Delta S^\ddagger$  (see Table 6) for the substitution reactions of  $\text{Fe}^{3+}$  and  $\text{FeOH}^{2+}$  are not substantially different from those for the water-exchange reaction of  $[\text{Fe}(\text{OH}_2)_6]^{3+}$  ( $\Delta H^\ddagger = 64 \text{ kJ mol}^{-1}$ ,  $\Delta S^\ddagger = 12 \text{ J K}^{-1} \text{ mol}^{-1}$ ) and of  $[\text{Fe}(\text{OH}_2)_5\text{OH}]^{2+}$  ( $\Delta H^\ddagger = 42.4 \text{ kJ mol}^{-1}$ ,  $\Delta S^\ddagger = 5.3 \text{ J K}^{-1} \text{ mol}^{-1}$ ).<sup>37</sup>

For  $\text{Al}^{\text{III}}$  the plots of  $k_{\text{obs}}$  versus  $[\text{Al}^{3+}]\{[\text{H}^+]/([\text{H}^+] + K_1)\}$  yielded [see equation (11)]  $k_f$  and  $k_r$ . The values of  $k_2$ ,  $k_3$  and  $k_4$  (see Table 5) were calculated by an identical procedure to that adopted for the  $\text{Fe}^{\text{III}}$  system. A comparison of the values of rate constants for the formation of  $\text{Al}^{\text{III}}$  complexes with some carboxylate ligands is given in Table 8. The magnitudes of the rate constants for the formation of  $[(\text{NH}_3)_5\text{Co}(\text{nta})\text{Al}]^{3+}$  via the reaction of  $\text{Al}^{3+}$  or  $\text{AlOH}^{2+}$  with the nta cobalt(III) substrate are comparable to those obtained for the formation of  $[(\text{NH}_3)_5\text{Co}(\text{sal})\text{Al}]^{4+}$  (ref. 41) but are lower than those for  $[(\text{NH}_3)_5\text{Co}(\text{C}_2\text{O}_4)\text{Al}]^{4+}$ .<sup>9</sup> The values of  $k_3$  (see Table 5) are smaller than the most recent data of water-exchange rate constant of  $[\text{Al}(\text{OH}_2)_6]^{3+}$  ( $k_{\text{ex}} = 16 \text{ s}^{-1}$  at 25 °C,<sup>42</sup>  $k_3/K_{\text{os}} =$

$1.2 \text{ s}^{-1}$  using  $K_{\text{os}} = 0.05 \text{ dm}^3 \text{ mol}^{-1}$ ). The fact that  $k_1 < k_{\text{ex}}$  eliminates the simple associative mechanism and supports an  $I_d$  mechanism. Similar conclusions were drawn previously for the complexation of  $\text{Al}^{\text{III}}$  with a number of co-ordinated carboxylate systems.<sup>43</sup> Much faster substitution reactions for  $[\text{Al}(\text{OH}_2)_5\text{OH}]^{2+}$ , as observed in the present work, are consistent with an  $I_d$  mechanism.


The binuclear species  $[(\text{NH}_3)_5\text{Co}(\text{nta})\text{Al}]^{3+}$  undergoes spontaneous dissociation at a rate comparable with that of  $[(\text{NH}_3)_5\text{Co}(\text{sal})\text{Al}]^{4+}$  or  $[(\text{NH}_3)_5\text{Co}(\text{C}_2\text{O}_4)\text{Al}]^{4+}$  (see Table 8). Since  $\text{Al}^{3+}$  is chelated by the bound salicylate and oxalate moieties, it is presumed that the nta moiety chelates the metal ion in at least a tridentate mode. However, complexation of  $[\text{Co}(\text{NH}_3)_5(\text{H}_2\text{nta})]^{2+}$  with trivalent metal ions, in contrast to that for bivalent metals, does not involve chelate-ring closure as the rate limiting step (step 2, Scheme 5). A similar conclusion was also drawn by Mentasti<sup>38</sup> for free-ligand complexation reactions.

The rate data for the acid-catalysed dissociation of  $[(\text{NH}_3)_5\text{Co}(\text{nta})\text{Fe}]^{3+}$  are available as supplementary data. The observed rate constants were corrected for the formation reaction utilising the values ( $k_2K_h + k_3K_1$ ),  $k_4K_hK_1$  and  $[\text{H}^+]$ .

**Table 9** Comparison of rate and activation parameters for the dissociation of some  $\text{Fe}^{\text{III}}$  complexes at 25 °C,  $I = 1.0 \text{ mol dm}^{-3}$ <sup>a</sup>


$\text{Fe}^{\text{III}}$ complex	$k_s^b/\text{s}^{-1}$	$\Delta H^\ddagger/\text{kJ mol}^{-1}$	$\Delta S^\ddagger/\text{J K}^{-1} \text{ mol}^{-1}$	$k_a^b/\text{dm}^3 \text{ mol}^{-1} \text{ s}^{-1}$	$\Delta H^\ddagger/\text{kJ mol}^{-1}$	$\Delta S^\ddagger/\text{J K}^{-1} \text{ mol}^{-1}$	Refs.
$[(\text{NH}_3)_5\text{Co}(\text{sal})\text{Fe}]^{4+}$	0.23	63 ± 6	-50 ± 17	0.05	72 ± 12	-29 ± 38	3
<i>cis</i> -[(en) <sub>2</sub> (NH <sub>3</sub> ) <sub>2</sub> Co(sal)Fe] <sup>4+</sup>	0.30	63 ± 2	-46 ± 8	0.09	64 ± 10	-50 ± 34	3
$\alpha\beta\text{S}-[(\text{en})_2\text{Co}(\text{sal})\text{Fe}]^{4+}$	0.72	62 ± 2	-42 ± 8	0.29	70 ± 4	-21 ± 13	3
$[(\text{NH}_3)_5\text{Co}(\text{C}_2\text{O}_4)_2\text{Fe}]^{4+}$	negligible			0.50	44 ± 3	-99 ± 8	2
$[(\text{NH}_3)_5\text{Co}(\text{nta})\text{Fe}]^{3+}$	0.24	61 ± 4	-31 ± 13	0.66	64 ± 4	-28 ± 14	This work
$[(\text{NH}_3)_5\text{Co}(\text{mal})\text{Fe}]^{4+}$	5.11	—	—	—	—	—	6
$[\text{Fe}(\text{pydca})]^+$	—	—	—	0.25	—	—	34
$[\text{Fe}(\text{O}_2\text{CCH}_2\text{Cl})]^{2+}$	20	—	—	—	—	—	32

<sup>a</sup> en = Ethylenediamine, tetren = tetraethylenepentamine. <sup>b</sup>  $k_s$  and  $k_a$  represent the spontaneous and acid-catalysed rate constants, respectively.

**Scheme 5**  $\text{M}^{3+} = \text{Fe}^{3+}$  or  $\text{Al}^{3+}$ 

The linear dependence of  $k_r (= k_{\text{obs}} - k_f[\text{M}^{3+}])$  with  $[\text{H}^+]$  (see Fig. 4) suggests that the last term in equation (7) is insignificant and can be neglected. The values of  $k_{-4}$  and  $(k_{-2} + k_{-3})$  derived from the intercepts and slopes, respectively are also available as supplementary data. A comparison of rate and activation parameters for the dissociation of analogous  $\text{Fe}^{\text{III}}$  complexes are given in Table 9. The rate constant for the spontaneous dissociation,  $k_4$  ( $= k_s$  in Table 9) at all temperatures compares satisfactorily with the value of  $k_r(\text{av.})$  obtained from the formation study. It is also evident from Table 9 that the value of  $k_s$  for the nta complex is comparable with that obtained for  $[(\text{NH}_3)_5\text{Co}(\text{sal})\text{Fe}]^{4+}$  but significantly lower than that of  $[\text{Fe}(\text{O}_2\text{CCH}_2\text{Cl})]^{2+}$ .<sup>38</sup> The relatively slow dissociation rate of the binuclear species may be taken as evidence that the  $\text{Fe}^{\text{III}}$  ion in  $[(\text{NH}_3)_5\text{Co}(\text{nta})\text{Fe}]^{3+}$  is chelated by the nta moiety in at least a tridentate co-ordination mode (see Scheme 5). A similar conclusion was reached by Cannon and Gardiner<sup>1</sup> in the study of  $\text{Fe}^{\text{III}}$ -catalysed aquation of  $[\text{Co}(\text{NH}_3)_5(\text{H}_2\text{nta})]^{2+}$ . The rate and activation parameters for the acid-catalysed dissociation of  $[(\text{NH}_3)_5\text{Co}(\text{nta})\text{Fe}]^{3+}$  are also comparable with the values reported for related binuclear species (see Table 9). The activation parameters also fall in the range of  $\Delta H^\ddagger$  ( $= 38-63 \text{ kJ mol}^{-1}$ ) and  $\Delta S^\ddagger$  ( $= -42$  to  $-126 \text{ J K}^{-1} \text{ mol}^{-1}$ ) values reported<sup>44</sup> for the dissociation of a series of  $[\text{FeL}]^{n+}$  ( $\text{L} = \text{F}^-, \text{N}_3^-, \text{Cl}^-, \text{SCN}^-, \text{SO}_4^{2-}$ ) complexes.

In conclusion, the difference in the rate-determining steps for the complexation of bivalent and trivalent metal ions with

**Fig. 4** Plots of  $k_r (= k_{\text{obs}} - k_f[\text{Fe}^{3+}])$  versus  $[\text{H}^+]$  for the acid-catalysed dissociation of  $[(\text{NH}_3)_5\text{Co}(\text{nta})\text{Fe}]^{3+}$  at 15 (a), 20 (b), 25 (c) and 35 °C (d)

$[\text{Co}(\text{NH}_3)_5(\text{H}_2\text{nta})]^{2+}$  may be explained as follows. The relatively greater coulombic repulsion between the  $\text{NH}^+$  centre and the carboxylate-bound  $\text{M}^{3+}$  ion (as compared to  $\text{M}^{2+}$ ) or favourable acid-base reaction of  $\text{MOH}^{2+}$  with the  $\text{NH}^+$  site may result in the enhanced rate of proton transfer from the  $\text{NH}^+$  centre of the binuclear species intermediate for  $\text{M}^{3+}$  ions [see Scheme 5 and equation (5)]. This may be responsible for proton transfer being equilibrium controlled for  $\text{M}^{3+}$  ions, while it is rate controlling for the  $\text{M}^{2+}$  ions (see Schemes 3 and 5). Accordingly general base catalysis was observed only for the complexation of the  $\text{M}^{2+}$  ions.

### Acknowledgements

A. C. D. is thankful to Department of Science and Technology, New Delhi for financial support to purchase a stopped flow spectrophotometer and R. D. is thankful for the award of a Junior Research Fellowship.

### References

- 1 R. D. Cannon and J. Gardiner, *Inorg. Chem.*, 1974, **13**, 390.
- 2 A. C. Dash and G. M. Harris, *Inorg. Chem.*, 1982, **21**, 1265.

3 A. C. Dash and G. M. Harris, *Inorg. Chem.*, 1982, **21**, 2336.

4 A. C. Dash, R. K. Nanda and A. N. Acharya, *Indian J. Chem., Sect. A*, 1991, **30**, 769.

5 A. C. Dash and J. Pradhan, *Int. J. Chem. Kinet.*, 1992, **24**, 155.

6 N. N. Das and R. K. Nanda, *Indian J. Chem., Sect. A*, 1991, **30**, 125.

7 A. C. Dash, R. K. Nanda and A. N. Acharya, *J. Chem. Soc., Dalton Trans.*, 1993, 1023.

8 R. Das and N. N. Das, *Transition Met. Chem.*, in the press.

9 P. Mohanty and A. C. Dash, *Transition Met. Chem.*, 1995, **20**, 153.

10 D. H. Bush and J. C. Bailar jun., *J. Am. Chem. Soc.*, 1953, **75**, 4574.

11 A. C. Dash and R. K. Nanda, *Inorg. Chem.*, 1974, **13**, 655.

12 H. M. Irving and H. S. Rossotti, *J. Chem. Soc.*, 1954, 2904.

13 R. G. Wilkins, *Kinetics and Mechanism of Reactions of Transition Metal Complexes*, 2nd edn., VCH, New York, 1991, pp. 202, Table 4.1.

14 Y. Ducommun, K. E. Newman and A. E. Merbach, *Inorg. Chem.*, 1980, **19**, 3697.

15 Y. Ducommun, W. L. Carl and A. E. Merbach, *Inorg. Chem.*, 1979, **18**, 2754.

16 A. E. Merbach, *Pure Appl. Chem.*, 1987, **59**, 161.

17 R. B. Jordan, *Reaction Mechanism of Inorganic and Organometallic Systems*, Oxford University Press, Oxford, 1991.

18 D. H. Powell, L. Helm and A. E. Merbach, *J. Chem. Phys.*, 1991, **95**, 9258.

19 R. M. Fuoss, *J. Am. Chem. Soc.*, 1958, **80**, 5059.

20 T. J. Bydalek and M. L. Blomster, *Inorg. Chem.*, 1964, **3**, 667.

21 N. Tanaka and M. Kimura, *Bull. Chem. Soc. Jpn.*, 1967, **40**, 2100.

22 N. Tanaka and M. Kimura, *Bull. Chem. Soc. Jpn.*, 1968, **41**, 2375.

23 J. Maguire, *Can. J. Chem.*, 1974, **52**, 4106.

24 D. L. Rabenstein, *J. Am. Chem. Soc.*, 1971, **93**, 2869.

25 J. C. Cassatt and R. G. Wilkins, *J. Am. Chem. Soc.*, 1968, **90**, 6045.

26 D. B. Rorabacher, T. S. Turan, J. Defever and W. G. Nickels, *Inorg. Chem.*, 1969, **8**, 1498.

27 R. M. Milburn, *J. Am. Chem. Soc.*, 1957, **79**, 537.

28 R. K. Schofield and A. W. Taylor, *J. Chem. Soc.*, 1954, 4445.

29 E. Mentasti, E. Pelizzetti and G. Saini, *Gazz. Chim. Ital.*, 1974, **104**, 201.

30 S. Funahashi, A. Adachi and M. Tanaka, *Bull. Chem. Soc. Jpn.*, 1973, **46**, 479.

31 R. N. Pandey and N. W. Smith, *Can. J. Chem.*, 1972, **50**, 194.

32 B. Perlmutter-Hayman and E. Tapuhi, *J. Coord. Chem.*, 1976, **6**, 31.

33 E. G. Moohead and N. Sutin, *Inorg. Chem.*, 1966, **5**, 1866.

34 K. Bridger, R. C. Patel and E. Matijevic, *Polyhedron*, 1982, **1**, 269.

35 T. Ozawa, K. Jitsukawa, H. Masuda and H. Einaga, *Ber. Bunsenges. Phys. Chem.*, 1994, **98**, 66.

36 M. Eigen and R. G. Wilkins, *Adv. Chem. Ser.*, 1965, **49**, 55.

37 M. Grant and R. B. Jordan, *Inorg. Chem.*, 1981, **20**, 51.

38 E. Mentasti, *Inorg. Chem.*, 1979, **18**, 1512.

39 K. Ishihara, S. Funahashi and M. Tanaka, *Inorg. Chem.*, 1983, **22**, 194.

40 H. W. Dodgen, G. Liu and J. P. Hunt, *Inorg. Chem.*, 1981, **20**, 1002.

41 A. C. Dash, *Inorg. Chem.*, 1983, **22**, 837.

42 F. Secco and M. Venturini, *Inorg. Chem.*, 1975, **14**, 1978.

43 D. W. Margerum, G. R. Cayley, D. C. Weatherburn and G. K. Pagenkopf, *ACS Monogr.*, 1978, **174(2)**, 5.

44 F. P. Cavasino, *J. Phys. Chem.*, 1968, **72**, 1378.

Received 3rd April 1995; Paper 5/02110C