

THE PERCEPTION OF LIGHT AND COLOUR AND THE PHYSIOLOGY OF VISION

Part VII. General Summary

BY SIR C. V. RAMAN

(*Memoir No. 125 of the Raman Research Institute, Bangalore-6*)

Received December 10, 1960

1. OBSERVATIONS OF THE RETINA

A SIMPLE but extremely powerful technique has been devised which enables an observer without any instrumental aid to see his own retina and study its functioning in varied circumstances. The observations made by this method have enabled important conclusions to be arrived at regarding the constitution of the retina and the mechanism by which light and colour are perceived. The observer sits facing a brightly lit white screen and views it through an appropriate colour filter held in front of his eye. After a sufficient interval of time, he fixes his vision on some particular point on the screen and then removes the filter. An enormously magnified picture of the retina then appears on the screen, the nature of which depends very much on the particular colour filter used. The explanation of the phenomenon is that the rays of the spectrum which in the first instance are absorbed by the filter, suddenly impinge on the retina when the filter is removed, and excite localised sensations over its different areas. These sensations project themselves on the observing screen as an enlarged image of the retina.

By correlating the absorption spectra of the filters used with the pictures of the retina perceived by the observer, it has been ascertained that the retina contains three visual pigments whose absorption spectra lie in different regions of the spectrum: Pigment A has an absorption lying in the spectral range 4000 to 5000 Å. Pigment B exerts an extremely powerful absorption in the wavelength range between 5000 and 6000 Å, while pigment C exerts a moderately powerful absorption in the region from 6000 to 7000 Å, but its absorption also extends into and partly overlaps the region covered by pigment B.

2. THE ROLE OF THE VISUAL PIGMENTS

The nature and properties of light when correlated with the observed facts concerning the perception of light and colour enable us to determine

the role played by the visual pigments in the retina. Each different monochromatic light in the spectrum represents radiation characterised by a different energy-quantum; the magnitude of the quantum increases progressively from the red to the violet end of the spectrum. It is a fact also that the colours observed in the spectrum change progressively and continuously from one end of it to the other. Some 250 different colours can be perceived over the whole range. A change of only one half of one per cent. in the magnitude of the light-quantum incident on and absorbed by the retinal pigments is usually sufficient to make an observable change of perceived colour, while in some parts of the spectrum a much smaller difference is thus detectable.

Monochromatic radiation is the fundamental entity with which the physicist is concerned in optics and spectroscopy. It follows from what has been stated that the fundamental visual sensations are also those excited by monochromatic radiation. The observed precision of colour perception would be inexplicable except on the hypothesis that the function of the visual pigments in the retina is to receive, absorb and then to pass on the absorbed energy-quanta to the central parts of the visual mechanism without any addition or subtraction, themselves returning to their original energy-states.

3. IDENTIFICATION OF THE VISUAL PIGMENTS

The identification of the visual pigments presents no particular difficulties. In day-light vision, there is a highly pronounced maximum of luminosity in the spectrum at about 5600 Å, on either side of which the brightness falls off rapidly. It follows that pigment B of which the absorption lies in the green plays the major role in human vision, a role somewhat analogous to that which the absorption in the red by chlorophyll plays in the photosynthesis by green leaves. We can, therefore, unhesitatingly identify our pigment B with ferroheme which exhibits a powerful absorption of light in the green sector located at the same position as the maximum of visual luminosity in the spectrum.

Oxidation-reduction mechanisms play a fundamental role in the chemistry of the living structures of aerobic organisms. The recognition of ferroheme as the principal visual pigment thus automatically involves the identification of pigment C which appears in the retina in close association with pigment B as ferriheme. It is known that the absorption of light by ferriheme is weaker than that of ferroheme, but extends much further towards longer wavelengths and is indeed sensible up to the extreme red end of the spectrum. These are the properties needed for the visual pigment which functions in that region.

The pigment A which absorbs light between 4000 and 5000 Å and is the mediator of vision in the blue and violet sectors of the spectrum can be

none other than the carotenoid pigment xanthophyll which gives the characteristic golden yellow colour to the yoke of the common hen's egg. Xanthophyll finds its way into the human body through the consumption of food products containing it, and its presence in the retina is therefore not a matter for surprise. Indeed, it is the same yellow pigment which led to the anatomical name of *macula lutea* being given to the physiologically most important area in the retina.

4. COLOUR AND LUMINOSITY IN THE SPECTRUM

The known features of the absorption spectra of the visual pigments enable a satisfactory explanation to be given of the distinctive features noticeable in the different sectors of the spectrum. The very steep rise in the absorption by xanthophyll which appears in the wavelength range around 4900 Å is responsible for the rapid change in the colour of the spectrum from green to blue and the very low value of the limen of wavelength alteration needed for observable change of colour appearing in that region. The similar but less striking fall of the limen in the region around 4400 Å where the colour of the spectrum changes from blue to violet is likewise attributable to the rapid fall of the absorption by xanthophyll with diminishing wavelength appearing in that region.

The overlap in the absorption spectra of ferroheme and ferriheme between 625 $m\mu$ and 566 $m\mu$ gives rise to the appearance of yellow and orange in the spectrum as an interpolation between the green and red sectors, the yellow where the absorptions of the two pigments are of comparable strength, and the orange where the absorption by ferriheme is stronger than by ferroheme. The steep fall in the absorption by ferroheme in the same region is responsible for the limen of wavelength change needed for an observable difference of colour reaching very low values in the region around 590 $m\mu$.

The progressive fall in the luminosity of the spectrum and the increasing limen of wavelength change for a perceptible colour difference manifested near the extreme violet and red ends of the spectrum appear as consequences respectively of the diminishing absorption and the diminishing slope of the absorption curves of xanthophyll and of ferriheme in those two regions.

Some remarkable effects are observed when a colour filter which transmits only the blue-violet part of the spectrum is held in front of the eye and a polaroid is placed and alternately taken out and put in before the filter. These effects cease to be observable when the illumination of the field under observation is diminished so as to fall below the photopic level. They afford

a conclusive demonstration that xanthophyll is the visual pigment which enables us to perceive the blue and violet colours of the spectrum.

5. NON-HOMOGENEOUS LIGHT

The visual sensations produced by heterogeneous light result from the synthesis by the centre of the sensations excited by the monochromatic radiations of which it is composed. The results of the synthesis may be either a chromatic or an achromatic sensation. Chromatic sensations arise when only two of the three visual pigments function, as for example when the radiations from the extreme red and violet ends of the spectrum are superposed, giving rise to the purples, or when the superposed radiations lie in the spectral range where xanthophyll has no absorption, *viz.*, in the red and green sectors; the resultant sensation is then a pure spectral colour.

The achromatic sensation arises when all the three visual pigments function in appropriate strengths. This enables a satisfactory explanation to be given of the fact that in a whole series of cases, the superposition of only two monochromatic radiations of appropriately chosen wavelengths results in the complete abolition of colour. In general, the sensations excited by heterogeneous light are a mixture of the chromatic and achromatic sensations which may be regarded as independent effects. The fact that all observable colours may be regarded as a superposition in appropriate proportions of white light and a pure spectral colour (including the pure purples) thus receives a satisfactory explanation.

The results obtained in experiments in which a pure spectral colour is sought to be reproduced by the superposition of other spectral colours may be interpreted in the same manner. The so-called spectral chromaticity coefficients determined in such experiments, when they have positive values, exhibit a parallelism with the absorption strength of the visual pigments which function in the respective spectral regions. *Per contra*, the appearance of negative values of the coefficients indicates that achromatic sensations are produced.

6. DEFECTIVE COLOUR VISION

The existence of both ferroheme and ferriheme as visual pigments in the retina presupposes that there is a biochemical mechanism which determines the proportions in which they are normally present. Any deviations of the mechanism from normality would result in ferriheme being either totally absent or else being present much in excess. There would also be intermediate cases. The existence of four types of inheritable defect in colour

vision would thus be explicable in terms of the biochemical mechanism which determines the ferroheme-ferriheme ratio in the retina. These are respectively the protanopic, protanomalous, deuteranomalous and deuteranopic types of colour vision. The features of these different types of defects are readily predictable and the results thus deduced are in agreement with the observed facts. Tritanopia is likewise explicable as due to the complete absence of xanthophyll from the retina.

7. REFERECENS

The literature on colour vision and related topics is very voluminous. The under-mentioned books were found very useful by the writer. Besides containing extensive bibliographies, they gave clear accounts of the present state of knowledge in their respective fields and factual information of value:—

- Bouma, P. J. .. *Physical Aspects of Colour*, Philips, Eindhoven, 1947.
 Yves Le Grand .. *Light, Colour and Vision*, Chapman and Hall, London, 1957.
 Goodwin, T. W. .. *The Comparative Biochemistry of the Carotenoids*, Chapman and Hall, London, 1952.
 Lemberg, R. and Legge, J. W. .. *Hematin Compounds and Bile Pigments*, Interscience, New York, 1949.

The following is a selected list of papers containing factual information of importance in relation to the subject-matter of the present memoir.

Luminous Efficiency in the Spectrum

- Walters and Wright .. *Proc. Roy. Soc., B*, 1943, **131**, 340.
 Hue Discrimination in the Spectrum.
 Tyndall, E. P. T. .. *J. Opt. Soc. Amer.*, 1933, **23**, 15.
 Wright and Pitt .. *Proc. Phys. Soc.*, 1934, **46**, 459.
 Haase, G. .. *Ann. Physik*, 1934, **20**, 75.
 Colour Mixing Experiments.
 Wright, W. D. .. *Jour. Sci. Inst.*, 1939, **16**, 10.
 _____ .. *Trans. Opt. Soc.*, 1929, **30**, 140.
 _____ .. *Ibid.*, 1930, **31**, 201.
 Defective Colour Vision.
 Pitt, F. H. G. .. *Proc. Roy. Soc. B*, 1945, **132**, 101.
 McKeon and Wright .. *Proc. Phy. Soc.*, 1940, **52**, 464.
 Nelson, J. H. .. *Ibid.*, 1938, **50**, 661.
 Judd, D. B. .. *Jour. Opt. Soc. Amer.*, 1943, **33**, 294.
 _____ .. *Ibid.*, 1945, **35**, 199.
 _____ .. *Jour. Res. Nat. Bur. St.*, 1948, **41**, 247.
 Wright, W. D. .. *Jour. Opt. Soc. Amer.*, 1952, **42**, 509.

Supplement to the

"Proceedings of the Indian Academy of Sciences"

INDEX TO VOL. LII—(A)

AUTHORS' INDEX

- Ambiye, Kumari, S. Y., Decay scheme of EU¹⁵⁶, 157.
 Bhagwat, S. D., Joshi,
 M. C., Sharma, R. P.
 and Subbarao, B. N.
- Bagavant, G. .. See Banerjee and Bagavant.
- Banerjee, D. K. and Isomerisations of substituted β -keto-esters, II, 165.
 Bagavant, G.
- Bhagavantam, S. and Elastic behaviour of matter under very high pressures.
 Chelam, E. V. Uniform compression, I.
- Bhagwat, S. D. .. See Ambiye and others.
- Chaturvedi, R. K. .. See Ramaiah and Chaturvedi.
- Chelam, E. V. .. See Bhagavantam and Chelam.
- Iwao, S. .. See Sudarshan and Iwao.
- Jayaraman, A. .. The luminescence spectra of diamond, 189.
- Joshi, M. C. .. See Ambiye and others.
- Kane, R. P. .. A critical appraisal of some cosmic ray data for the
 I.G.Y., 69.
- Kane, R. P. and Thakore, S. R. Latitude dependence of the daily variation of cos-
 mic ray intensity, 122.
- Krishnamurti, D. .. The Raman spectra of Polycrystalline solids, 20.
- Lalitha Ramaswamy .. See Puranik and Lalitha Ramaswamy.
- Mahadevan, C. .. See Rao and Mahadevan.
- Mathai, I. M. .. The influence of substituents on the additive reactivity
 of ethylene derivatives, VI, 182.
- Moosath, S. S. .. See Sosamma and Moosath.
- Narasimhan, R. .. On the system and engineering design of the general
 purpose electronic digital computer at T.I.F.R., 47.

- Prakash, Nirmala and Ram Behari Deviations from parallelism and equidistance in Finsler space, 209.
- Puranik, P. G. and Lalitha Ramaswamy Potential constants and calculated thermodynamic properties for diiodoethylene, 135.
- Radha, T. K. .. See Ramakrishnan and others.
- Ramachandran, G. N. .. A problem in probability related to the passage of light through a cloud of particles, 87.
.. Analysis of the X-ray diffraction pattern of helical structures, 240.
- Ramaiah, N. A. and Chaturvedi, R. K. Conductometric determination of a few physico-chemical constants of murexide, 197.
- Ramakrishnan, Alladi, Radha, T. K. and Thunga, R. On the decomposition of the Feynman propagator, 228.
- Raman, C. V. .. The perception of light and colour and the physiology of vision, I, II, III, IV, V, VI, VII, 255, 267, 281, 292, 305, 314, 324.
- Ram Behari .. See Prakash and Ram Behari.
- Rangaswami, S. and Rao, E. Venkata Chemical components of *Pulmiera alba* Linn., 173.
- Rao, B. D. Nageswara and Venkateswarlu, Putcha The nuclear magnetic resonance spectra of four spin systems, I, 109.
- Rao, B. N. Subba .. Levels in CO⁵⁹, 130.
- Rao, E. Venkata .. See Rangaswami and Rao.
- Rao, K. Kameswara and Mahadevan, C. Studies on lead-zinc mineralization in Karempudi area, Guntur District, Andhra Pradesh, 143.
- Sharma, R. P. .. See Ambiye and others.
- Sosamma Chandy and Moosath, S. S. Studies on metallic halide dioxanates, III, 58.
- Subbarao, B. N. .. See Ambiye and others.
- Sudarshan, E. C. G. and Iwao, S. A model for hyperfragments, 27.
- Swaminathan, S. .. See Viswanathan and Swaminathan.
- Thakore, S. R. .. See Kane and Thakore.
- Thunga, R. .. See Ramakrishnan and others.
- Thyagarajan, G. .. See Venkateswarlu and Thyagarajan.

- Venkataraman, S. .. Studies on collagen, VI, 80.
- Venkateswarlu, K. and Thyagarajan, G. Potential constants of ethylene oxide, ethylene oxide- d_4 and ethylene sulphide, 101.
- Venkateswarlu, Putcha .. See Rao and Venkateswarlu.
- Viswanathan, K. and Swaminathan, S. *dl*-Marindinin (dihydrokawain) and some related 6-aryl-5, 6-dihydro-4-methoxy-2-pyrones, 63.
- Viswanathan, K. S. .. The Dirac equation for many-electron systems, 35.

TITLE INDEX

- CO⁶⁰, levels in (Rao), 130.
- Collagen, studies, VI (Venkataraman), 80.
- Cosmic ray data for the I.G.Y., some, a critical appraisal of (Kane), 69.
- Cosmic ray intensity, latitude dependence of the daily variation of (Kane and Thakore), 122.
- Elastic behaviour of matter under very high pressures. Uniform compression (Bhagavantam and Chelam), 1.
- Electronic digital computer at T.I.F.R., the general purpose, on the system and engineering design of (Narasimhan), 47.
- Ethylene derivatives, additive reactivity of, the influence of substituents on, VI (Mathai), 182.
- Ethylene oxide, ethylene oxide-*d*₄ and ethylene sulphide, potential constants of (Venkateswarlu and Thyagarajan), 101.
- EU¹⁵⁶, decay scheme of (Ambiye and others), 157.
- Feynman propagator, on the decomposition of (Ramakrishnan and others), 228.
- Finsler space, deviations from parallelism and equidistance in (Prakash and Ram Behari), 209.
- Hyperfragments, a model for (Sudarshan and Iwao), 27.
- Isomerisations of substituted β -keto-esters, II (Banerjee and Bagavant), 165.
- Diiodoethylene, potential constants and calculated thermodynamic properties for (Puranik and Lalitha Ramaswamy), 135.
- Dirac equation for many-electron systems (Viswanathan), 35.
- Lead-zinc mineralization in Karempudi area, Guntur District, Andhra Pradesh, studies, 143.
- Light, passage of, through a cloud of particles, a problem in probability related to the (Ramachandran), 87.
- Luminescence spectra of diamond (Jayaraman), 189.
- dl*-Marindinin (dihydrokawain) and some related 6-aryl-5, 6-dihydro-4-methoxy-2-pyrones (Viswanathan and Swaminathan), 63.
- Metallic halide dioxanates, studies, III (Sosamma Chandy and Moosath), 58.

Murexide, conductometric determination of a few physico-chemical constants of (Ramaiah and Chaturvedi), 197.

Nuclear magnetic resonance spectra of four spin systems, I (Rao and Venkateswarlu), 109.

Physiology of vision, the perception of light and colour and the, I, II, III, IV, V, VI, VII (Raman), 255, 267, 281, 292, 305, 314, 324.

Pulmiera alba Linn., chemical components of (Rangaswami and Rao), 173.

Raman spectra of polycrystalline solids (Krishnamurti), 20.

X-ray diffraction pattern of helical structures, analysis of (Ramachandran), 240.