

Nitrogen uptake rates and new production in the northern Indian Ocean

Naveen Gandhi¹, Satya Prakash^{1,2}, R. Ramesh^{1*} and Sanjeev Kumar^{1,3}

¹Physical Research Laboratory, Navrangpura, Ahmedabad – 380 009 India

²Present address: Indian National Centre for Ocean Information Services, Hyderabad-500 055, India.

³Present address: Environmental Science Research Centre, St. Francis Xavier University, Box 5000, Antigonish, Nova Scotia, Canada

Received 9 June 2010; revised 13 October 2010

Nitrogen-15 based new production measurements initiated during the last decade in the northern Indian Ocean are summarized. Two different biogeochemical provinces in eastern Arabian Sea have been recognized during the late winter monsoon: less productive southern (non-bloom) and more productive northern (*Noctiluca scintillans* bloom) regions. The southern sector is characterized by low column N-uptake and very low *f*-ratio (~5.5 mmol N m⁻²d⁻¹ and 0.4 respectively). The *f* ratio, although low, increased progressively towards north. This increase may be the effect of more intense winter cooling towards the north. The northern part is a highly productive zone, with very high N-uptake and significantly high *f* ratio (~19 mmol N m⁻²d⁻¹ and 0.8 respectively). New production during the pre-monsoon in the Bay of Bengal is higher than that in the post-monsoon. New production for the region during the pre-monsoon averages around 5 (±4) mmol N m⁻²d⁻¹, almost twice the average value observed during the post-monsoon (2.6 mmol N m⁻²d⁻¹). Average *f* ratio for the entire region increases to 0.70 (±0.1) during pre-monsoon from 0.5 during the post-monsoon.

[Keywords: New productivity; *f* ratio; ¹⁵N tracer technique; northern Indian Ocean.]

Introduction

Nitrogenous nutrients are considered to be the major limiting factor of oceanic primary production in many regions¹. Primary production is partitioned on the basis of the nitrogen source: 'new' production² supported by the N-sources, principally, nitrate brought into the euphotic zone from the deep, and secondarily, N₂-fixation, riverine and atmospheric inputs; 'regenerated' production² supported by ammonium and urea, derived from biological processes occurring within the euphotic zone.

The ratio of new to total production is called the *f* ratio³. It represents the probability that a nitrogen atom is assimilated by phytoplankton due to new production; likewise (1-*f*) is the probability of assimilation by regenerated production. Thus, (1-*f*)/*f* provides a measure of number of times nitrogen recycles in the euphotic zone before sinking out of the system as particulate matter³. The *f* ratio also defines the strength of the biological pump, the magnitude of the rate of export of organic matter. Theoretically, *f* can vary between 0 and 1. The calculation of regenerated production², as proposed originally, did not include the uptake of urea-N. They defined the

primary production associated with ammonium alone as regenerated production and those associated with NO₃⁻ and NO₂⁻ as new production. Eppley and Peterson³ qualitatively showed that in some cases the uptake of urea and other organic-N contributes up to ~30% of the total nitrogen uptake and therefore non-inclusion of urea uptake overestimates the *f* ratio. Available data for the northern Indian Ocean⁴ was used to check the effect of non inclusion of urea uptake on the *f* ratio (Fig. 1). In general *f* ratio appears to be overestimated by up to 20% (i.e. 40% is estimated as 60%).

The total and new productions are often linearly correlated and the relation can be expressed in terms of a straight line⁵.

$$\rho_N = \rho_T (f_M) + \rho_M$$

where ρ_N and ρ_T are the new and total production, the intercept, ρ_M , is the minimum possible regenerated production in the absence of extraneous nitrate supply and the slope, f_M , gives the upper bound of the *f* ratio for a particular region. The significance of this linear relationship lies in the quantification of new production from satellite data on total production⁵ (e.g. derived from ocean color monitor).

*Author for correspondence
Email : rramesh@prl.res.in




Fig 1—Comparison of f ratios calculated by including and excluding urea uptake rates in the total nitrogen fixation in the northeastern Arabian Sea and the BOB in various seasons⁴. 1:1 lines are also shown in above plots.

The Indian Ocean has been identified as a net sink⁶ for atmospheric CO₂. It takes up ~330-430 Tg C yr⁻¹ by solubility pump, accounting for nearly 20% of the global oceanic uptake of CO₂. Biological uptake of CO₂ by the Indian Ocean has been estimated⁷ to be ~750-1320 Tg C yr⁻¹. The Indian peninsula divides the northern Indian Ocean into the Bay of Bengal (BOB) and the Arabian Sea. Although situated at the same latitude, they are significantly different in oceanographic properties such as sea surface temperature (SST), emission of biogenic gases, vertical mixing in the upper layers, mixed layer depth (MLD), nutrients, and productivity⁸. The Indian Ocean is warming faster than any other ocean basin⁹. This makes the northern Indian Ocean important for the oceanographers to study. Some major international scientific programmes such as JGOFS (Joint Global Ocean Flux Studies) aimed at assessing the role of global ocean in the carbon cycle. India also actively participated in this programme and a number of studies were carried out during the Indian JGOFS in the eastern Arabian Sea to assess its role in the global carbon cycle. The BOB is less sampled and studied compared to its western counterpart, the Arabian Sea, particularly for its biogeochemical aspects. However, some recent programs such as the BOBMEX (Bay of Bengal Monsoon Experiment), MLR (Marine Living Resources) and BOBPS (Bay of Bengal Process Studies) have focused on various oceanographic processes occurring in the Bay. Nitrogen uptake rates and new production measurements have been initiated recently under the BOBPS program⁵. In this paper we discuss the ¹⁵N

based marine productivity and new production measurements in the northern Indian Ocean.

Materials and Methods

Experimental Method

The ¹⁵N tracer technique is used to measure new and regenerated productivity^{2,10}. In this, new production is measured as the uptake rate of ¹⁵N-labelled nitrate by the phytoplankton during deck incubation and regenerated production as a sum of ¹⁵N-labelled ammonium and urea uptake rates.

Water samples were collected by a CTD rosette fitted with Niskin/Go-Flo bottles taking care to avoid trace metal contamination. Six sampling depths, 100, 80, 64, 20, 5 and 1% of surface irradiance measured using a hyperspectral radiometer (Hyperpro-II; Satlantic Inc.), were chosen to cover the euphotic zone. Individual water samples were collected for nitrate (2L volume), ammonium (2L) and urea (1L) tracer experiments. Prior to incubation at 10.00 Hrs. local time, tracers containing 99 atom% ¹⁵N (sodium nitrate, ammonium chloride and urea, obtained from SIGMA-ALDRICH, USA) were added to the bottles. After the addition of tracer the samples were incubated for 4 hrs on deck under light conditions similar to their depths of origin and filtered through pre-combusted (12 hrs at 400-450°C) 47mm diameter and 0.7µm pore size GF/F filters under a low vacuum (<70 mm Hg), dried in oven overnight at 50°C and brought to the laboratory for mass spectrometric analysis.

A CarloErba elemental analyser interfaced via conflo III to a Finnigan Delta Plus mass spectrometer

was used to analyse all the samples, using a technique for sub-microgram level ^{15}N determination¹¹. The mass spectrometer was calibrated using international standards such as IAEA- NO_3 (KNO_3 , #213) and IAEA-N-2($(\text{NH}_4)_2\text{SO}_4$, #342) along with some laboratory standards (Bovine serum albumin and Casein). Suitable and rigorous calibration steps were carried out and used to calculate the atom% ^{15}N and organic nitrogen content of the sample. A good agreement has been found between the atom% of ^{15}N measured on duplicate sets of water samples processed (Fig. 2).

Results and Discussion

^{15}N based studies in different parts of the northern Indian Ocean

Since the formulation of the concept of new and regenerated productivity² and the use of new productivity as a measure of export production³, the ^{15}N tracer technique has been extensively used to characterize the biogeochemistry of the surface ocean and to assess the ocean's role in carbon sequestration. Global scientific programmes have been carried out in the past to estimate the relationship between new and export productivity. The nitrate uptake rates (new production) and f ratios obtained from different

programmes/areas using different methods are listed in Table 1. A large spatial-temporal variation in the new productivity and f ratios are observed in the different part of the world's ocean. To assess the new productivity and f ratios in the northern Indian Ocean ^{15}N based new productivity measurements were carried out in the western and central Arabian Sea under US JGOFS programme. Since 2003 our group is involved in the eastern part of the Arabian Sea to carry out such measurements. In addition, we have also initiated new production measurements⁵ in the Bay of Bengal (BOB) under BOBPS. New production and f -ratio data from the Arabian Sea and the BOB are listed in the Table 2.

Significant seasonal and geographical variations in the new production and f -ratios are observed in the Arabian Sea. Owens *et al*¹² reported a large variation in the nitrate uptake rates, from $2.7 \text{ mmol N m}^{-2}\text{d}^{-1}$ in the central Arabian Sea to $88.9 \text{ mmol N m}^{-2}\text{d}^{-1}$ in the coastal upwelling region during Autumn, 1986. The f ratio varied from a low of 0.09 at an open ocean station to as high as 0.92 at a coastal station. N-uptake rates varied from $9.2 \text{ mmol N m}^{-2}\text{d}^{-1}$ to $40 \text{ mmol N m}^{-2}\text{d}^{-1}$ during winter monsoon and from $3.9 \text{ mmol N m}^{-2}\text{d}^{-1}$ to $24 \text{ mmol N m}^{-2}\text{d}^{-1}$ during the late summer/early winter monsoon for the central




Fig 2—Comparison between the atom% of ^{15}N measured on duplicate sets of water samples processed.

Table 1—New or export production and *f* ratios in different regions^{12,13}.

Region/season	New productivity (mmol N m ⁻² d ⁻¹)	<i>f</i> ratio
Hawaii Ocean Time-series (HOT) ¹⁴	0.6	0.10
Bermuda Atlantic Time-series Study (BATS) ¹⁵	0.9	0.15
North Atlantic Bloom Experiment (NABE) ^{16,17}	7.0	0.51
Equatorial Pacific (150°W) ^{18,19}		
November	0.5-2.7	0.08-0.21
August	0.5-4.8	0.05-0.2
Sub-Arctic Pacific ¹⁹⁻²¹		
Station-P winter	1.0-4.5	
Station-P summer	0.8-4.0	
Peru	18.3-24.2	0.30-0.42
Sub-Arctic Atlantic ²²⁻²⁴		
Iceland Basin-July	0.9-5.8	
North Atlantic	3-9	0.25-0.56
Greenland polynya	2.5	0.56
Southern Ocean ¹²		
170°W/summer	0.9-12.5	0.05-0.48
Ross Sea		

Arabian Sea¹³. Here the N-uptake rate is significantly higher in the winter (~26 mmol N m⁻²d⁻¹) than late summer (11 mmol N m⁻²d⁻¹). The *f* ratio varied from 0.03 to 0.31 and from 0.04 to 0.29 during winter and the late summer monsoons, respectively. Large variations in the N-uptake rate from 1.1 mmol N m⁻²d⁻¹ to 23.6 mmol N m⁻²d⁻¹ have also been found¹⁴ for the north-western Arabian Sea during an inter-monsoon period. It is found that the *f* ratios varied from a low of 0.07 in the open ocean region to a high of 0.52 at a coastal station. A large variation in the N-uptake rate, ranging from 0.1 to 13 mmol N m⁻²d⁻¹ has also been reported¹⁵ during the spring inter-monsoon and the summer monsoon for the northern Arabian Sea.

Very high new production has been reported¹⁶ from the north-eastern Arabian Sea during winter ranging from 1 to 4.3 mmol N m⁻²d⁻¹ with a lower *f* ratio, averaging around 0.19. A general trend of spatial increase in the new production from south to north has been observed. During late winter monsoon, new production and *f* ratio were significantly different in bloom (23 mmol N m⁻²d⁻¹, 0.65) and non-bloom

Table 2—Region, period of sampling, new or export production and *f* ratios for the Arabian Sea.

Region	Period	New productivity (mmol N m ⁻² d ⁻¹)	<i>f</i> -ratio
NW and central Arabian Sea	Autumn inter-monsoon ²⁵	2.7-5.3	0.09-0.14
	Spring inter-monsoon ²⁶	0.4-5.2	0.05-0.35
	SW monsoon ²⁶	4.3-6.4	0.05-0.4
Omani coast	Autumn inter-monsoon ²⁷	1.3-9.8	0.20-0.41
	NE monsoon ²⁸	0.56-6.8	0.01-0.27
	Autumn inter-monsoon ²⁸	0.63-2.9	0.14-0.29
	Autumn inter-monsoon ²⁵	4.1-88.9	0.12-0.92
	Spring inter-monsoon ²⁶	3.1* (n=5)	<0.05
	SW monsoon ²⁶	4.8* (n=5)	<0.05
Gulf of Oman	Autumn inter-monsoon ²⁷	0.3-2.5	0.17-0.52
	NE monsoon ²⁸	0.87-7.2	0.03-0.31
	Autumn inter-monsoon ²⁸	0.34-4.4	0.06-0.24
NE Arabian Sea	Autumn inter-monsoon ²⁷	0.3-5.1	0.07-0.41
	NE monsoon ²⁹	1.0-4.3	0.11-0.53
	Late NE monsoon ²⁹	5.7-23.2	0.33-0.61
	Late NE monsoon ³⁰	0.63-20.91	0.17-0.91
Bay of Bengal	NE monsoon ³⁰	1.95-19.70	0.46-0.87
	Spring inter-monsoon	0.35-1.58 [#]	0.51-0.82 [#]
	Autumn inter-monsoon ⁵	0.17-8.85	0.11-0.81
	Spring inter-monsoon ⁵	0.98-10.67	0.50-0.87

*Average values
#unpublished data

(5.7 mmol N m⁻²d⁻¹, 0.54) regions. An increasing trend from south to north in N-uptake rates, 2.7 to 23.0 mmol N m⁻²d⁻¹, also has been observed by Prakash *et al.*¹⁷ in the eastern Arabian Sea during late winter monsoon. In addition, they have found the presence of two different biogeochemical provinces in eastern Arabian Sea during the late winter monsoon: less productive southern (non-bloom) and more productive northern (*Noctiluca scintillans* bloom) regions. The southern sector is characterized by low column N-uptake and very low *f*-ratio (~5.5 mmol N m⁻²d⁻¹ and 0.4 respectively). The *f* ratio, although low, increased progressively towards north. This increase may be the effect of more intense winter cooling towards the north^{18,19}. The northern part is a highly productive zone, with very high N-uptake and significantly high *f* ratio (~19 mmol N m⁻²d⁻¹ and 0.8 respectively).

New production during the pre-monsoon in the BOB was higher than that in the post-monsoon (Table 2). Overall, new production for the region during the pre-monsoon averaged around 5 (±4) mmol N m⁻²d⁻¹, that was almost twice the average value observed during the post-monsoon (2.6 mmol N m⁻²d⁻¹). However, the average *f* ratio estimated for the entire region increased to 0.70 (±0.1) during pre-monsoon as compared to 0.5 during the post-monsoon⁵.

The maximum uptake observed at the surface reported by Kumar *et al.*⁵ (¹⁵N tracer method) is in good agreement with that reported by Qasim²⁰ and Radhakrishna²¹ (¹⁴C method). The surface productivity (1m depth) in the BOB (4.9 tonnes carbon km⁻² yr⁻¹) is greater than that in the Arabian Sea (3.9 tonnes carbon km⁻² yr⁻¹), but reverse is the case for column productivity²⁰. The reason offered³³ for this difference is the greater cloud cover over BOB (annual range ~ 4.1-5.1 oktas) compared to the Arabian Sea (annual range ~ 1.5-3.7 oktas). It has been argued that the cloud cover present over the Bay attenuates the excess light intensity that would otherwise reach the sea surface and result in photoinhibition of plankton in surface water as in the Arabian Sea²⁰. Another reason cited for the higher surface production in the BOB is significant nitrogen and phosphorus brought by large river runoff and rainfall into the BOB²⁰. The effect of such supply of nutrients is limited to the surface as the runoff can only dilute the upper 25 m of the Bay by 5% without influencing the waters below this depth²⁰. On the

other hand, it seems that addition of nutrients through the river runoff and rainfall in the Arabian Sea is not significant in comparison to the Bay²⁰.

The average rate of photosynthetic fixation of carbon by marine phytoplankton (primary productivity) is more than a factor of two higher in the Arabian Sea than in the BOB²². However, the time averaged sediment trap data indicate that on the basin scale, the downward flux of organic carbon in the Arabian Sea is not proportionately higher than that of the BOB, except for the upwelling region in the northwestern Arabian Sea²³⁻²⁶.

The comparable downward organic carbon fluxes in the highly productive Arabian Sea and the moderately productive BOB can be understood with the help of independent estimates of new production based on nitrogen uptake because, on a longer time scale, new production is considered theoretically equal to the export production³. It has been suggested²⁷ that new production and particle sinking are coupled over longer (annual) time scales. Values reported in two different seasons consistently show a high new production which could be one of the reasons for relatively higher downward organic carbon flux in the moderately/lowly productive BOB⁵. Prasanna Kumar *et al.*²⁸ proposed eddy pumping by the subsurface cold core eddies as a possible mechanism of nutrient injection to the oligotrophic waters in the Bay of Bengal during summer. It has been found that eddy enhances the biological productivity 2 to 8 times compared to the non-eddy region²⁸. Madhu *et al.*²⁹ observed enhanced primary production along the south-western BOB due to the October 1999 super cyclone. Therefore, such mixing by cyclones or by eddies could be a reason for higher new production in the Bay. Hence, such oceanic regions may play a more significant role in removing the excess anthropogenic CO₂ from the atmosphere, than considered so far. In addition algal blooms could sporadically increase the removal of CO₂ from the atmosphere to the deep ocean¹⁷.

Summary and Conclusions

¹⁵N technique has tremendous potential to help understand the marine nitrogen cycle. As nitrogen has a great importance for marine phytoplankton, a large number of studies have focused on elucidating the details of the marine nitrogen cycle. Presently available data show that the marine productivity is highly variable, both temporally and spatially, ranging over several orders of magnitude. The southern sector

of eastern Arabian Sea is characterized by low column N-uptake and very low f -ratio (~ 5.5 mmol N m⁻²d⁻¹ and 0.4 respectively). The f ratio, although low, increased progressively towards north. This increase may be the effect of more intense winter cooling towards the north. The north-eastern Arabian Sea is a highly productive zone, with very high N-uptake and significantly high f ratio (~ 19 mmol N m⁻²d⁻¹ and 0.8 respectively). New production during the pre-monsoon in the BOB is higher than that in the post-monsoon. Overall, new production for the region during the pre-monsoon averages around $5 (\pm 4)$ mmol N m⁻²d⁻¹, almost twice the average value observed during the post-monsoon (2.6 mmol N m⁻²d⁻¹). However, the average f ratio for the entire region increases to $0.70 (\pm 0.1)$ during pre-monsoon from 0.5 during the post-monsoon. This is in good agreement with sediment trap based export production. Though some global programs have focused on the northern Indian Ocean, still this region remains inadequately characterized for its biogeochemical aspects, particularly the BOB and north-eastern Arabian Sea. Therefore, to obtain large scale estimations for the carbon export fluxes, sustained observations over several years of ¹⁵N based export productivity measurements coupled with biogeochemical models and remote sensing are needed.

Acknowledgements

Authors are grateful to thank R. R. Navalgund, Director, Space Applications Centre (SAC, Ahmedabad) for facilities. Financial support from ISRO-GBP is gratefully acknowledged.

References

- Harrison W G, Platt T & Lewis M R, f -ratio and its relationship to ambient nitrate concentration in coastal waters, *J Plank Res*, 9(1) (1987) 235-248.
- Dugdale R C & Goering J J, Uptake of new and regenerated forms of nitrogen in primary productivity, *Limnol Oceanogr*, 12 (1967) 196-206.
- Eppley R W & Petersen B J, Particulate organic matter flux and planktonic new production in the deep ocean, *Nature* 282 (1979) 677-680.
- Kumar S., Biogeochemistry of Nitrogen Isotopes in Northern Indian Ocean, PhD thesis (2005), M.S.University Baroda, Vadodara, India.
- Kumar S, Ramesh R, Sardesai S & Sheshshayee M S, High new production in the Bay of Bengal: Possible causes and implications, *Geophys Res Lett*, 31 (2004) L18304, doi:10.1029/2004GL021005.
- Takahashi T, Sutherland S C, Sweeney C, Poisson A, Metzl N, Tilbrook B, Bates N, Wanninkhof R, Feely R A, Sabine C, Olafsson J & Nojiri Y, Global sea-air CO₂ flux based on climatological surface ocean pCO₂, and seasonal biological and temperature effects, *Deep-Sea Res-II*, 49 (2002) 1601-1622.
- Bates N R, Pequignet A C & Sabine C L, Ocean carbon cycling in the Indian Ocean: 1. Spatio-temporal variability of inorganic carbon and air-sea CO₂ gas exchange, *Global Biogeochem Cycles*, 20 (2006a) GB3020, doi: 10.1029/2005GB002491.
- Kumar M D, Biogeochemistry of the north Indian Ocean, IGBP-WCRP-SCOPE Report Series: 1 (2006), Indian National Science Academy.
- Levitus S, Antonov J I, Boyer T P & Stephens C, Warming of the world Ocean, *Science*, 287 (2000) 2225-2229.
- JGOFS Report No. 19, Protocol for the Joint Global Ocean Flux Study (JGOFS) core measurements, (1996)145-150, SCOR, Bergen, Norway.
- Owens N J P & Rees A P, Determination of nitrogen-15 at sub-microgram levels of nitrogen using automated continuous-flow isotope ratio mass spectrometer, *Analyst*, 114 (1989) 1655-1657.
- Owens N J P, Burkill P H, Mantoura R F C, Woodward E M S, Bellan I E, Aiken J, Howland R J M & Llewellyn C A, Size-fractionated primary production and nitrogen assimilation in the northwestern Indian Ocean, *Deep-Sea Res-II*, 40(3) (1993) 607-709.
- McCarthy J J, Garside C & Nevins J, Nitrogen dynamics during the Arabian Sea northeast monsoon, *Deep-Sea Res-II*, 46 (1999) 1623-1664.
- Watts L J & Owens N J P, Nitrogen assimilation and f -ratio in the northwestern Indian Ocean during an Intermonsoon period, *Deep-Sea Res-II*, 46 (1999) 725-743.
- Sambrotto R N, Nitrogen production in the Arabian Sea during spring intermonsoon and southwest monsoon seasons, *Deep-Sea Res-II*, 48 (2001) 1173-1198.
- Kumar S, Ramesh R, Dwivedi R M, Raman M, Sheshshayee M S & D'souza W, Effect of winter cooling on nitrogen uptake in the northeastern Arabian Sea (Under review: International Journal of Oceanography)
- Prakash S, Ramesh R, Sheshshayee M S, Dwivedi R M & Raman M, Quantification of New Production during a winter *Noctiluca scintillans* bloom in the Arabian Sea, *Geophys Res Lett*, 35 (2008) doi: 10.1029/2008GL033819.
- Madhupratap MS, Prasanna Kumar S, Bhattathiri P M A *et al.*, Mechanism of the biological response to winter cooling in the northeastern Arabian Sea, *Nature*, 384 (1996) 549-552.
- Prasanna Kumar S & Prasad T G, Winter cooling in the northern Arabian Sea, *Current Science*, 71 (11) (1996) 834-841.
- Qasim S Z, Biological productivity of the Indian Ocean, *Ind J Mar Sci*, 6 (1977) 122-137.
- Radhakrishna K, Primary productivity of the Bay of Bengal during March-April, 1975, *Ind J Mar Sci*, 7 (1978a) 58-60.
- Madhupratap M, Gauns M, Ramaiah N, Prasannakumar S, Muraleedharan P M, DeSouza S N, Sardesai S & Muraleedharan U, Biogeochemistry of the Bay of Bengal: physical, chemical and primary productivity characteristics of the central and western Bay of Bengal during summer monsoon 2001, *Deep-Sea Res-II*, 50 (2003) 881-896.

- 23 Haake B, Ittekkot V, Rixen T, Ramaswami V, Nair R R & Curry W B, Seasonality and interannual variability of particle fluxes to the deep Arabian Sea, *Deep-Sea Res-I*, 40 (1993) 1323-1344.
- 24 Smith W O *et al.*, New production in the northeast water Pohnya, *J Marine Systems*, 10 (1997) 199-209.
- 25 Honjo S, Dymond J, Prell W & Ittekkot V, Monsoon-controlled export fluxes to the interior of the Arabian Sea, *Deep-Sea Res-II*, 46 (1999) 1859-1902.
- 26 Lee C, Murray D M, Barber R T, Buesseler K O, Dymond J, Hedges J I, Honjo S, Manganini S J, Marra J, Moser C, Peterson M L, Prell W L & Wakeham S G, Particulate organic carbon fluxes: Compilation of results from the 1995 US JGOFS Arabian Sea Process Study, *Deep-Sea Res-II*, 45 (1998) 2489-2501.
- 27 Sarmiento J L, Dunne J & Armstrong R A, Do we now understand the Ocean's Biological Pump?, U.S. JGOFS NEWS, 12 No. 4 (2004).
- 28 Prasannakumar S, Nuncio M, Narvekar J, Kumar A, Sardesai S, De Souza S N, Gauns M, Ramaiah N & Madhupratap M, Are eddies nature's trigger to enhance biological productivity in the Bay of Bengal?, *Geophys Res Lett*, 31 L07309 (2004) doi: 10.1029/2003GL019274.
- 29 Madhu N V, Maheswaran P A, Jyothibabu R, Sunil V, Ravichandran C, Balasubramanian T, Gopalakrishnan T C & Nair K K C, Enhanced biological production off Chennai triggered by October 1999 super cyclone (Orissa), *Current Science*, 82 (12) (2002) 1472-1479.