

Studies on herbicidal activity of parthenin, a constituent of *Parthenium hysterophorus*, towards billgoat weed (*Ageratum conyzoides*)

Daizy R. Batish, R. K. Kohli, H. P. Singh and D. B. Saxena*

Department of Botany, Panjab University, Chandigarh 160 014, India

*Division of Agricultural Chemicals, Indian Institute of Agricultural Sciences, New Delhi 110 012, India

Parthenin, a constituent of *Parthenium hysterophorus*, was evaluated for herbicidal activity against *Ageratum conyzoides* (billgoat weed) under *in vitro* conditions. It inhibited/retarded germination of *A. conyzoides* at concentrations ranging from 0.02 to 0.1 mg/ml. Radicle elongation and seedling length were severely reduced. The germination of wheat, however, remained unaffected at similar concentrations. The study, therefore, paves a way for possible exploitation of parthenin as a selective herbicide.

NATURAL plant products are fast catching attention of scientists for their use as herbicides to control weeds¹. Unlike synthetic herbicides, they do not cause pollution and undesired side effects on environment. Sesquiterpene lactone parthenin – a constituent of the noxious weed *Parthenium hysterophorus*^{2,3}, has been shown to be phytotoxic against some aquatic weeds⁴. A work was, therefore, initiated to explore its herbicidal activity against a terrestrial weed *Ageratum conyzoides* (billgoat weed) commonly found in agricultural fields in north-western India.

Parthenin was extracted from healthy, shade-dried leaves of *Parthenium hysterophorus* (collected locally from wildly growing stands as per the method of Saxena

*et al.*⁵). The parthenin, so obtained, was dissolved in a few drops of ethanol and the final volume was made with distilled water to get solution ranging from concentrations 0.02 to 0.1 mg/ml (in increments of 0.02 mg/ml). To test the herbicidal property of parthenin, seeds of *A. conyzoides* (collected locally from wild stands) and wheat *Triticum aestivum* var. HD-2329 (procured from Punjab Agricultural University, Ludhiana, India) were subjected to germination in solutions of parthenin. Fifty seeds each of *A. conyzoides* and wheat were dipped in treatment solutions for 16 h. Treatment with distilled water served as control. The dipped seeds were then placed on the upper surface of Whatman filter paper no. 1 in a 12-cm petri plate. Below the filter paper a thin cotton pad soaked in the treatment solution was placed. The petri plates were incubated at 27°C, 16 h photoperiod and 75% relative humidity in a germination chamber. The entire experiment was arranged in a randomized block with four replications. The number of seeds germinated were counted and radicle length was measured after 72 h. After 10 days of treatment, the seedling length was measured. Mean values and the standard errors of the data were calculated and presented with respect to control. Besides, values of correlation coefficient between different concentrations and parameters were also calculated using polynomial regression analysis.

The results show that in response to different concentrations of parthenin, all the seeds of wheat germinated like that of water-treated control, whereas, seeds of *A. conyzoides* responded differently (Table 1). At the concentration of 0.02 mg/ml parthenin, the germination of *A. conyzoides* was nearly 50% whereas it was reduced to less than 20% at the concentrations ranging from 0.04 to 0.08 mg/ml. None of the seeds, however, could germinate at 0.1 mg/ml parthenin (Table 1).

Radicle length of *A. conyzoides* after 72 h was drastically reduced at all the concentrations of parthenin used and was nearly 50% of the control at the lowest concentration, i.e. 0.02 mg/ml. In contrast, a slight stimulation in radicle elongation of wheat was observed at this concentration (Figure 1). Almost similar trends were observed in case of seedling length (Figure 2).

Based on these results, a concentration of 0.04 mg/ml parthenin is recommended for further studies since at this concentration, germination of *A. conyzoides* was reduced by 90% whereas that of wheat remained unaffected.

From the present study it is clear that sesquiterpene lactone parthenin exhibits selective phytotoxicity and may, thus, find a potential use in the field of agriculture. In fact, a number of natural products such as cineole¹, 1,3,7-trimethylxanthine⁶, ailanthone⁷, including the sesquiterpene lactones such as artemisinin^{8,9} are fast emerging as selective phytotoxins/herbicides. The greater biological activity profiles of sesquiterpene

RESEARCH COMMUNICATIONS

Table 1. Germination of *A. conyzoides* and wheat in response to different concentrations of parthenin ($n = 4$)

Concentration (mg/ml)	Germination		% change from control	
	<i>A. conyzoides</i>	Wheat	<i>A. conyzoides</i>	Wheat
0 (control)	43.75a	50a	—	—
0.02	22.50b	50a	48.57	—
0.04	11.25c	50a	74.86	—
0.06	5.00d	50a	88.57	—
0.08	4.50d	50a	89.94	—
0.1	0	50a	100	—

Similar alphabets along a column represent insignificant difference at $p = 0.05$.




Figure 1. Effect of different concentrations of parthenin on % radicle elongation. r value along each line represents correlation coefficient between concentration and radicle length.




Figure 2. Effect of different concentrations of parthenin on % seedling length. r value along each line represents correlation coefficient between concentration and seedling length.

lactones can be attributed to the presence of α -methylene- γ -lactone and/or β -unsubstituted cyclopentenone, functionalization, ability to adopt different spatial arrangements and greater accessibility of groups for alkylations¹⁰. They react with $-SH$ groups of amino acids and proteins by non-reversible alkylation of key enzymes, thereby modifying their original properties.

Parthenin is known to possess both α -methylene- γ -lactone moiety as well as β -unsubstituted cyclopentenone which enhance its reactivity to various biological nucleophiles, thus imparting an array of properties to it. Its exact mechanism of action is not known but has been reported to change the respiratory metabolism¹¹, damage

cell membrane⁴ and even possesses growth regulatory activity comparable to indole-3-acetic acid (IAA)¹². The observed reduction in radicle elongation and seedling length in the present study could also be due to the effect of parthenin on cell elongation as a number of phytotoxins/sesquiterpenes are reported to inhibit gibberellin and IAA-induced functions¹³.

From the present results, it could be concluded that parthenin possesses herbicidal activity which can be exploited but much needs to be done. Simple and easy protocols of its extraction and enhancement in its production by using biotechnological and tissue culture techniques need to be tried. Further, its herbicidal activ-

ity can be enhanced by making suitable derivatives since it possesses exocyclic methylene double bond of lactone ring and a cyclopentenone moiety in its skeleton which provide an excellent site for structural modifications⁵.

1. Duke, S. O., *Rev. Weed Sci.*, 1986, **2**, 15-44.
2. Towers, G. H. N., Rodriguez, E., Bennett, F. D. and Rao, P. V. S., *J. Sci. Ind. Res.*, 1977, **36**, 672-684.
3. Kohli, R. K. and Daizy Rani, *Res. Bull. Panjab Univ.*, 1994, **44**, 105-149.
4. Pandey, D. K., *J. Chem. Ecol.*, 1996, **22**, 151-160.
5. Saxena, D. B., Dureja, P., Kumar, B., Daizy Rani and Kohli, R. K., *Indian J. Chem.*, 1991, **30**, 849-858.
6. Rizvi, S. J. H., Mukerji, D. and Mathur, S. N., *Agric. Biol. Chem.*, 1981, **45**, 1255-1256.
7. Heisey, R. M., *Am. J. Bot.*, 1996, **83**, 192-200.
8. Duke, S. O., Vaughin, K. C., Croom, E. M. Jr. and Elsholy, H. N., *Weed Sci.*, 1987, **35**, 499-505.
9. Chen, P. K., Polatnick, M. and Leather, G., *J. Agric. Food Chem.*, 1991, **39**, 991-994.
10. Macais, F. A., Torres, A., Molinillo, J. M. G., Varela, R. M. and Castellano, D., *Phytochemistry*, 1996, **43**, 1205-1215.
11. Kohli, R. K., Daizy Rani and Verma, R. C., *Biol. Plant.*, 1993, **33**, 567-576.
12. Batish, D. R., Kohli, R. K., Saxena, D. B. and Singh, H. P., *Plant Growth Regul.*, 1997, in press.
13. Tomaszewski, M. and Thimann, K. V., *Plant Physiol.*, 1966, **41**, 1443-1454.

ACKNOWLEDGEMENTS. D.R.B. thanks Panjab University, Chandigarh for a minor project from the unassigned grant of the UGC, R.K.K. thanks DOEn, Government of India for a project on *Ageratum* and H.P.S. thanks CSIR.

Received 24 April 1997; revised accepted 23 June 1997