

Fungal diversity inside caves of Southern India

A. John Koilraj[†], G. Marimuthu^{†,**},
K. Natarajan*, S. Saravanan*, P. Maran* and
Minna J. Hsu[‡]

[†]Department of Animal Behaviour and Physiology, School of Biological Sciences, Madurai Kamaraj University, Madurai 625 021, India

*Centre for Advanced Studies in Botany, Department of Mycology, University of Madras, Chennai 600 025, India

[‡]Department of Biological Sciences, National Sun Yat-Sen University, Kaohsiung 80424, Taiwan

Thirty-five species of sporulating mesophilous fungi and seven types of non-sporulating fungi were isolated from the soil samples collected at the entrance, twilight and dark zones of six caves. Of these, 27 species belong to Deuteromycetes, 8 species belong to Zygomycetes and one species belongs to Ascomycetes. The genus *Aspergillus* and *Penicillium* were present in large numbers in all the three zones. Fungi like *Aspergillus flavus*, *Penicillium cyclopium* and *Chaetomium* sp. were predominantly available from all the samples in all the six caves. Ten species of fungi such as *A. chevalieri*, *A. versicolor*, *A. sydowi*, *Aspergillus* sp., *Curvularia brachyspora*, *Fusarium* sp., *Geotrichum candidum*, *P. fellutanum*, *Mucor* sp. and *Rhizopus stolonifer* were isolated from light available zones. The number of species of fungi and the number of propagules/gram of soil obtained from the entrance zone soil sample were significantly more compared to that of the twilight and dark zone soil samples of the caves. The fungi, *Cunninghamella echinata* was isolated only from the dark zone of caves 2 and 5, which were occupied exclusively by a colony of carnivorous bat *Megaderma lyra*. Insectivorous bats such as *Hipposideros speoris*, *H. fulvus* and *Rhinopoma hardwickei* occupied other areas. *Syncephalis* sp., a rare fungus, has been isolated only from the dark zone of a cave.

CAVES are divided into different zones based on the prevailing light and temperature¹. According to Poulson and White², each cave has three zones: (i) twilight zone, which is located at the entrance area; (ii) middle zone, in which relative darkness prevails with fluctuating temperature; and (iii) dark zone, in which total darkness and constant temperature prevails³.

Many sets of organisms live inside caves; generally cave life is divided into three groups^{2,4}. The organisms that live inside the cave but come out periodically everyday for feeding are known as *trogloxenes* (e.g. bats). The second group of animals live their entire lives inside the cave but their conspecifics found outside the

cave are also named as *troglophiles* (e.g. cockroaches). The third group is the peculiar and most specialized for the cave environment, especially to the dark zone and are not found elsewhere. They are known as *troglobites* (e.g. cave millipede, *Glyphiulus cavernicolus*). The constancy of temperature, humidity and darkness⁵ might be the favourable factors for cave life. Another interesting feature of the cave environment is the absence of green plants deep inside caves³, due to lack of sunlight.

Fungi are remarkable for their antiquity, diversity, ubiquitous distribution and longevity⁶. Fungi are known to occur in almost all environments including soils⁷⁻⁹, among seeds¹⁰, marshy plants¹¹⁻¹³, plants in lakes¹⁴, sewage and polluted waters¹⁵, peat soils¹⁶ and also in caves^{17,18}.

Ingold¹⁹ has stated that the majority of fungi known to develop in culture are mesophiles growing between 5° and 37°C. Apparently no report is available on fungi from caves in India. An attempt was made to isolate and culture the mesophilic fungi from soil samples collected at different zones from six caves in South India.

We collected soil samples from six different caves: (1) Samanar cave, (2) Pannian cave, (3) KKB cave I, (4) KKB cave II, (5) Ramanathapuram cave and (6) Veerasihamani cave. The Samanar cave is towards the south-east, whereas Pannian cave, KKB cave I and KKB cave II are towards the north-west and all are at a distance of about 10 km from the Madurai Kamaraj University campus (9°58'N, 78°10'E). The Ramanathapuram cave and Veerasihamani cave are located at a distance of about 140 km from Tirunelveli (8°44'N, 77°42'E) towards north. The temperature and humidity were recorded inside the first four caves continuously for a period of one year^{5,20,21}. The physical factors such as temperature, relative humidity and light intensity fluctuate at the entrance and twilight zones of the caves. At the dark region, the temperature and humidity showed relatively constant values at 27°C and 95%, respectively³. We arbitrarily divided the cave into three zones in terms of the prevailing light, temperature and humidity: the entrance zone near the cave mouth into which the environmental light penetrates with varying temperature and humidity, the twilight zone in which diffused light is available and the dark zone, characterized by constant temperature and humidity with mysterious total darkness³. Cave 3 does not have a dark zone since the environmental light can be perceived even from the deeper area. The cave soil samples (bat guano) were collected from the entrance, twilight and dark zones of caves between September 1997 and August 1998. They were screened for fungi using soil dilution plating method²². Czapek-Dox agar medium (pH 4.5) was used for isolating mesophilous fungi. Six replicates were prepared for each sample. A total of 102 plates were prepared. Petri dishes were incubated at 25 ± 1°C for one

**For correspondence (e-mail: gmarimuthu@hotmail.com)

RESEARCH COMMUNICATIONS

Table 1. Quantitative estimation of the number of propagules/gram of dry soil ($\times 10^3$) of entrance, twilight and dark zones in six caves

Fungi	Entrance zone						Twilight zone						Dark zone					
	C ₁	C ₂	C ₃	C ₄	C ₅	C ₆	C ₁	C ₂	C ₃	C ₄	C ₅	C ₆	C ₁	C ₂	C ₄	C ₅	C ₆	
Deuteromycetes																		
<i>Aspergillus japonicus</i>	3	2	1	7	—	4	3	1	2	—	2	—	2	3	3	—	2	
<i>A. flavus</i>	19	34	3	5	2	21	8	27	2	4	4	19	1	2	3	—	5	
<i>A. versicolor</i>	—	11	—	—	5	—	—	7	—	—	3	—	—	—	—	—	—	
<i>A. tamari</i>	2	7	2	6	—	11	3	5	—	—	—	6	1	2	3	—	5	
<i>A. sydowii</i>	7	9	—	—	—	10	2	5	—	—	—	3	—	—	—	—	—	
<i>A. chevalieri</i>	4	—	—	—	—	7	—	—	—	—	—	—	—	—	—	—	—	
<i>A. ochraceous</i>	1	—	—	4	—	—	2	—	—	—	—	—	—	—	—	—	2	
<i>A. niger</i>	11	22	—	18	—	6	8	16	3	10	—	6	5	14	8	—	2	
<i>A. parasiticus</i>	2	21	3	5	—	9	6	10	1	2	—	—	4	4	1	—	5	
<i>A. fumigatus</i>	21	9	1	9	—	14	16	4	1	2	—	6	16	14	1	—	6	
<i>A. terreus</i>	5	2	—	—	—	—	5	3	—	—	—	—	2	2	—	—	—	
<i>A. wenti</i>	—	—	—	—	—	—	—	—	—	—	—	—	4	—	—	—	—	
<i>Aspergillus</i> sp.	14	—	—	—	—	—	—	3	—	—	2	2	—	—	—	—	—	
<i>Chrysosporium</i> sp.	—	—	—	—	—	—	—	—	—	—	—	—	2	—	—	1	1	
<i>Curvularia brachyspora</i>	7	9	—	11	5	—	7	11	6	—	5	—	—	—	—	—	—	
<i>Dactylella</i> sp.	6	9	—	4	—	11	—	—	—	—	—	—	—	—	—	—	4	
<i>Fusarium</i> sp.	4	2	1	7	—	14	12	22	—	—	—	—	6	3	4	—	10	
<i>Geotrichum candidum</i>	—	2	—	—	6	—	—	—	—	—	4	—	—	—	—	—	—	
<i>Humicola</i> sp.	2	3	—	5	—	—	—	—	—	—	—	—	—	—	—	7	5	
<i>Mycogone</i> sp.	1	2	6	9	—	14	2	2	7	11	—	16	3	4	9	—	19	
<i>Penicillium cyclopium</i>	19	11	2	8	9	10	15	8	2	7	5	9	3	5	2	3	4	
<i>P. fellutanum</i>	—	4	—	—	6	7	—	—	—	—	7	—	—	—	—	—	—	
<i>P. citreonigrum</i>	12	7	4	2	—	6	—	5	—	—	—	—	1	4	2	—	4	
<i>Penicillium</i> sp.	5	14	—	1	5	11	4	—	—	—	4	9	—	—	—	—	4	
<i>Sepedonium</i> sp.	6	—	—	—	7	—	—	—	—	—	15	—	—	—	—	—	6	
<i>Trichoderma viride</i>	—	—	—	—	—	6	—	—	—	—	—	—	—	1	—	—	—	
<i>Trichophyton</i> sp.	—	—	—	—	—	—	—	—	—	—	—	—	—	2	—	—	1	
Zygomycetes																		
<i>Absidia corymbifera</i>	—	—	19	6	—	—	—	—	16	—	—	—	—	—	14	—	—	
<i>Absidia</i> sp.	—	—	—	—	—	—	—	—	—	12	—	—	14	—	—	—	—	
<i>Cunninghamella echinata</i>	—	—	—	—	—	—	—	—	—	—	—	—	—	6	—	4	—	
<i>Mucor</i> sp.	—	—	—	9	—	—	—	—	14	—	—	—	—	—	—	—	—	
<i>Rhizopus stolonifer</i>	12	15	—	—	—	6	14	19	6	—	4	3	—	—	—	—	—	
<i>Rhizopus</i> sp.	7	—	—	—	12	—	10	—	—	—	—	—	16	—	—	—	—	
<i>Syncephalis</i> sp.	—	—	—	—	—	—	—	—	—	—	—	—	—	12	—	—	—	
Ascomycetes																		
<i>Chaetomium</i> sp.	5	16	5	9	14	18	8	21	8	9	5	19	12	29	10	14	21	
Non-sporulating fungus																		
A	—	—	—	—	—	—	—	—	—	—	—	—	—	—	3	5	11	
B	—	—	—	—	—	—	—	—	—	—	—	—	14	—	23	2	—	
C	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	29	
D	—	—	—	—	—	—	—	—	—	—	—	—	—	25	—	22	7	
E	—	—	—	—	—	—	—	—	—	—	—	—	6	2	16	9	—	
F	—	—	—	—	—	—	—	—	—	—	—	—	3	—	27	9	—	
G	—	—	—	—	—	—	—	—	—	—	—	—	4	17	—	—	—	

— Indicates absence of fungi; C₁ represents cave 1; C₂ represents cave 2; and so on.

week and the mycoflora were observed under microscope. The data were quantified to obtain the relative numbers of colony forming units (cfu) of different species of sporulating and non-sporulating fungi per gram of dry cave soil from various zones. The number of propagules of fungi was calculated by the following formula²³

$$\frac{\text{Average number of colonies per plate}}{\text{Weight of the soil}} \times \text{Dilution factor}$$

A total of 35 species of sporulating fungi belonging to 18 genera and seven species of non-sporulating fungi was isolated from the soil samples of six caves. Table shows the distribution of all the species in different




Figure 1. Light photomicrographs of a sporulating fungus *Syncephalis* sp. at a, $\times 40$; b, $\times 100$.

zones of the caves as indicated cfu. Among them, the fungi belonging to Deuteromycetes were more than those belonging to Zygomycetes and Ascomycetes. The genus *Aspergillus* was represented by 13 species, *Penicillium* represented by 4 species, *Absidia* and *Rhizopus* represented each by 2 species and *Chrysosporium*, *Curvularia*, *Dactylella*, *Fusarium*, *Geotrichum*, *Humicola*, *Mycogone*, *Sepedonium*, *Trichoderma*, *Trichophyton*, *Cunninghamella*, *Mucor*, *Syncephalis* and *Chaetomium* were represented each with a single species. The fungus *A. chevalieri*, *A. versicolor*, *A. sydowi*, *Aspergillus* sp., *C. brachyspora*, *Fusarium* sp., *G. candidum*, *P. cyclopium*, *Mucor* sp. and *R. stolonifer* were isolated only from light available areas (entrance and twilight zones). Four species of fungi, *A. wentii*, *Chrysosporium*, *Trichophyton* sp. and *Syncephalis* sp. (Figure 1a and b) were isolated only from dark zone samples. The fungi like *A. flavus*, *P. cyclopium* and *Chaetomium*

sp. were isolated invariably from all the samples in all the six caves.

The distribution of number of colonies of species of the predominantly occurring genera *Aspergillus* and *Penicillium* decreased from the entrance towards the interior, whereas, there was an increase in the number of colonies *Mycogone*, *Sepedonium*, *Absidia*, *Rhizopus*, *Mucor* and *Chaetomium* towards the interior. One way completely randomized ANOVA shows that the number of species found in the entrance zone of all the caves was significantly higher than the twilight and dark zones ($F_{2,645} = 6.18$, $P < 0.005$). The fungus *Cunninghamella echinata* was isolated in the dark zone of caves 2 and 5, occupied by a colony of carnivorous bat *Megaderma lyra*. Insectivorous bats, *Hipposideros speoris*, *H. fulvus* and *Rhinopoma hardwickei* occupied other areas. All seven different types of non-sporulated fungi were isolated only from dark zone samples of the caves.

The results of the present study provide the comparative analysis of the diversity of fungal species as well as propagules available at the entrance, twilight and dark zones of different caves. Natural agents such as flood and air introduce fungal spores and mycelium into the cave environment²⁴. The spores might also enter into the cave through organic substances such as plant and animal remains which are carried into the cave by troglodines^{4,25,26}. The spores that enter into a cave begin to grow on suitable substrate like remnants of insects¹⁷ and bat guano²⁷. The total number of colonies of fungi/gram cave soil at the entrance zone was presumably higher when compared to twilight and dark zones. The entrance and twilight zones accommodated more number of fungal species. This may be due to the higher temperature, lower relative humidity and the availability of light. In addition to physical parameters, the guano of carnivorous and insectivorous bats also plays an important role in the occurrence of specific fungal species at different zones of the caves. It was clearly pointed out that many of the fungi recorded from caves would eventually be found in the epigeal domain also because their substrates may not restrict them to the hypogean domain²⁸. Fungi form a component of the food chain that prevails in the cave ecosystem. They depend on the troglodines and troglodiles for food and are in turn eaten by other organisms²⁹. Some invertebrates like springtails³⁰ and glow worm larvae feed largely on fungus⁴. Seven species of non-sporulated fungi that were isolated from the samples collected only at the dark zone substantiate the essential role of light on sporulation. The constancy of temperature and humidity in addition to darkness might presumably be conducive for the growth of fungi.

Five species of *Syncephalis* are reported from different habitats in India³¹⁻³⁴. In the present study *Syncephalis* sp. was isolated for the first time from the dark area of a cave. According to an earlier report, the

growth from spores was disrupted at less than 90% relative humidity and germination occurred readily in total darkness than under normal conditions³⁵. The reason for the isolation of well-documented, highly sporulating species in the present study may be due to the isolation technique used, viz. the dilution plate technique. There is a possibility that many slow-growing fungi belonging to Basidiomycetes might have been eliminated by using this technique.

34. Mehotra, B. S. and Prasad, R., *Mycologia*, 1964, **56**, 905-908.
35. Headlee, T. J. and McColloch, J. W., *Kansas State Agri. College Bull.*, 1913, **191**, 287-353.

ACKNOWLEDGEMENTS. We thank the unknown referee for providing suggestions. The work is supported by a CSIR-SRF to A.J.K. and by a research project of MOEn to G.M.

Received 21 June 1999; revised accepted 16 September 1999

1. Mohr, C. E. and Poulson, T. L., *The Life of the Cave*, McGraw-Hill, New York, 1966.
2. Poulson, T. L. and White, W. B., *Science*, 1969, **165**, 971-981.
3. Koilraj, J. A. and Marimuthu, G., *Curr. Sci.*, 1998, **75**, 1111-1113.
4. Waltham, T., *Caves*, Macmillan, London, 1974.
5. Marimuthu, G., *Oecologia*, 1984, **61**, 352-357.
6. Maheshwari, R., *J. Genet.*, 1996, **75**, 239-243.
7. Gilman, J. C., *A Manual of Soil Fungi*, State College Press, Ames, 1957.
8. Barron, G. L., *The Genera of Hypomycetes from Soil*, Williams and Wilkins Co., Baltimore, 1968.
9. Paul, E. Q. and Clark, F. E., *Soil Microbiology and Biochemistry*, Academic Press, London, 1996.
10. Paul, M. C. and Mishra, R. R., *Acta Bot. Indica*, 1993, **25**, 16-42.
11. Apinis, A. E., Chesters, C. G. C. and Toligoola, H. K., *Nova Hedwigia*, 1972, **23**, 113-124.
12. Pugh, G. J. F. and Mulder, J. L., *Trans. Br. Mycol. Soc.*, 1971, **57**, 273-282.
13. Panwar, M. R. S., Ph D thesis, Meerut University, Meerut, 1979.
14. Mishra, R. C. and Dwivedi, R. S., *Acta Bot. Indica*, 1983, **11**, 28-31.
15. Udaiyan, K. and Krishnamurthy, R., *Acta Bot. Indica*, 1991, **19**, 176-184.
16. Girivasan, K. P., Rajagopal, K., Muruganandam, V. and Suryanarayanan, T. S., *Curr. Sci.*, 1998, **74**, 359-360.
17. Marvanova, L., Kalouskova Hanulakova, D. and Schanel, L., *Mikromycety Zbrasouskych jeskyni*, 1992, **46**, 243-250.
18. Semikolennykh, A. A., in *Proceedings of 12th International Congress on Speleology, Switzerland, 1997*, vol. 3, pp. 293-296.
19. Ingold, C. T., *The Biology of Fungi*, Ancho Press, London, 1975, pp. 11-33.
20. Usman, K., Ph D thesis, Madurai Kamaraj University, Madurai, 1981.
21. Habersetzer, J., Ph D thesis, Goethe University, Frankfurt, 1983.
22. Garrett, S. D., *Soil Fungi and Soil Fertility*, Pergamon, Oxford, 1963, pp. 1-165.
23. Michael, P., *Ecological Methods for Field and Laboratory Investigation*, Tata McGraw Hill, New Delhi, 1984, pp. 24-45.
24. Vandel, A., *Biospeleology*, Pergamon Press, Oxford, 1965, pp. 57-60.
25. Cubbon, B. D., in *The Science of Speleology* (eds Ford, T. D. and Cullingford, C. H. D.), Academic Press, London, 1976, pp. 423-452.
26. Biswas, J. and Kanoje, R. S., *J. Ravishankar Univ.*, 1992, **45**, 33-41.
27. Cubbon, B. D., *Trans. Cave Res. Group G.B.*, 1970, **12**, 57-74.
28. Tomaselli, R., *Atti. Ist. Bot. Lab. Crittogon. Pavia, Ser.*, 1956, **12**, 203-212.
29. Newrick, J. A., *Trans. Cave Res. Group G.B.*, 1957, **5**, 35-51.
30. Walton, G. A., in *Proceedings of University of Bristol Speleological Society, Bristol, 1944*, vol. 5, pp. 127-137.
31. Ginai, M. A., *J. Indian Bot. Soc.*, 1936, **15**, 269-284.
32. Ramakrishnan, K., *Proc. Indian Acad. Sci.*, 1955, **41**, 110-116.
33. Mehotra, B. S., *Proc. Natl. Acad. Sci.*, 1959, **29**, 94-96.