

Geometrical features of a nonlinear wavefront

Phoolan Prasad

Department of Mathematics, Indian Institute of Science, Bangalore 560 012, India

We use the equations of weakly nonlinear ray theory (WNLRT), developed by us over a number of years, to study all possible shapes which a nonlinear wavefront in a polytropic gas can have. As seen in experiments, a converging nonlinear wavefront avoids folding itself in a caustic region of a linear theory and emerges unfolded with a pair of kinks. We review the work of Baskar, Potadar and Szeftel showing the way in which the solution of a Riemann problem of the conservation form of the equations of WNLRT can be used to study the formation of new shapes of a nonlinear wavefront from a single singularity on it. We also study the ultimate result of interactions of elementary shapes on the front.

In a linear theory of wave propagation, the rays starting from various points of an initial wavefront may envelop a surface which is called caustic. For a small amplitude wave (so that linear theory is valid) propagating in a polytropic gas, which is in uniform state and at rest, the rays are straight lines orthogonal to the successive positions of the wavefront. Therefore every concave wavefront leads to the formation of a caustic and the front itself folds in the caustic region with cusp type of singularities as seen in Figure 1.

The figure represents a very interesting case of a caustic, which starts from the arête $(2, 0)$ and ends on both sides at a finite distance at points $(5, \pm 2)$ at a time $4\sqrt{2}$, whereas the front continues to have cusp type of singularity even after the time $t = 4\sqrt{2}$ due to discontinuities in the curvature of the initial wavefront at points $(1, \pm 2)$.

Experimental results¹ showed that the caustic is resolved for a front which is a moderately strong shock front. This, of course, was predicted as early as in 1957 (ref. 2) using a theory based on heuristic arguments and has been subject of discussion by us over a number of years³⁻⁷, see Figure 2. An important conclusion from these investigations is that (i) the resolution of the caustic by nonlinearity is accompanied by appearance of a new type of singularity on the front, *kink* (which was called shock-shock by Whitham) across which the amplitude and the normal direction of the front change discontinuously⁸ and (ii) the geometrical features of a weakly nonlinear wavefront^{5,9,10} and a weak shock front (moderately weak in the sense of Strutevant and Kulkarni) are qualitatively the

same. Therefore, in order to study geometrical features of a shock front, we study those of a weakly nonlinear wavefront by a simpler set of conservation laws. In this process, we also review some unpublished results obtained by Baskar, Potadar and Szeftel during 1999. A study of geometrical features of a shock using the conservation forms of the equations of a weakly nonlinear shock ray theory is in progress.

Basic equations

Consider a two-dimensional pulse of small amplitude propagating in a polytropic gas, which before the arrival of the pulse, is in uniform state and at rest. Under a short wave or high frequency approximation, the pulse can be described by a one-parameter family of nonlinear wavefronts. If a shock front appears, these weakly nonlinear wavefronts ahead of and behind the shock will keep on interacting with the shock and then disappear after that. However, we ignore this interaction and trace the history of one of these nonlinear wavefronts, as it would have been without the interaction. We use appropriate non-dimensionalization of dependent and independent variables and denote by m the Mach number of the wavefront. Then m is the wavefront velocity divided by the constant sound velocity of the undisturbed medium. Let \mathbf{q} be the




Figure 1. Linear wavefront propagation in an isotropic homogeneous medium with speed of propagation unity.

*e-mail: prasad@math.iisc.ernet.in

angle which the normal to the nonlinear wavefront (which we shall refer to simply as the front) at time t makes with a fixed direction, say the x -axis.

The position of the wavefront at time t is given parametrically by $(x(\mathbf{x}, t), y(\mathbf{x}, t))$, where \mathbf{x} is chosen such that when \mathbf{x} is fixed and t varies we move along a ray. Then (\mathbf{x}, t) represents a ray coordinate system. We note that for the waves under consideration, the $\mathbf{x} = \text{constant}$ curves and the $t = \text{constant}$ curves form an orthogonal system. Therefore, a point on the ray moving with the wavefront satisfies

$$x_t = m \cos \mathbf{q}, \quad y_t = m \sin \mathbf{q}. \tag{1}$$

m is the metric along the rays, i.e. $m dt$ represents an element of distance along a ray in the (x, y) -plane. Let g be a function such that $g d\mathbf{x}$ is an element of distance along the wavefront. Morton *et al.* showed that the equations of the weakly nonlinear ray theory (WNLRT) for m and \mathbf{q} can be written as conservation laws⁵

$$(g \sin \mathbf{q})_t + (m \cos \mathbf{q})_x = 0, \tag{2}$$

and

$$(g \cos \mathbf{q})_t - (m \sin \mathbf{q})_x = 0, \tag{3}$$

where g satisfies

$$g = (m - 1)^{-2} e^{-2(m-1)}. \tag{4}$$

Equations (1) to (4) form the complete set of equations of our WNLRT. Though the expression for g is actually given as $g = f(\mathbf{x})(m - 1)^{-2} e^{-2(m-1)}$, where $f(\mathbf{x})$ depends on the initial position of the wavefront and amplitude distribution m on it, we can choose \mathbf{x} suitably⁵ (as a function of




Figure 2. Successive positions of a shock front starting from an initial shape of the type shown in Figure 1. Rays are shown by broken lines and kinks by dots.

the arclength along the wavefront) so that $f(\mathbf{x}) = 1$. Then $g(\mathbf{x}, 0)$ is determined from eq. (4) and we can set up an initial value problem for the eqs (1) to (3):

$$x(\mathbf{x}, 0) = x_0(\mathbf{x}), \quad y(\mathbf{x}, 0) = y_0(\mathbf{x}), \tag{5}$$

$$m(\mathbf{x}, 0) = m_0(\mathbf{x}) \quad \text{and} \quad \mathbf{q}(\mathbf{x}, 0) = \mathbf{q}_0(\mathbf{x}). \tag{6}$$

Equations (2) and (3) with initial conditions (6) can be solved first and then the position of the wavefront at any time can be found by integrating (1) with respect to t (i.e. along the rays) with initial condition (5).

The system, eqs (2) and (3), is hyperbolic when $m > 1$, i.e. when the gas pressure in the wave is greater than that in the unperturbed medium. We restrict our discussion only to the situation when $m > 1$. The characteristic curves are

$$d\mathbf{x}/dt = -\sqrt{\{(m - 1)/(2g^2)\}} = I_1, \quad \text{say}; \tag{7}$$

$$d\mathbf{x}/dt = \sqrt{\{(m - 1)/(2g^2)\}} = I_2, \quad \text{say}.$$

An interesting exact solution of eqs (1) to (6) is available⁵ using two receding simple waves in $\mathbf{x} < 0$ and $\mathbf{x} > 0$. The




Figure 3. Comparison of the successive positions of the shock fronts (by NTSD and Whitham's theory) and a nonlinear wavefront starting with same initial front ($y^2 = 8x$ for $0 < x < 1$ and $y - 1 = \pm(4x - 0.5)$ for $x > 1$) and same initial amplitude distribution.

solution shows that the caustic is completely resolved due to nonlinearity and the wavefront emerges unfolded. Further, extensive numerical solutions⁶ of these equations again lead to the same result: converging rays starting from concave parts of an initial wavefront are not allowed to converge due to nonlinearity and the nonlinear wavefront, which emerges unfolded, develops kinks. Kinks are images in the (x, y) -plane of shocks in (\mathbf{x}, t) -plane of the conservation laws (2) and (3). Figure 3 shows results of one numerical computation showing the shapes of a nonlinear wavefront⁶ and a shock front by the new theory of shock dynamics (NTSD)⁷ verifying the assertion that geometrical features of these two types of fronts are qualitatively similar. For a shock front by NTSD, we need an additional initial data which is the distribution of normal derivative of a quantity (say density) behind the shock.

Elementary wave solutions and their interpretation as elementary shapes

Elementary wave solutions of eqs (2) and (3) are solutions of the form $m(\mathbf{x}, t) = m(\mathbf{x}/t)$, $\mathbf{q}(\mathbf{x}, t) = \mathbf{q}(\mathbf{x}/t)$. These are centred rarefaction wave solutions with centre at the origin and shock waves passing through the origin.




Figure 4. *a*, Example of the 1-R wave, i.e. centred simple wave of the first family in the (\mathbf{x}, t) -plane. The fan of characteristic curves is shown; *b*, Geometrical features of the front associated with the solution in *a*.

We denote centred rarefaction waves of first and second characteristic family by 1-R and 2-R, respectively. In a 1-R wave, the corresponding Riemann invariant is constant, i.e. $\mathbf{q} + \sqrt{8(m-1)} = \text{constant}$. Suppose the constant state on the left of the 1-R wave in the (\mathbf{x}, t) -plane is (m_l, \mathbf{q}_l) , then by rotation of the coordinate axes we can always choose $\mathbf{q}_l = 0$, i.e. for the 1-R wave we have (see relation (6.57), ref. 5)

$$\mathbf{q} + \sqrt{8(m-1)} = \sqrt{8(m_l-1)}. \tag{8}$$

If the state on any straight characteristic in the 1-R wave in the (\mathbf{x}, t) -plane be (m, \mathbf{q}) , then $I_1(m_l) < I(m)$ which implies $m_l > m$. Then the relation (8) gives $\mathbf{q} > 0$. At the trailing end of the 1-R wave in \mathbf{x} -space, the wave merges into a constant state (m_r, \mathbf{q}_r) and these inequalities remain valid i.e. $m_r < m_l$ and $\mathbf{q}_r > 0$. Figure 4 *a* represents a typical 1-R wave solution in the (\mathbf{x}, t) -plane and Figure 4 *b* represents its image in the (x, y) -plane. Similarly, Figure 5 *a* represents a typical 2-R wave solution in the (\mathbf{x}, t) -plane and Figure 5 *b* its image in the (x, y) -plane, where we note that that $m_r > m_l$ and $\mathbf{q}_r > 0$. We call a shape of a front in the (x, y) -plane obtained from an elementary wave




Figure 5. *a*, Example of the 2-R wave, i.e. centred simple wave of the second family in the (\mathbf{x}, t) -plane. The fan of characteristic curves is shown; *b*, Geometrical features of the front associated with the solution in *a*.

solution in the (\mathbf{x}, t) -plane as an *elementary shape*. We observe that the elementary shapes in Figure 4 b (we denote it by κ_1) and Figure 5 b (we denote it by κ_2) are convex smooth wavefronts and look almost the same geometrically but κ_1 in Figure 4 b propagates downwards on the wavefront whereas κ_2 in Figure 5 b moves upwards. Note that the rays in Figure 4 b cross the κ_1 region from below whereas in Figure 5 b they cross κ_2 from above.

When $(m_l, 0)$ and (m_r, \mathbf{q}_r) satisfy appropriate jump conditions (see relation (6.67), reference 5), we get one of the two shocks 1-S and 2-S joining two constant states $(m_l, 0)$ and (m_r, \mathbf{q}_r) and passing through $\mathbf{x} = 0$ at $t = 0$. The jumps in \mathbf{q} and m across a shock satisfy (since $\mathbf{q}_l = 0$)

$$\cos \mathbf{q}_r = (m_r g_r + m_l g_l) / (m_l g_r + m_r g_l). \tag{9}$$

Since the Lax shock inequality implies $I_1(m_r) < I_1(m_l)$ for 1-S and $I_2(m_r) < I_2(m_l)$, for 2-S, we get $m_r > m_l$ for 1-S and $m_r < m_l$ for 2-S. From the expression for g it follows that g decreases after crossing the shock ($g_r < g_l$ for 1-S and $g_r > g_l$ for 2-S). The jump relations from (2.2) and (2.3) give

$$s g_r \sin \mathbf{q}_r = g_r (m_r^2 - m_l^2) \mp (m_l g_r + m_r g_l), \tag{10}$$

where s is the shock velocity in the (\mathbf{x}, t) -plane which is negative for 1-S and positive for 2-S. This relation shows that for both shocks $\mathbf{q}_r < 0$. The images of 1-S and 2-S in the (\mathbf{x}, t) -plane to the (x, y) -plane are elementary shapes of a front, which are 1-kink (denoted by κ_1) and 2-kink (denoted by κ_2) as shown in Figure 6 a and Figure 6 b, respectively.

Solution of the Reimann problem and interpretation

In this section we briefly review some recent work of Baskar, Potadar and Szeftel (1999). A Riemann problem for the system, eqs (2) and (3) consists of solving the system with following initial conditions

$$(m, \mathbf{q})|_{t=0} = \begin{cases} (m_l, \mathbf{q}_l), & \text{we choose } \mathbf{q}_l = 0, \mathbf{x} < 0, \\ (m_r, \mathbf{q}_r), & \mathbf{x} > 0, \end{cases} \tag{11}$$

where m_l, m_r and \mathbf{q}_r are constant.

We define curves R_a and S_a ($a = 1, 2$) as loci of the points (m_r, \mathbf{q}_r) which can be joined to the point $(m_l, 0)$ by a -R and a -S waves. Figure 7 shows these curves for a typical value of $m_l = 1.2$. We note that nonlinear ray theory is valid only for small values of $m - 1$, say for $0 < m - 1 < 0.25$.





Figure 6 a, b. Rays are neither created nor lost across a kink but suddenly change their direction and since g decreases after it crosses the kink path, the rays emerge compressed.




Figure 7. R_a and S_a ($a = 1, 2$) curves in the (m, \mathbf{q}) -plane for $m = 1.2$.

If we do not go into the question of existence of the curves into consideration, the method of solution of the Riemann problem is simple. Suppose (m_r, q_r) lies in the domain A bounded by curves R_1 and R_2 as shown in Figure 7. We draw a curve R_{12} which represents the set of points joining (m_r, q_r) by 2-R wave to an intermediate state (m_i, q_i) , which lies on the R_1 curve. Thus, in this case the solution consists of the state $(m_l, 0)$ on the left of a 1-R wave continuing up to an intermediate constant state (m_i, q_i) , which ends into a 2-R wave to the right of which we have the final state (m_r, q_r) (see Figure 8 a). The shape of the wavefront at $t = 0$ and $t = t_1 > 0$ is shown in Figure 8 b. Since $(m_l, 0)$ is a state on the left, it can be joined to an intermediate state (m_i, q_i) on its right only if (m_i, q_i) lies on R_1 and not on R_2 .

We describe this result symbolically as




Figure 8. a, Solution of the Riemann problem when (m_r, q_r) is in A ; b, Shape of the wavefront at $t = 0$ and $t = t_1 > 0$ when (m_r, q_r) is in A .

$$(m_r, q_r) \in A \rightarrow R_1 R_2, \tag{12}$$

which means that when (m_r, q_r) is in A , the resultant wavefront has an elementary shape R_1 propagating below, and R_2 propagating above and these two are separated by a section of plane (or straight) front. Similarly we get the result

$$(m_r, q_r) \in B \rightarrow K_1 R_2, \tag{13}$$

as shown in Figure 9. Other results are,

$$(m_r, q_r) \in C \rightarrow K_1 K_2, \tag{14}$$

$$(m_r, q_r) \in D \rightarrow R_1 K_2. \tag{15}$$

Asymptotic result of a nonlinear wavefront⁶ when the initial wavefront is as in Figure 2 can be easily obtained. We note that when we observe the wavefront from a very large length scale, the central curved part of the initial wavefront tends to a point and the initial data reduces to

$$m(x, 0) = \begin{cases} (m_l, q_l) & x < 0, \\ (m_l, -q_l) & x > 0. \end{cases} \tag{16}$$

Choosing the direction of the x -axis perpendicular to the lower part the wavefront, solving the corresponding Riemann problem and rotating back the x -direction we get the following solution of eqs (2), (3) and (16)

$$m(x, t) = \begin{cases} (m_l, q_l) & x < -st, \\ (m_i, 0) & -st < x < st, \\ (m_l, -q_l) & st < x, \end{cases} \tag{17}$$




Figure 9. When (m_r, q_r) is in B , the front consists of a K_1 propagating downward and R_2 propagating upward.

where m_i is given by the equation

$$m_i g_{mi} + m_l g_{ml} = (m_i g_{ml} + m_l g_{mi}) \cos \mathbf{q}_i, \quad (18)$$

and

$$s = \sqrt{\{(m_i^2 - m_l^2)/(g_{ml}^2 - g_{mi}^2)\}}. \quad (19)$$

This solution when mapped into the (x, y) -plane gives the shape of the wavefront as shown in Figure 10.

Transition from one shape of the wavefront to another shape (e.g. from $R_1 R_2$ to $K_1 K_2$) as the point (m_r, \mathbf{q}_r) crosses curves R_1, R_2, S_1 or S_2 has also been discussed. The results of transitions lead to beautiful geometrical patterns.

Interaction of elementary shapes

Elementary shapes on a nonlinear wave propagate on the front. Two elementary shapes, separated by a plane portion of the front, may or may not interact. The process of interaction may take finite or infinite time depending on




Figure 10. Limiting shape, as t tends to infinity, of the nonlinear wavefront originating from an initial front as in Figure 2.




Figure 11. Successive positions of an initially sinusoidal shock front and rays plotted at $t = 0, 1, 2, \dots, 10$. Kinks have been shown by dots.

the strengths of the two elementary shapes. It is not possible to visualize the shape during the process of interaction without full numerical solution of the conservation laws, eqs (2) and (3). However, when the interaction period is finite we can easily obtain the final results, which will again consist of a pair of elementary shapes. All these geometrically beautiful results can be studied from the corresponding results on the interaction of simple waves and shock waves in the (\mathbf{x}, t) -plane^{12,13}. We can use Figure 7 for this purpose, where we note that the curves R_1, R_2, S_1 and S_2 are meaningful for more general simple waves (not just for centred waves) and shock waves (not necessarily passing through the origin in the (\mathbf{x}, t) -plane). No distinction has been made between the waves, in which characteristics converge (corresponding to compression waves




Figure 12. *a*, The front $R_1 K_1$ before the interaction; *b*, The front $K_1 R_2$ after the interaction.

in gas dynamics), and a corresponding shock. This is justified because we are considering only small changes in m . We use the symbols introduced in the previous section with a slight modification. k_2k_1 would mean a kink of second family on the lower part of the front (smaller values of \mathbf{x}) separated by a plane part (m_j, \mathbf{q}_j) of the front from a kink of the first family on the upper part of the front. To reach a state (m_r, \mathbf{q}_r) from $(m_i, 0)$ through k_2k_1 , we need to move along S_2 from $(m_i, 0)$ up to (m_j, \mathbf{q}_j) and then move along S_{j1} from (m_j, \mathbf{q}_j) up to the point (m_r, \mathbf{q}_r) . Clearly (m_r, \mathbf{q}_r) is in the region C , which implies

$$k_2k_1 \rightarrow k_1k_2, \quad (20)$$

with obvious physical interpretation. Such interactions of kinks are clearly seen in the case of propagation of an initially sinusoidal front⁷, reproduced here in Figure 11.

All possible interactions of elementary shapes, namely $k_1k_1, k_2k_2, R_1k_1, R_2k_2, k_1R_1, k_2R_2, R_2R_1, R_2k_1$ and k_2R_1 in addition to k_2k_1 mentioned earlier, have been discussed. A geometrical representation of one of these cases, namely

$$R_1k_1 \textcircled{R} k_1R_2$$

when k_1 is strong compared to R_1 has been shown in Figure 12. Note that the scales for x and y used in Figure 12 *a* and *b* are very different.

1. Sturtevant, B. and Kulkarni, V. A., *J. Fluid Mech.*, 1976, **73**, 651–671.
2. Whitham, G. B., *Linear and Nonlinear Waves*, Wiley, 1974.
3. Ramanathan, T. M., Ph D thesis, Indian Institute of Science, Bangalore, 1985.
4. Ravindran, R. and Prasad, P., *Advances in Nonlinear Waves* (ed. Debnath, L.), Pitman, 1985, vol. II, pp. 77–99.
5. Prasad, P., *Pitman Research Notes in Mathematics*, Longman, 1993, No. 293.
6. Prasad, P. and Sangeeta, K., *J. Fluid Mech.*, 1999, **385**, 1–20.
7. Monica, A. and Prasad, P., *Propagation of a curved weak shock front*, Preprint No. 01/1999, Department of Mathematics, Indian Institute of Science, Bangalore, communicated to *J. Fluid Mech.*
8. Prasad, P., *J. Indian Inst. Sci.*, 1995, **75**, 517–535.
9. Prasad, P., *J. Math. Anal. Appl.*, 1975, **50**, 470–482.
10. Prasad, P., *Wave Motion*, 1994, **20**, 21–31.
11. Morton, K. W., Prasad, P. and Ravindran, R., Technical Report 2, Department of Mathematics, Indian Institute of Science, Bangalore, 1992.
12. Courant, R. and Friedrichs, K. O., *Supersonic Flow and Shock Waves*, Interscience Publishers, reprinted by Springer Verlag, 1948.
13. Smoller, J., *Shock Waves and Reaction–Diffusion Equations*, Springer-Verlag, 1983.

ACKNOWLEDGEMENTS. This article is based on an unpublished work done by three students, namely S. Baskar and N. Potadar of the Department of Mathematics, IISc, Bangalore and Jérémie Szeftel of Ecole Normale Supérieure, Lyon. I thank Prof. Renuka Ravindran, who jointly supervised the work of Jérémie and Potadar.

Received 3 May 2000; revised accepted 7 August 2000