

Reprinted from the *Indian Journal of Chemistry*, 1971, Vol. 9, No. 1, pp. 80-81

**Structures of Fischeric Acid & Isofischeric Acid,
Two New Sesquiterpenes from
*Neolitsea fischeri****

B. S. JOSHI & V. N. KAMAT

CIBA Research Centre, Goregaon, Bombay 63

Manuscript received 3 September 1970

From the roots of *Neolitsea fischeri* Gamble two new isomeric sesquiterpenes designated as fischeric acid (I) and isofischeric acid (II) have been isolated. These belong to the germacrane and elemene types of furano terpenes and their structures have been deduced on the basis of spectroscopic and degradative evidences. Fischeric acid (I) undergoes Cope rearrangement to give isofischeric acid (II).

IN continuation of our studies on the constituents of *Neolitsea* species¹, we wish to report on the structures of two new furano sesquiterpenes isolated from the roots of *Neolitsea fischeri* Gamble. Fischeric acid (I), C₁₅H₁₈O₃, m.p. 158°, M⁺ at *m/e* 246, λ_{max} 212 nm (log ε 4.2), ν_{max}^{nujol} 1690 and 1660 cm⁻¹, and isofischeric acid (II), C₁₅H₁₈O₃, m.p. 120°, M⁺ at *m/e* 246, [α]_D 0°, ν_{max}^{nujol} 1700 cm⁻¹, were isolated

*Contribution No. 212 from CIBA Research Centre, Bombay.

from the hexane extract of the roots by chromatography on silica gel†.

Both the compounds are soluble in aq. NaHCO₃ and show positive Ehrlich test². The presence of a β-methylfuran ring could be inferred from the NMR and mass spectra. The gross structures of fischeric acid and isofischeric acid are evident from their NMR spectra (Table 1) and the mass spectral fragmentations.

Fischeric acid on hydrogenation over PtO₂ in acetic acid gave an octahydro acid (III), m.p. 163°, identical with the acid obtained by the hydrogenolysis of linderolactone³. Treatment of (I) with diazomethane afforded a pyrazoline, m.p. 117-18°, to which we assign the structure (IV), on the basis of the available spectral data.

Isofischeric acid (II) on hydrogenation gave an octahydro acid, m.p. 107-8°, formulated as (V). It also formed a pyrazoline when treated with diazomethane.

From the essential oil of the leaves of *Neolitsea sericea* Koidz., Hayashi *et al.* have isolated the sesquiterpenes sericenine (VI)^{4,5}, sericenic acid (VII)^{4,5} and isocericenine (VIII)⁶. Since fischeric acid (I) differs in its physical and spectral properties from VII, it is apparently a geometrical isomer of sericenic acid. The carboxyl and the hydrogen on

†Satisfactory elementary analyses were obtained for all new compounds.

