

**Observation and Properties of the $X(3872)$ Decaying to $J/\psi \pi^+ \pi^-$
in $p\bar{p}$ Collisions at $\sqrt{s}=1.96$ TeV**

V.M. Abazov,³² B. Abbott,⁶⁹ M. Abolins,⁶⁰ B.S. Acharya,²⁶ D.L. Adams,⁶⁷ M. Adams,⁴⁷ T. Adams,⁴⁵ M. Agelou,¹⁶ J.-L. Agram,¹⁷ S.N. Ahmed,³¹ S.H. Ahn,²⁸ G.D. Alexeev,³² G. Alkhazov,³⁶ A. Alton,⁵⁹ G. Alverson,⁵⁸ G.A. Alves,² S. Anderson,⁴¹ B. Andrieu,¹⁵ Y. Arnaud,¹² A. Askew,⁷² B. Åsman,³⁷ C. Autermann,¹⁹ C. Avila,⁷ L. Babukhadia,⁶⁶ T.C. Bacon,³⁹ A. Baden,⁵⁶ S. Baffioni,¹³ B. Baldin,⁴⁶ P.W. Balm,³⁰ S. Banerjee,²⁶ E. Barberis,⁵⁸ P. Bargassa,⁷² P. Baringer,⁵³ C. Barnes,³⁹ J. Barreto,² J.F. Bartlett,⁴⁶ U. Bassler,¹⁵ D. Bauer,⁵⁰ A. Bean,⁵³ S. Beauceron,¹⁵ F. Beaudette,¹⁴ M. Begel,⁶⁵ S.B. Beri,²⁵ G. Bernardi,¹⁵ I. Bertram,³⁸ M. Besançon,¹⁶ A. Besson,¹⁷ R. Beuselinck,³⁹ V.A. Bezzubov,³⁵ P.C. Bhat,⁴⁶ V. Bhatnagar,²⁵ M. Bhattacharjee,⁶⁶ M. Binder,²³ A. Bischoff,⁴⁴ K.M. Black,⁵⁷ I. Blackler,³⁹ G. Blazey,⁴⁸ F. Blekman,³⁰ D. Bloch,¹⁷ U. Blumenschein,²¹ A. Boehnlein,⁴⁶ T.A. Bolton,⁵⁴ P. Bonamy,¹⁶ F. Borchering,⁴⁶ G. Borissov,³⁸ K. Bos,³⁰ T. Bose,⁶⁴ C. Boswell,⁴⁴ A. Brandt,⁷¹ G. Briskin,⁷⁰ R. Brock,⁶⁰ G. Brooijmans,⁶⁴ A. Bross,⁴⁶ D. Buchholz,⁴⁹ M. Buehler,⁴⁷ V. Buescher,²¹ S. Burdin,⁴⁶ T.H. Burnett,⁷⁴ E. Busato,¹⁵ J.M. Butler,⁵⁷ J. Bystricky,¹⁶ F. Canelli,⁶⁵ W. Carvalho,³ B.C.K. Casey,⁷⁰ D. Casey,⁶⁰ N.M. Cason,⁵¹ H. Castilla-Valdez,²⁹ S. Chakrabarti,²⁶ D. Chakraborty,⁴⁸ K.M. Chan,⁶⁵ A. Chandra,²⁶ D. Chapin,⁷⁰ F. Charles,¹⁷ E. Cheu,⁴¹ L. Chevalier,¹⁶ D.K. Cho,⁶⁵ S. Choi,⁴⁴ S. Chopra,⁶⁷ T. Christiansen,²³ L. Christofek,⁵³ D. Claes,⁶² A.R. Clark,⁴² C. Clément,³⁷ Y. Coadou,⁵ D.J. Colling,³⁹ L. Coney,⁵¹ B. Connolly,⁴⁵ W.E. Cooper,⁴⁶ D. Coppage,⁵³ M. Corcoran,⁷² J. Coss,¹⁸ A. Cothenet,¹³ M.-C. Cousinou,¹³ S. Crépe-Renaudin,¹² M. Cristetiu,⁴⁴ M.A.C. Cummings,⁴⁸ D. Cutts,⁷⁰ H. da Motta,² B. Davies,³⁸ G. Davies,³⁹ G.A. Davis,⁶⁵ K. De,⁷¹ P. de Jong,³⁰ S.J. de Jong,³¹ E. De La Cruz-Burelo,²⁹ C. De Oliveira Martins,³ S. Dean,⁴⁰ K. Del Signore,⁵⁹ F. Déliot,¹⁶ P.A. Delsart,¹⁸ M. Demarteau,⁴⁶ R. Demina,⁶⁵ P. Demine,¹⁶ D. Denisov,⁴⁶ S.P. Denisov,³⁵ S. Desai,⁶⁶ H.T. Diehl,⁴⁶ M. Diesburg,⁴⁶ M. Doidge,³⁸ H. Dong,⁶⁶ S. Doulas,⁵⁸ L. Duflot,¹⁴ S.R. Dugad,²⁶ A. Duperrin,¹³ J. Dyer,⁶⁰ A. Dyshkant,⁴⁸ M. Eads,⁴⁸ D. Edmunds,⁶⁰ T. Edwards,⁴⁰ J. Ellison,⁴⁴ J. Elmsheuser,²³ J.T. Eltzroth,⁷¹ V.D. Elvira,⁴⁶ S. Eno,⁵⁶ P. Ermolov,³⁴ O.V. Eroshin,³⁵ J. Estrada,⁴⁶ D. Evans,³⁹ H. Evans,⁶⁴ A. Evdokimov,³³ V.N. Evdokimov,³⁵ J. Fast,⁴⁶ S.N. Fatakia,⁵⁷ D. Fein,⁴¹ L. Feligioni,⁵⁷ T. Ferbel,⁶⁵ F. Fiedler,²³ F. Filthaut,³¹ H.E. Fisk,⁴⁶ F. Fleuret,¹⁵ M. Fortner,⁴⁸ H. Fox,²¹ W. Freeman,⁴⁶ S. Fu,⁶⁴ S. Fuess,⁴⁶ C.F. Galea,³¹ E. Gallas,⁴⁶ E. Galyaev,⁵¹ M. Gao,⁶⁴ C. Garcia,⁶⁵ A. Garcia-Bellido,⁷⁴ J. Gardner,⁵³ V. Gavrilov,³³ D. Gelé,¹⁷ R. Gelhaus,⁴⁴ K. Genser,⁴⁶ C.E. Gerber,⁴⁷ Y. Gershtein,⁷⁰ G. Geurkov,⁷⁰ G. Ginther,⁶⁵ K. Goldmann,²⁴ T. Golling,²⁰ B. Gómez,⁷ K. Gounder,⁴⁶ A. Goussiou,⁵¹ G. Graham,⁵⁶ P.D. Grannis,⁶⁶ S. Greder,¹⁷ J.A. Green,⁵² H. Greenlee,⁴⁶ Z.D. Greenwood,⁵⁵ E.M. Gregores,⁴ S. Grinstein,¹ J.-F. Grivaz,¹⁴ L. Groer,⁶⁴ S. Grünendahl,⁴⁶ M.W. Grünewald,²⁷ W. Gu,⁶ S.N. Gurzhiev,³⁵ G. Gutierrez,⁴⁶ P. Gutierrez,⁶⁹ A. Haas,⁷⁴ N.J. Hadley,⁵⁶ H. Haggerty,⁴⁶ S. Hagopian,⁴⁵ I. Hall,⁶⁹ R.E. Hall,⁴³ C. Han,⁵⁹ L. Han,⁴⁰ K. Hanagaki,⁴⁶ P. Hanlet,⁷¹ K. Harder,⁵⁴ J.M. Hauptman,⁵² R. Hauser,⁶⁰ C. Hays,⁶⁴ J. Hays,⁴⁹ C. Hebert,⁵³ D. Hedin,⁴⁸ J.M. Heinmiller,⁴⁷ A.P. Heinson,⁴⁴ U. Heintz,⁵⁷ C. Hensel,⁵³ G. Hesketh,⁵⁸ M.D. Hildreth,⁵¹ R. Hirosky,⁷³ J.D. Hobbs,⁶⁶ B. Hoeneisen,¹¹ M. Hohlfeld,²² S.J. Hong,²⁸ R. Hooper,⁵¹ S. Hou,⁵⁹ Y. Hu,⁶⁶ J. Huang,⁵⁰ Y. Huang,⁵⁹ I. Iashvili,⁴⁴ R. Illingworth,⁴⁶ A.S. Ito,⁴⁶ S. Jabeen,⁵³ M. Jaffré,¹⁴ S. Jain,⁶⁹ V. Jain,⁶⁷ K. Jakobs,²¹ A. Jenkins,³⁹ R. Jesik,³⁹ Y. Jiang,⁵⁹ K. Johns,⁴¹ M. Johnson,⁴⁶ P. Johnson,⁴¹ A. Jonckheere,⁴⁶ P. Jonsson,³⁹ H. Jöstlein,⁴⁶ A. Juste,⁴⁶ M.M. Kado,⁴² D. Käfer,¹⁹ W. Kahl,⁵⁴ S. Kahn,⁶⁷ E. Kajfasz,¹³ A.M. Kalinin,³² J. Kalk,⁶⁰ D. Karmanov,³⁴ J. Kasper,⁵⁷ D. Kau,⁴⁵ Z. Ke,⁶ R. Kehoe,⁶⁰ S. Kermiche,¹³ S. Kesisoglou,⁷⁰ A. Khanov,⁶⁵ A. Kharchilava,⁵¹ Y.M. Kharzheev,³² K.H. Kim,²⁸ B. Klima,⁴⁶ M. Klute,²⁰ J.M. Kohli,²⁵ M. Kopal,⁶⁹ V. Korablev,³⁵ J. Kotcher,⁶⁷ B. Kothari,⁶⁴ A.V. Kotwal,⁶⁴ A. Koubarovsky,³⁴ A. Kouchner,¹⁶ O. Kouznetsov,¹² A.V. Kozelov,³⁵ J. Kozminski,⁶⁰ J. Krane,⁵² M.R. Krishnaswamy,²⁶ S. Krzywdzinski,⁴⁶ M. Kubantsev,⁵⁴ S. Kuleshov,³³ Y. Kulik,⁴⁶ S. Kunori,⁵⁶ A. Kupco,¹⁶ T. Kurča,¹⁸ V.E. Kuznetsov,⁴⁴ S. Lager,³⁷ N. Lahrachi,¹⁶ G. Landsberg,⁷⁰ J. Lazoflores,⁴⁵ A.-C. Le Bihan,¹⁷ P. Lebrun,¹⁸ S.W. Lee,²⁸ W.M. Lee,⁴⁵ A. Leflat,³⁴ C. Leggett,⁴² F. Lehner,^{46,*} C. Leonidopoulos,⁶⁴ P. Lewis,³⁹ J. Li,⁷¹ Q.Z. Li,⁴⁶ X. Li,⁶ J.G.R. Lima,⁴⁸ D. Lincoln,⁴⁶ S.L. Linn,⁴⁵ J. Linnemann,⁶⁰ R. Lipton,⁴⁶ L. Lobo,³⁹ A. Lobodenko,³⁶ M. Lokajicek,¹⁰ A. Lounis,¹⁷ J. Lu,⁶ H.J. Lubatti,⁷⁴ A. Lucotte,¹² L. Lueking,⁴⁶ C. Luo,⁵⁰ M. Lynker,⁵¹ A.L. Lyon,⁴⁶ A.K.A. Maciel,⁴⁸ R.J. Madaras,⁴² P. Mättig,²⁴ A.-M. Magnan,¹² M. Maity,⁵⁷ P.K. Mal,²⁶ S. Malik,⁵⁵ V.L. Malyshev,³² V. Manankov,³⁴ H.S. Mao,⁶ Y. Maravin,⁴⁶ T. Marshall,⁵⁰ M. Martens,⁴⁶ M.I. Martin,⁴⁸ S.E.K. Mattingly,⁷⁰ A.A. Mayorov,³⁵ R. McCarthy,⁶⁶ R. McCroskey,⁴¹ T. McMahon,⁶⁸ D. Meder,²² H.L. Melanson,⁴⁶ A. Melnitchouk,⁷⁰ X. Meng,⁶ M. Merkin,³⁴ K.W. Merritt,⁴⁶ A. Meyer,¹⁹ C. Miao,⁷⁰ H. Miettinen,⁷² D. Mihalcea,⁴⁸ C.S. Mishra,⁴⁶ J. Mitrevski,⁶⁴ N. Mokhov,⁴⁶ J. Molina,³ N.K. Mondal,²⁶ H.E. Montgomery,⁴⁶ R.W. Moore,⁵ M. Mostafa,¹ G.S. Muanza,¹⁸ M. Mulders,⁴⁶ Y.D. Mutaf,⁶⁶ E. Nagy,¹³

F. Nang,⁴¹ M. Narain,⁵⁷ V.S. Narasimham,²⁶ N.A. Naumann,³¹ H.A. Neal,⁵⁹ J.P. Negret,⁷ S. Nelson,⁴⁵ P. Neustroev,³⁶ C. Noeding,²¹ A. Nomerotski,⁴⁶ S.F. Novaes,⁴ T. Nunnemann,²³ E. Nurse,⁴⁰ V. O'Dell,⁴⁶ D.C. O'Neil,⁵ V. Oguri,³ N. Oliveira,³ B. Olivier,¹⁵ N. Oshima,⁴⁶ G.J. Otero y Garzón,⁴⁷ P. Padley,⁷² K. Papageorgiou,⁴⁷ N. Parashar,⁵⁵ J. Park,²⁸ S.K. Park,²⁸ J. Parsons,⁶⁴ R. Partridge,⁷⁰ N. Parua,⁶⁶ A. Patwa,⁶⁷ P.M. Perea,⁴⁴ E. Perez,¹⁶ O. Peters,³⁰ P. Pétrouff,¹⁴ M. Petteni,³⁹ L. Phaf,³⁰ R. Piegai,¹ P.L.M. Podesta-Lerma,²⁹ V.M. Podstavkov,⁴⁶ B.G. Pope,⁶⁰ E. Popkov,⁵⁷ W.L. Prado da Silva,³ H.B. Prosper,⁴⁵ S. Protopopescu,⁶⁷ M.B. Przybycien,^{49,†} J. Qian,⁵⁹ A. Quadt,²⁰ B. Quinn,⁶¹ K.J. Rani,²⁶ P.A. Rapidis,⁴⁶ P.N. Ratoff,³⁸ N.W. Reay,⁵⁴ J.-F. Renardy,¹⁶ S. Reucroft,⁵⁸ J. Rha,⁴⁴ M. Ridel,¹⁴ M. Rijssenbeek,⁶⁶ I. Ripp-Baudot,¹⁷ F. Rizatdinova,⁵⁴ C. Royon,¹⁶ P. Rubinov,⁴⁶ R. Ruchti,⁵¹ B.M. Sabirov,³² G. Sajot,¹² A. Sánchez-Hernández,²⁹ M.P. Sanders,⁴⁰ A. Santoro,³ G. Savage,⁴⁶ L. Sawyer,⁵⁵ T. Scanlon,³⁹ R.D. Schamberger,⁶⁶ H. Schellman,⁴⁹ P. Schieferdecker,²³ C. Schmitt,²⁴ A. Schukin,³⁵ A. Schwartzman,⁶³ R. Schwienhorst,⁶⁰ S. Sengupta,⁴⁵ E. Shabalina,⁴⁷ V. Shary,¹⁴ W.D. Shephard,⁵¹ D. Shpakov,⁵⁸ R.A. Sidwell,⁵⁴ V. Simak,⁹ V. Sirotenko,⁴⁶ D. Skow,⁴⁶ P. Slattery,⁶⁵ R.P. Smith,⁴⁶ K. Smolek,⁹ G.R. Snow,⁶² J. Snow,⁶⁸ S. Snyder,⁶⁷ S. Söldner-Rembold,⁴⁰ X. Song,⁴⁸ Y. Song,⁷¹ L. Sonnenschein,⁵⁷ A. Sopczak,³⁸ V. Sorín,¹ M. Sosebee,⁷¹ K. Soustruznik,⁸ M. Souza,² N.R. Stanton,⁵⁴ J. Stark,¹² J. Steele,⁶⁴ G. Steinbrück,⁶⁴ K. Stevenson,⁵⁰ V. Stolin,³³ A. Stone,⁴⁷ D.A. Stoyanova,³⁵ J. Strandberg,³⁷ M.A. Strang,⁷¹ M. Strauss,⁶⁹ R. Ströhmer,²³ M. Strovink,⁴² L. Stutte,⁴⁶ A. Sznajder,³ M. Talby,¹³ P. Tamburello,⁴¹ W. Taylor,⁶⁶ P. Telford,⁴⁰ J. Temple,⁴¹ S. Tentindo-Repond,⁴⁵ E. Thomas,¹³ B. Thooris,¹⁶ M. Tomoto,⁴⁶ T. Toole,⁵⁶ J. Torborg,⁵¹ S. Towers,⁶⁶ T. Trefzger,²² S. Trincas-Duvoid,¹⁵ T.G. Trippe,⁴² B. Tuchming,¹⁶ C. Tully,⁶³ A.S. Turcot,⁶⁷ P.M. Tuts,⁶⁴ L. Uvarov,³⁶ S. Uvarov,³⁶ S. Uzunyan,⁴⁸ B. Vachon,⁴⁶ R. Van Kooten,⁵⁰ W.M. van Leeuwen,³⁰ N. Varelas,⁴⁷ E.W. Varnes,⁴¹ I. Vasilyev,³⁵ M. Vaupel,²⁴ P. Verdier,¹⁴ L.S. Vertogradov,³² M. Verzocchi,⁵⁶ F. Villeneuve-Segulier,³⁹ J.-R. Vlimant,¹⁵ E. Von Toerne,⁵⁴ M. Vreeswijk,³⁰ T. Vu Anh,¹⁴ H.D. Wahl,⁴⁵ R. Walker,³⁹ N. Wallace,⁴¹ Z.-M. Wang,⁶⁶ J. Warchol,⁵¹ M. Warsinsky,²⁰ G. Watts,⁷⁴ M. Wayne,⁵¹ M. Weber,⁴⁶ H. Weerts,⁶⁰ M. Wegner,¹⁹ A. White,⁷¹ V. White,⁴⁶ D. Whiteson,⁴² D. Wicke,²⁴ D.A. Wijngaarden,³¹ G.W. Wilson,⁵³ S.J. Wimpenny,⁴⁴ J. Wittlin,⁵⁷ T. Wlodek,⁷¹ M. Wobisch,⁴⁶ J. Womersley,⁴⁶ D.R. Wood,⁵⁸ Z. Wu,⁶ T.R. Wyatt,⁴⁰ Q. Xu,⁵⁹ N. Xuan,⁵¹ R. Yamada,⁴⁶ T. Yasuda,⁴⁶ Y.A. Yatsunenko,³² Y. Yen,²⁴ K. Yip,⁶⁷ S.W. Youn,²⁸ J. Yu,⁷¹ A. Yurkewicz,⁶⁰ A. Zabi,¹⁴ A. Zatserklyaniy,⁴⁸ M. Zdrzil,⁶⁶ C. Zeitnitz,²² B. Zhang,⁶ D. Zhang,⁴⁶ X. Zhang,⁶⁹ T. Zhao,⁷⁴ Z. Zhao,⁵⁹ H. Zheng,⁵¹ B. Zhou,⁵⁹ Z. Zhou,⁵² J. Zhu,⁵⁶ M. Zielinski,⁶⁵ D. Zieminska,⁵⁰ A. Zieminski,⁵⁰ R. Zitoun,⁶⁶ V. Zutshi,⁴⁸ E.G. Zverev,³⁴ and A. Zylberstejn¹⁶

(DØ Collaboration)

¹ *Universidad de Buenos Aires, Buenos Aires, Argentina*

² *LAFEX, Centro Brasileiro de Pesquisas Físicas, Rio de Janeiro, Brazil*

³ *Universidade do Estado do Rio de Janeiro, Rio de Janeiro, Brazil*

⁴ *Instituto de Física Teórica, Universidade Estadual Paulista, São Paulo, Brazil*

⁵ *University of Alberta and Simon Fraser University, Canada*

⁶ *Institute of High Energy Physics, Beijing, People's Republic of China*

⁷ *Universidad de los Andes, Bogotá, Colombia*

⁸ *Charles University, Center for Particle Physics, Prague, Czech Republic*

⁹ *Czech Technical University, Prague, Czech Republic*

¹⁰ *Institute of Physics, Academy of Sciences, Center for Particle Physics, Prague, Czech Republic*

¹¹ *Universidad San Francisco de Quito, Quito, Ecuador*

¹² *Laboratoire de Physique Subatomique et de Cosmologie, IN2P3-CNRS, Université de Grenoble 1, Grenoble, France*

¹³ *CPPM, IN2P3-CNRS, Université de la Méditerranée, Marseille, France*

¹⁴ *Laboratoire de l'Accélérateur Linéaire, IN2P3-CNRS, Orsay, France*

¹⁵ *LPNHE, Universités Paris VI and VII, IN2P3-CNRS, Paris, France*

¹⁶ *DAPNIA/Service de Physique des Particules, CEA, Saclay, France*

¹⁷ *IReS, IN2P3-CNRS, Univ. Louis Pasteur Strasbourg, and Univ. de Haute Alsace, France*

¹⁸ *Institut de Physique Nucléaire de Lyon, IN2P3-CNRS, Université Claude Bernard, Villeurbanne, France*

¹⁹ *RWTH Aachen, III. Physikalisches Institut A, Aachen, Germany*

²⁰ *Universität Bonn, Physikalisches Institut, Bonn, Germany*

²¹ *Universität Freiburg, Physikalisches Institut, Freiburg, Germany*

²² *Universität Mainz, Institut für Physik, Mainz, Germany*

²³ *Ludwig-Maximilians-Universität München, München, Germany*

²⁴ *Fachbereich Physik, University of Wuppertal, Wuppertal, Germany*

²⁵ *Panjab University, Chandigarh, India*

²⁶ *Tata Institute of Fundamental Research, Mumbai, India*

²⁷ *University College Dublin, Dublin, Ireland*

²⁸ *Korea Detector Laboratory, Korea University, Seoul, Korea*

²⁹ *CINVESTAV, Mexico City, Mexico*

- ³⁰FOM-Institute NIKHEF and University of Amsterdam/NIKHEF, Amsterdam, The Netherlands
³¹University of Nijmegen/NIKHEF, Nijmegen, The Netherlands
³²Joint Institute for Nuclear Research, Dubna, Russia
³³Institute for Theoretical and Experimental Physics, Moscow, Russia
³⁴Moscow State University, Moscow, Russia
³⁵Institute for High Energy Physics, Protvino, Russia
³⁶Petersburg Nuclear Physics Institute, St. Petersburg, Russia
³⁷Lund University, Royal Institute of Technology, Stockholm University, and Uppsala University, Sweden
³⁸Lancaster University, Lancaster, United Kingdom
³⁹Imperial College, London, United Kingdom
⁴⁰University of Manchester, Manchester, United Kingdom
⁴¹University of Arizona, Tucson, Arizona 85721
⁴²Lawrence Berkeley National Laboratory and University of California, Berkeley, California 94720
⁴³California State University, Fresno, California 93740
⁴⁴University of California, Riverside, California 92521
⁴⁵Florida State University, Tallahassee, Florida 32306
⁴⁶Fermi National Accelerator Laboratory, Batavia, Illinois 60510
⁴⁷University of Illinois at Chicago, Chicago, Illinois 60607
⁴⁸Northern Illinois University, DeKalb, Illinois 60115
⁴⁹Northwestern University, Evanston, Illinois 60208
⁵⁰Indiana University, Bloomington, Indiana 47405
⁵¹University of Notre Dame, Notre Dame, Indiana 46556
⁵²Iowa State University, Ames, Iowa 50011
⁵³University of Kansas, Lawrence, Kansas 66045
⁵⁴Kansas State University, Manhattan, Kansas 66506
⁵⁵Louisiana Tech University, Ruston, Louisiana 71272
⁵⁶University of Maryland, College Park, Maryland 20742
⁵⁷Boston University, Boston, Massachusetts 02215
⁵⁸Northeastern University, Boston, Massachusetts 02115
⁵⁹University of Michigan, Ann Arbor, Michigan 48109
⁶⁰Michigan State University, East Lansing, Michigan 48824
⁶¹University of Mississippi, University, Mississippi 38677
⁶²University of Nebraska, Lincoln, Nebraska 68588
⁶³Princeton University, Princeton, New Jersey 08544
⁶⁴Columbia University, New York, New York 10027
⁶⁵University of Rochester, Rochester, New York 14627
⁶⁶State University of New York, Stony Brook, New York 11794
⁶⁷Brookhaven National Laboratory, Upton, New York 11973
⁶⁸Langston University, Langston, Oklahoma 73050
⁶⁹University of Oklahoma, Norman, Oklahoma 73019
⁷⁰Brown University, Providence, Rhode Island 02912
⁷¹University of Texas, Arlington, Texas 76019
⁷²Rice University, Houston, Texas 77005
⁷³University of Virginia, Charlottesville, Virginia 22901
⁷⁴University of Washington, Seattle, Washington 98195

We report the observation of the $X(3872)$ in the $J/\psi \pi^+ \pi^-$ channel, with J/ψ decaying to $\mu^+ \mu^-$, in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV. Using approximately 230 pb^{-1} of data collected with the Run II DØ detector, we observe 522 ± 100 $X(3872)$ candidates. The mass difference between the $X(3872)$ state and the J/ψ is measured to be 774.9 ± 3.1 (stat) ± 3.0 (syst) MeV/ c^2 . We have investigated the production and decay characteristics of the $X(3872)$, and find them to be similar to those of the $\psi(2S)$ state.

PACS numbers: 14.40.Gx, 14.40.Lb, 13.25.Gv, 13.25.Jx, 13.85.Ni

A new particle, the $X(3872)$, was recently discovered by the Belle Collaboration [1] in the decay mode $B^\pm \rightarrow XK^\pm$ ($X \rightarrow J/\psi \pi^+ \pi^-$) where the mass of the $X(3872)$ was measured to be 3872.0 ± 0.6 (stat) ± 0.5 (syst) MeV/ c^2 . The existence of the $X(3872)$ state has been confirmed (also decaying in the $J/\psi \pi^+ \pi^-$ mode) in

$p\bar{p}$ collisions by the CDF Collaboration [2]. At this time, it is still unclear whether this particle is a $c\bar{c}$ state, or a more complex object. See for example [3, 4, 5, 6].

The charmonium state $\psi(2S)$, with mass $m_{\psi(2S)} = 3685.96 \pm 0.09$ MeV/ c^2 [7], has the same decay mode, providing a good benchmark for comparison with the

$X(3872)$. The $\psi(2S)$ mesons produced in $p\bar{p}$ collisions can originate either from decays of B hadrons or from direct production. The $\psi(2S)$ mesons from B decays have longer effective decay lengths and tend to be less isolated than directly produced $\psi(2S)$ mesons [8].

We examine the production rate of the $X(3872)$ relative to $\psi(2S)$ as a function of the transverse momentum with respect to the beam axis (p_T), isolation and decay length, as well as a function of rapidity ($y = \frac{1}{2} \log \frac{E+P_L}{E-P_L}$, where E is the energy and P_L is the longitudinal momentum with respect to the beam axis), to determine whether the production characteristics of the $X(3872)$ are similar to those of the $\psi(2S)$. We also compare the angular decay distributions of the $\pi^+\pi^-$ and $\mu^+\mu^-$ systems in $X(3872)$ decays with those from $\psi(2S)$, to check for any differences in helicities of these two states.

The data set used in this Letter was collected in $p\bar{p}$ collisions at $\sqrt{s}=1.96$ TeV between April 2002 and January 2004, and corresponds to an integrated luminosity of approximately 230 pb^{-1} . The $D\emptyset$ detector is described elsewhere [9]. The components most important to this analysis include the vertex, central tracking and muon systems. The $D\emptyset$ tracking system consists of a silicon microstrip tracker (SMT) and a central fiber tracker (CFT), both within a 2 T solenoidal magnetic field.

The SMT has approximately 800,000 individual strips, with typical pitch of 50–80 μm , and a design optimized for tracking and vertexing over the range $|\eta| < 3$, where $\eta = -\ln[\tan(\theta/2)]$ is the pseudorapidity and θ is the polar angle measured relative to the proton beam direction. The system has a six-barrel longitudinal structure, each with a set of four layers arranged axially around the beam pipe, and interspersed with 16 radial disks. The system provides a resolution, in the plane transverse to the beam axis, for the distance of closest approach of a charged particle relative to the primary vertex of $\approx 50 \mu\text{m}$ for tracks with $p_T \approx 1 \text{ GeV}/c$, improving asymptotically to 15 μm for tracks with $p_T \geq 10 \text{ GeV}/c$.

The CFT comprises eight thin coaxial barrels, each supporting two doublets of overlapping scintillating fibers of 0.835 mm diameter, one doublet being parallel to the collision axis, and the other alternating by $\pm 3^\circ$ to provide information along the beam axis.

The muon system is located outside the calorimeters, and consists of a layer of tracking detectors and scintillation trigger counters in front of 1.8 T toroidal magnets, followed by two similar layers behind the toroids. Tracking in the muon system in the range $|\eta| < 1$ relies on 10 cm wide drift tubes [10], while 1 cm mini-drift tubes are used for $1 < |\eta| < 2$.

$J/\psi \rightarrow \mu^+\mu^-$ decays are selected by triggering on dimuons using a three-tier trigger system. The first trigger level uses hardware to form roads defined by hits in two layers of the muon scintillator system. The second trigger level uses digital signal processors to form track stubs defined by hits in the muon drift-chamber and muon scintillator systems. The third level comprises a farm of computer processors with access to the entire

event. Events passing the third-level trigger are recorded for analysis.




FIG. 1: $\Delta M = M(\mu^+\mu^-\pi^+\pi^-) - M(\mu^+\mu^-)$ for all candidates satisfying the selection requirements. The solid curve is a fit to the data and the dashed curve represents the background under each peak. The insert shows the mass distribution of the J/ψ candidates used in the analysis.

Muons are identified by extrapolating charged particle tracks from the central tracking system that match with muon track segments formed from hits in the muon system. Oppositely charged muons are combined to form J/ψ candidates, which are then combined with two oppositely charged particles assumed to be pions. At least two of these four tracks are required to have at least one hit in the SMT. To reduce background from combinatorics, events are required to satisfy the following selection criteria. An event is required to have less than 100 tracks. J/ψ candidates are selected by requiring the invariant mass of the $\mu^+\mu^-$ system to be between 2.80 and 3.35 GeV/c^2 , and the transverse momentum with respect to the beam axis (p_T) of the J/ψ is required to be greater than 5 GeV/c . In addition, the p_T of each of the two pions must be greater than 0.7 GeV/c , and the spatial separation, ΔR , between the momentum vector of the $J/\psi \pi^+\pi^-$ system and each pion momentum vector is required to have $\Delta R < 0.4$, where ΔR is defined as $\sqrt{(\Delta\phi)^2 + (\Delta\eta)^2}$, with ϕ being the azimuthal angle. The invariant mass of the two pions, $M(\pi^+\pi^-)$, is required to be greater than 0.52 GeV/c^2 , and the χ^2 of a fit to the $\mu^+\mu^-\pi^+\pi^-$ vertex is required to be less than 16 (for five degrees of freedom).

Figure 1 shows the distribution in the mass difference $\Delta M = M(\mu^+\mu^-\pi^+\pi^-) - M(\mu^+\mu^-)$, after all selections.

Superimposed is a fit to the data, where Gaussians are used to describe the $\psi(2S)$ and the $X(3872)$, and a third-order polynomial is used to account for the background. The fitted width of the $X(3872)$ peak is $17 \pm 3 \text{ MeV}/c^2$, which is consistent with detector resolution. The results of the Gaussian fit yield 522 ± 100 $X(3872)$ candidates with $\Delta M = 771.9 \pm 3.1$ (stat) MeV/c^2 . The yield of the $X(3872)$ is dependent on the value of the $M(\pi^+\pi^-)$ requirement. When we tighten the requirement on $M(\pi^+\pi^-)$ to be greater than $0.60 \text{ GeV}/c^2$, $0.65 \text{ GeV}/c^2$ and $0.70 \text{ GeV}/c^2$ and fix the mean of the $X(3872)$ to its nominal value, we obtain yields of 433 ± 98 , 336 ± 78 and 170 ± 54 events, respectively.

The position of the J/ψ mass in Fig. 1 is shifted relative to the currently accepted value [7]. This is within the uncertainties of our momentum scale, but we correct the measured ΔM by the ratio of the currently accepted mass difference between the $\psi(2S)$ and the J/ψ [7] and our corresponding measured mass difference. We assign a 100% systematic uncertainty to this mass scale correction, giving a final measurement of $\Delta M = 774.9 \pm 3.1$ (stat) ± 3.0 (syst) MeV/c^2 . We investigated systematic effects by using different background parameterizations and bin widths in our fits, and found the largest change in the central value of the mass difference to be $0.6 \text{ MeV}/c^2$.

To investigate the characteristics of the $X(3872)$ state, we study its production and decay properties, and compare the signal yields of the $X(3872)$ to the $\psi(2S)$. For example, Fig. 2 shows the ΔM distribution in two separate regions of rapidity of the $X(3872)$. The data is also separated into regions of effective proper decay length, transverse momentum of the $X(3872)$ candidate, isolation, and decay angle.

The effective proper decay length, dl , is defined as the distance in the transverse plane from the primary vertex to the decay vertex of the J/ψ scaled by the mass of the $\mu^+\mu^-\pi^+\pi^-$ system divided by the $\mu^+\mu^-\pi^+\pi^-$ system p_T .

The isolation is defined as the ratio of the $X(3872)$ momentum to the sum of the momentum of the $X(3872)$ and the momenta of all other reconstructed charged particles within a cone of radius $\Delta R = 0.5$, about the direction of the $X(3872)$ momentum.

The helicity of the $\pi^+\pi^-$ ($\mu^+\mu^-$) system can be inferred by boosting one of the pions (or muons) and the $X(3872)$ into the dipion (dimuon) rest frame, and measuring the angle θ_π (θ_μ) between them. The cosine of this angle is used for the comparison between the $X(3872)$ and $\psi(2S)$.

The widths of the $\psi(2S)$ and $X(3872)$ are fixed to the values obtained in the fit to the full sample, and the fitted number of $\psi(2S)$ and $X(3872)$ candidates in each region of chosen variables is given in Table I. The data in all regions are well represented by our choice of fitting function, with fit probabilities all greater than 15%. In Fig. 3, we present the results of Table I in a graphical




FIG. 2: ΔM distribution for two ranges of rapidity of the $\mu^+\mu^-\pi^+\pi^-$ system.

form. For example, we compare the event yield fraction:

$$f_{X(3872)} = \frac{N(X(3872)) |y| < 1}{N(X(3872)) |y| < 2},$$

to that for $\psi(2S)$:

$$f_{\psi(2S)} = \frac{N(\psi(2S)) |y| < 1}{N(\psi(2S)) |y| < 2}.$$

From the ratios of isolation and decay lengths, it is apparent that the production of the $X(3872)$ has a similar mixture of prompt and long-lived contributions as the $\psi(2S)$.

In summary, we observe the $X(3872)$ particle in approximately 230 pb^{-1} of data with a significance of 5.2 standard deviations. The mass difference between the $X(3872)$ particle and the J/ψ is $\Delta M = 774.9 \pm 3.1$ (stat) ± 3.0 (syst) MeV/c^2 . When the data are separated according to production and decay variables, we find no significant differences between the $X(3872)$ and the $c\bar{c}$ state $\psi(2S)$.

We thank Estia Eichten, Ken Lane, and Chris Quigg, for fruitful discussions. We thank the staffs at Fermilab and collaborating institutions, and acknowledge support from the Department of Energy and National Science Foundation (USA), Commissariat à l'Energie Atomique and CNRS/Institut National de Physique Nucléaire et de Physique des Particules (France), Ministry of Education and Science, Agency for Atomic Energy and RF President Grants Program (Russia), CAPES, CNPq, FAPERJ, FAPESP and FUNDUNESP (Brazil), Departments of Atomic Energy and Science and Technology

TABLE I: Number of $\psi(2S)$ and $X(3872)$ candidates for different ranges of variables. The fitted widths of the $X(3872)$ and $\psi(2S)$ are not constrained in the initial selection, but in the different ranges they are fixed to the values obtained from the full sample. Apart from the full sample, uncertainties are only from the normalization of the fitted Gaussian function and do not include small contributions from uncertainties in the background.

Regions	Number of $\psi(2S)$	Number of $X(3872)$
Initial selection	1192 ± 55	522 ± 100
(a) $p_T(J/\psi \pi^+ \pi^-) > 15 \text{ GeV}/c$	396 ± 26	179 ± 39
$p_T(J/\psi \pi^+ \pi^-) \leq 15 \text{ GeV}/c$	796 ± 41	358 ± 64
(b) $ y < 1$	741 ± 37	316 ± 57
$1 \leq y \leq 2$	449 ± 31	204 ± 49
(c) $\cos(\theta_\pi) < 0.4$	589 ± 34	288 ± 53
$\cos(\theta_\pi) \geq 0.4$	606 ± 34	244 ± 53
(d) $dl < 0.01 \text{ cm}$	838 ± 41	351 ± 66
$dl \geq 0.01 \text{ cm}$	359 ± 26	164 ± 41
(e) isolation = 1	257 ± 20	85 ± 22
isolation < 1	942 ± 44	438 ± 72
(f) $\cos(\theta_\mu) < 0.4$	593 ± 33	232 ± 46
$\cos(\theta_\mu) \geq 0.4$	602 ± 35	288 ± 60




FIG. 3: Comparison of event-yield fractions for $X(3872)$ and $\psi(2S)$ in the regions: (a) $p_T(J/\psi \pi^+ \pi^-) > 15 \text{ GeV}/c$; (b) $|y|$ of $J/\psi \pi^+ \pi^- < 1$; (c) $\cos(\theta_\pi) < 0.4$; (d) effective proper decay length, $dl < 0.01 \text{ cm}$; (e) isolation = 1; (f) $\cos(\theta_\mu) < 0.4$.

(India), Colciencias (Colombia), CONACyT (Mexico), KRF (Korea), CONICET and UBACyT (Argentina), The Foundation for Fundamental Research on Matter (The Netherlands), PPARC (United Kingdom), Ministry of Education (Czech Republic), Natural Sciences and Engineering Research Council and WestGrid Project (Canada), BMBF (Germany), A.P. Sloan Foundation, Civilian Research and Development Foundation, Research Corporation, Texas Advanced Research Program, and the Alexander von Humboldt Foundation.

[*] Visitor from University of Zurich, Zurich, Switzerland.

[†] Visitor from Institute of Nuclear Physics, Krakow, Poland.

[1] S. K. Choi *et al.* (Belle Collaboration), Phys. Rev. Lett.

91, 262001 (2003).

[2] D. Acosta *et al.* (CDF II Collaboration), submitted to Phys. Rev. Lett., hep-ex/0312021.

[3] F. E. Close, P. R. Page, “The $D^{*0}\bar{D}^0$ threshold reso-

- nance”, Phys. Lett. **B578**, 119 (2004), hep-ph/0309253.
- [4] E. J. Eichten, K. Lane, and C. Quigg, “Charmonium levels near threshold and the narrow state $X(3872) \rightarrow \pi^+\pi^-J/\psi$,” FERMILAB-Pub-04/001-T, hep-ph/0401210, and references therein.
- [5] C. Quigg, “Quarkonium: New developments,” hep-ph/0403187.
- [6] N. Törnqvist, “Isospin breaking of the narrow charmonium state of Belle at 3872 MeV as a deuson”, hep-ph/0402237.
- [7] K. Hagiwara *et al.*, Phys. Rev. D **66**, 010001 (2002) and 2003 off-year partial update for the 2004 edition available on the PDG WWW pages (URL: <http://pdg.lbl.gov/>).
- [8] S. Abachi *et al.* (DØ Collaboration), Phys. Lett. **B370**, 239 (1996); F. Abe *et al.* (CDF Collaboration), Phys. Rev. Lett. **79**, 572 (1997).
- [9] V. Abazov, *et al.*, in preparation for submission to Nucl. Instrum. Methods Phys. Res. A, and T. LeCompte and H.T. Diehl, “The CDF and D0 Upgrades for Run II”, Ann. Rev. Nucl. Part. Sci. **50**, 71 (2000).
- [10] S. Abachi, *et al.*, Nucl. Instrum. Methods Phys. Res. A **338**, 185 (1994).