

Relations Between Quantum Maps and Quantum States

M. Asorey

Departamento de Física Teórica, Universidad de Zaragoza
50009 Zaragoza, Spain
e-mail: asorey@saturno.unizar.es

A. Kossakowski

Institute of Physics, Nicolaus Copernicus University
Toruń 87–100, Poland
e-mail: kossak@phys.uni.torun.pl

G. Marmo

Dipartimento di Scienze Fisiche, Università Federico II di Napoli
and
INFN, Sezione di Napoli
Complesso Univ. di Monte Sant'Angelo, Via Cintia, 80125 Napoli, Italy
e-mail: marmo@na.infn.it

E. C. G. Sudarshan

Department of Physics, University of Texas at Austin
Austin, Texas 78712-1081
e-mail: sudarshan@physics.utexas.edu

February 1, 2008

Abstract

The relation between completely positive maps and compound states is investigated in terms of the notion of quantum conditional probability.

1 Introduction

In quantum information two problems play a relevant role, the first one concerns the study of the dynamical change of states of a system by means of completely positive maps, commonly called channels, the second one is to describe corre-

lations between the initial and final states; such correlations are described by compound states.

The connection between the two concepts is based on a very general principle. Indeed, in any Hilbert space \mathcal{H} there is an one-to-one correspondence between the set \mathcal{P}_q^p of p -contravariant q -covariant tensors and the set \mathcal{P}_{p+q} of $p+q$ covariant tensors. The equivalence is due to the identification of \mathcal{H} and its dual space \mathcal{H}^* by means of the hermitian product. Consequently, endomorphisms of \mathcal{H} which are in \mathcal{P}_1^1 are in one-to-one correspondence with 2-covariant tensors in \mathcal{P}_2 . In particular, if we consider the Hilbert spaces of Hilbert-Schmidt operators on \mathcal{H}_1 and \mathcal{H}_2 , any map of the Hilbert-Schmidt operators on \mathcal{H}_1 into those on \mathcal{H}_2 is associated to a state on the tensor product of the spaces of Hilbert-Schmidt operators on \mathcal{H}_1 and \mathcal{H}_2 .

The correspondence can also be formulated at C^* -algebraic level. However, only the finite-dimensional case will be considered here.

Let us consider two systems described by $(M_n, S(M_n))$ and $(M_m, S(M_m))$. The first one describes an initial (*input*) system and the second one a final (*output*) system. The symbol M_n stands for the algebra of $n \times n$ complex matrices and the symbol $S(M_n)$ stands for the set of all states on M_n , i.e. the set of all density matrices. Moreover, I_n denotes the identity matrix in M_n . Let us consider a map $\varphi^*: S(M_n) \rightarrow S(M_m)$, such that its dual map $\varphi: M_m \rightarrow M_n$ is completely positive and normalized, i.e. $\varphi(I_m) = I_n$.

For an initial state $\rho \in M_n$ and final state $\varphi^*(\rho) \in M_m$, a composite state $\omega \in S(M_n \otimes M_m)$ should satisfy the following two conditions:

- i) $\omega(a \otimes I_m) = \rho(a)$, for all $a \in M_n$
- ii) $\omega(I_n \otimes b) = \varphi^*(\rho)(b)$ for all $b \in M_m$.

It is well known that joint probability measures do not generally exist for quantum systems, therefore it is difficult to define a compound state ω satisfying the above conditions.

The first construction of a compound state ω satisfying the above two conditions has been given by Ohya [1,2]. Let $\rho \in M_n$, then ρ has the following spectral decomposition

$$\rho = \sum_k \lambda_k m_k \rho_k, \quad (1.1)$$

where

$$\rho_k = \frac{1}{m_k} p_k, \quad m_k = \text{Tr } p_k, \quad (1.2)$$

λ_k are the eigenvalues of ρ , and p_k are eigenprojectors of ρ , respectively.

Then for any $\varphi^*: S(M_n) \rightarrow S(M_m)$ the compound state $\omega_\varphi \in S(M_n \otimes M_m)$ has the form

$$\omega_\varphi = \sum_k \lambda_k m_k \rho_k \otimes \varphi^*(\rho_k). \quad (1.3)$$

Let us observe that ω_φ is a separable state on $M_n \otimes M_m$, φ is a positive normalized map $\varphi: M_m \rightarrow M_n$, and the construction of ω_φ is non-linear with respect to ρ .

We notice that the Ohya compound state can be constructed in the case of general C^* -algebraic setting.

The construction of compound states which will be studied in the present paper can be described as follows.

Let $\sigma \in S(M_n \otimes M_m)$, and suppose that

$$\mathrm{tr}_{\mathbb{C}^m} \sigma = \sigma_1 > 0. \quad (1.4)$$

Then, one can define the following operator

$$\begin{aligned} \pi(\sigma) &: \mathbb{C}^n \otimes \mathbb{C}^m \longrightarrow \mathbb{C}^n \otimes \mathbb{C}^m \\ \pi(\sigma) &= (\sigma_1^{-1/2} \otimes I_m) \sigma (\sigma_1^{-1/2} \otimes I_m), \end{aligned} \quad (1.5)$$

which has the properties

$$\pi(\sigma) \geq 0 \quad (1.6)$$

$$\mathrm{tr}_{\mathbb{C}^m} \pi(\sigma) = I_m. \quad (1.7)$$

It follows from (1.4) and (1.5) that the operator π is the quantum analogue of classical conditional probability. Another definition of quantum conditional probability has been given in [3].

Definition 1 A map

$$\pi: \mathbb{C}^n \otimes \mathbb{C}^m \longrightarrow \mathbb{C}^n \otimes \mathbb{C}^m \quad (1.8)$$

is a quantum conditional probability (QCPO) iff satisfies (1.6) and (1.7).

For a given π and any $\rho \in S(M_n)$ one can define a compound state

$$\omega = (\rho^{1/2} \otimes I_m) \pi (\rho^{1/2} \otimes I_m), \quad (1.9)$$

which has the following properties

$$\mathrm{tr}_{\mathbb{C}^m} \omega = \rho \quad (1.10)$$

and

$$\mathrm{tr}_{\mathbb{C}^n} \omega = \mathrm{tr}_{\mathbb{C}^n} \pi(\rho \otimes I_m) = \varphi^*(\rho), \quad (1.11)$$

where

$$\varphi^*: S(M_n) \longrightarrow S(M_m).$$

The study of the relation between φ^* and π is based on duality between quantum maps and composite states which has been investigated in detail in [4,11] (see also references therein).

2 Classification of Composite States and Positive Maps

It has been shown above that the construction of composite states is based on the notion of quantum conditional probability operator (*QCPO*) $\pi: \mathbb{C}^n \otimes \mathbb{C}^m \rightarrow \mathbb{C}^n \otimes \mathbb{C}^m$. In what follows the case $m = n$ will be considered for simplicity.

Let \mathcal{H} be a Hilbert space and $\mathcal{H}_1 = \mathcal{H}_2 = \mathcal{H}$, the following order in the tensor product $\mathcal{H} \otimes \mathcal{H} = \mathcal{H}_2 \otimes \mathcal{H}_1$ will be used, and the partial trace with respect to the Hilbert space \mathcal{H}_a will be denoted by tr_a .

Let $\{e_1, \dots, e_n\}$ be a fixed orthonormal basis in \mathbb{C}^n and $e_{kl} = e_k(e_l, \cdot)$ be the corresponding basis in M_n , then $a \in M_n$ can be written in the form

$$a = \sum_{i,j=1}^n (e_i, a e_j) e_{ij} = \sum_{i,j=1}^n \text{tr}(a e_{ij}^*) e_{ij}, \quad (2.1)$$

and the transpose map $\mathcal{T}: M_n \rightarrow M_n$ (with respect to the basis $\{e_1, \dots, e_n\}$) has the form

$$\mathcal{T}(a) = \sum_{i,j=1}^n (e_j, a e_i) e_{ij} = \sum_{i,j=1}^n e_{ij} a e_{ij}. \quad (2.2)$$

A generic element $x \in \mathbb{C}^n \otimes \mathbb{C}^n$ can be written in the form

$$x = \sum_{i=1}^n x_i \otimes e_i = \sum_{i=1}^n (a e_i) \otimes e_i, \quad (2.3)$$

where $x_1, x_2, \dots, x_n \in \mathbb{C}^n$ and $a \in M_n$.

With every $a \in M_n$, such that $\text{tr}(a^* a) = 1$, one can associate one-dimensional projections $p_a: \mathbb{C}^n \otimes \mathbb{C}^n \rightarrow \mathbb{C}^n \otimes \mathbb{C}^n$

$$p_a = \sum_{i,j=1}^n a e_{ij} a^* \otimes e_{ij}. \quad (2.4)$$

Moreover, two projections p_a and p_b are orthogonal provided that $\text{tr}(a^* b) = 0$. As a consequence of (2.4) any positive operator $\hat{A}: \mathbb{C}^n \otimes \mathbb{C}^n \rightarrow \mathbb{C}^n \otimes \mathbb{C}^n$ has the form

$$\hat{A} = \sum_{i,j=1}^n \sigma_{ij} \otimes e_{ij} = \sum_{i,j=1}^n \varphi(e_{ij}) \otimes e_{ij}, \quad (2.5)$$

where

$$\varphi(e_{ij}) = \sum_{\alpha=1}^{n^2} \lambda_\alpha a_\alpha e_{ij} a_\alpha^* \quad (2.6)$$

and

$$\lambda_\alpha \geq 0, \quad \text{tr}(a_\alpha a_\beta^*) = \delta_{\alpha\beta},$$

i.e. $\{a_1, a_2, \dots, a_{n^2}\}$ is an orthonormal basis in M_n .

Relation (2.5) can also be rewritten in the form

$$\hat{\sigma} = (\varphi \otimes \text{id}) \sum_{i,j=1}^n e_{ij} \otimes e_{ij}, \quad (2.7)$$

which gives the relation between elements of $(M_n \otimes M_n)^+$ and completely positive maps in M_n . In order to classify the states on $M_n \otimes M_n$ it is convenient to introduce the following cones in $M_n \otimes M_n$:

$$V_s = \text{conv} \left\{ \sum_{i,j=1}^n a e_{ij} a^* \otimes e_{ij} : a \in M_n, \text{rank } a \leq s \right\} \quad (2.8)$$

where $\text{conv } X$ means convex (not normalized) set generated by elements of X , and

$$V^s = (\text{id} \otimes \mathcal{T})V_s, \quad (2.9)$$

where \mathcal{T} is the transpose map on M_n , i.e.

$$V^s = \text{conv} \left\{ \sum_{i,j}^n a e_{ij} a^* \otimes e_{ji} : a \in M_n, \text{rank } a \leq s \right\}. \quad (2.10)$$

It follows from (2.8) and (2.9) that the following chains of inclusions

$$\begin{aligned} V_1 \subset V_2 \subset \dots \subset V_n, \quad V^1 \subset V^2 \subset \dots \subset V^n, \\ V_1 \cap V^1 \subset V_2 \cap V^2 \subset \dots \subset V_n \cap V^n \end{aligned} \quad (2.11)$$

hold true.

It is clear that V_n coincides with the cone $(M_n \otimes M_n)^+$ of all positive semidefinite elements of $M_n \otimes M_n$, $V_1 = V^1$ is the cone generated by elements $a \otimes b$, where a, b are positive elements of M_n , i.e. V_1 coincides with separable (not normalized) states on $M_n \otimes M_n$, while $V_n \cap V^n$ is the set of all (not normalized) PPT states on $M_n \otimes M_n$ (by definition).

Using the results of [4, 5, 6] the above cones can be used to classify positive maps. Let P_s , P^s and $P_s \cup P^r$ be the cones of s -positive, s -copositive maps and sums of s -positive and r -copositive ones, respectively. One can verify that

$$\begin{aligned} \varphi \in P_s &\iff (\varphi \otimes \text{id})V_s \in (M_n \otimes M_n)^+ \\ \varphi \in P^s &\iff (\varphi \otimes \text{id})V^s \in (M_n \otimes M_n)^+ \end{aligned} \quad (2.12)$$

and

$$\varphi \in P_s \cup P^r \iff (\varphi \otimes \text{id})V_s \cap V^r \in (M_n \otimes M_n)^+.$$

Relations (2.12) can be considered as an extension of the Horodecki theorem [7] which gives the characterization of the cone V_1 in terms of positive maps. It should be pointed out that our knowledge concerning the above mentioned cones is rather poor. In fact only the structure of cones V_n and V^n is known.

In the n -dimensional case an example of an element $V_2 \cap V^2$ which is not separable has been given in [8].

The cones V_r and V^r and $V_r \cap V^s$ can be used for classification of completely positive maps.

Definition 2 A completely positive map $\varphi : M_n \longrightarrow M_n$ is said to be *s-completely positive* if

$$\sum_{i,j=1}^n \varphi(e_{ij}) \otimes e_{ij} \in V_s \quad (2.13)$$

and *(r, s)-completely positive* if

$$\sum_{i,j=1}^n \varphi(e_{ij}) \otimes e_{ij} \in V_r \otimes V^s.$$

Let us observe that the set \mathcal{P}_{rs} of all (r, s) -completely positive maps is a subset of $P_n \cap P^n$, the subset of maps which are completely positive and completely copositive. On the other hand, (r, s) -completely positive maps generate PPT-states since the inclusion $V_r \cap V^s \subseteq V_n \cap V^n$ holds.

It is also convenient to consider *s-completely positive* maps which are *k-copositive*, i.e. elements of the set $P_n \cap P^k$ ($k < n$) which generate NPT states. The construction of composite states in terms of QCPO will be used to find out some classes of PPT and NPT states.

Indeed, taking into account (2.5), one finds out that the general form of π is given by

$$\pi = \sum_{i,j}^n \varphi(e_{ij}) \otimes e_{ij}, \quad (2.14)$$

where φ is a completely positive normalized map in M_n , i.e., $\varphi(I_n) = I_n$. It follows from (2.14) and (1.8) that the composite state can be written in the form

$$\omega = \sum_{i,j=1}^n \rho^{1/2} \varphi(e_{ij}) \rho^{1/2} \otimes e_{ij} \quad (2.15)$$

and the relations

$$\begin{aligned} \text{tr}_1 \omega &= \rho, \\ \text{tr}_2 \omega &= (\mathcal{T} \circ \varphi^*)(\rho) = \psi^*(\rho), \end{aligned} \quad (2.16)$$

hold true.

The dual map ψ can be written as

$$\psi(e_{ij}) = \varphi(e_{ji}) = (\varphi \circ \mathcal{T})(e_{ij}) \quad \text{or} \quad \psi^* = \mathcal{T} \circ \varphi^*. \quad (2.17)$$

The properties of the composite state ω can be summarized as follows.

Corollary 1 The composite state ω is a PPT iff φ is (r, s) -completely positive.

Corollary 2 The composite state ω is NPT iff φ is r -completely positive and k -copositive ($k < n$) provided $\text{rank } \rho = n$.

The above results imply that the construction of PPT-states is reduced to normalized completely positive and completely copositive maps, while, entangled but not PPT-states are induced by normalized completely positive and k -completely copositive maps.

Next we will analyze some examples of normalized completely positive and completely copositive maps.

Example 1 Let us consider the following QCPO:

$$\pi_\lambda = \frac{1-\lambda}{n} I_n \otimes I_n + \lambda \sum_{i,j=1}^n e_{ij} \otimes e_{ij}, \quad (2.18)$$

where

$$-\frac{1}{n^2-1} \leq \lambda \leq \frac{1}{n+1}. \quad (2.19)$$

The above π is, up to normalization, the Horodecki state [8], and for λ satisfying (2.19) π is separable, i.e., $\pi_\lambda \in V^1 \cap V_1$.

Let $\psi: M_n \rightarrow M_n$ be a normalized positive map, then

$$\sigma_\lambda = (\psi \otimes \text{id}) \pi_\lambda \quad (2.20)$$

is a QCPO which is separable, and the relation

$$\sum_{ij} \varphi_\lambda(e_{ij}) \otimes e_{ij} = (\psi \otimes \text{id}) \pi_\lambda \quad (2.21)$$

defines completely positive and completely copositive maps of the form

$$\varphi_\lambda(e_{ij}) = \frac{1-\lambda}{n} I_n \delta_{ij} + \lambda \psi(e_{ij}), \quad (2.22)$$

which is $(1, 1)$ -completely positive.

Example 2 Let us consider the following QPCO:

$$\pi_\gamma = N_\gamma^{-1} \left(n I_n \otimes I_n + \sum_{i,j=1}^n a_{ij} \otimes e_{ij} \right), \quad (2.23)$$

where

$$a_{ij} = n e_{ij}, \quad i \neq j, \quad (2.24)$$

$$a_{ij} = \left(1 - \frac{1}{\gamma^2} \right) (\gamma^2 e_{i+1, i+1} - e_{n+i-1, n+i-1}) \pmod{n}, \quad (2.25)$$

$$N_\gamma = n^2 + \left(1 - \frac{1}{\gamma^2} \right) (\gamma^2 - 1), \quad \gamma^2 > 0. \quad (2.26)$$

It has been shown in [4] that $\pi_\gamma \in V_2 \cap V^2$ but it is not separable.

The relation

$$\sum_{i,j=1}^n \varphi_\gamma(e_{ij}) \otimes e_{ij} = \pi_\gamma \quad (2.27)$$

defines a normalized (2, 2)-completely positive map φ which has the form

$$\varphi_\gamma(e_{ij}) = \frac{1}{N_\gamma} (n I_n \delta_{ij} + a_{ij}). \quad (2.28)$$

Example 3 The map φ given by (2.28) has the following form

$$\varphi(a) = \sum_{i,j}^n c_{ij} e_{ij} a e_{ij}^* + \mu a, \quad (2.29)$$

where

$$\varphi(I_n) = \sum_{i=1}^n e_{ii} \left(\sum_{j=1}^n c_{ij} + \mu \right). \quad (2.30)$$

Taking into account (2.2) one finds that the relation

$$\varphi(\mathcal{T}(a)) = \mathcal{T}(\varphi(a)) \quad (2.31)$$

holds.

Theorem 1 *The map φ , as in (2.29), is completely positive iff the following conditions*

$$c_{ij} \geq 0, \quad i \neq j \quad (2.32)$$

and

$$[c_{ii} \delta_{ij} + \mu] \geq 0 \quad (2.33)$$

are satisfied.

Proof. The map φ can be written in the form

$$\varphi(a) = \sum_{i,j=1}^n (\delta_{ij} c_{ii} + \mu) e_{ii} a e_{jj} + \sum_{i \neq j} c_{jj} e_{ij} a e_{ij}^* \quad (2.34)$$

on the other hand the map ψ is completely positive iff it has the form

$$\psi(a) = \sum_{\alpha,\beta=1}^{n^2} \lambda_{\alpha\beta} f_\alpha a f_\beta^*, \quad (2.35)$$

where

$$\text{tr}(f_\alpha f_\beta^*) = \delta_{\alpha\beta} \quad (2.36)$$

and

$$[\lambda_{\alpha\beta}] \geq 0. \quad (2.37)$$

Taking into account (2.35)–(2.37) and (2.34), one finds conditions (2.33) and (2.34).

Theorem 2 *The map φ is completely copositive iff the following conditions*

$$c_{ii} + \mu \geq 0, \quad (2.38)$$

$$c_{ij} + c_{ji} \geq 2\mu, \quad i \neq j, \quad (2.39)$$

$$c_{ij} + c_{ji} \geq -2\mu, \quad i \neq j, \quad (2.40)$$

$$c_{ij}c_{ji} \geq \mu^2, \quad i \neq j \quad (2.41)$$

are satisfied.

Proof. The map φ is completely copositive iff the map $\mathcal{T}\varphi$ is completely positive.

Taking into account (2.2) and (2.29) one finds

$$(\mathcal{T} \circ \varphi)(a) = \sum_{i=1}^n (c_{ii} + \mu) e_{ii} a e_{ii} + \sum_{i \neq j} (c_{ij} e_{ij} a e_{ij}^* + \mu e_{ij} a e_{ij}). \quad (2.42)$$

From (2.42) one finds the condition (2.38). Let us introduce trace orthonormal operators

$$\begin{aligned} f_{ij} &= \frac{1}{\sqrt{2}}(e_{ij} + e_{ji}), \\ g_{ij} &= \frac{-i}{\sqrt{2}}(e_{ij} - e_{ji}), \quad i < j. \end{aligned} \quad (2.43)$$

Then the equality

$$\begin{aligned} \sum_{i \neq j} (c_{ij} e_{ij} a e_{ij}^* + \mu e_{ij} a e_{ij}) &= \sum_{k < l} \left\{ \left(\frac{1}{2}(c_{kl} + c_{lk}) + \mu \right) f_{kl} a f_{kl} \right. \\ &\quad + \left(\frac{1}{2}(c_{kl} + c_{lk}) - \mu \right) g_{kl} a g_{kl} - \frac{i}{2}(c_{kl} - c_{lk}) f_{kl} a g_{kl} \\ &\quad \left. + \frac{i}{2}(c_{kl} - c_{lk}) g_{kl} a f_{kl} \right\} \end{aligned} \quad (2.44)$$

holds true.

Taking into account (2.35) and (2.37) one finds that $\mathcal{T} \circ \varphi$ is completely positive iff the matrices

$$\Delta_{kl} = \begin{bmatrix} \frac{1}{2}(C_{kl} + C_{lk}) + \mu & -\frac{i}{2}(C_{kl} - C_{lk}) \\ \frac{i}{2}(C_{kl} - C_{lk}) & \frac{1}{2}(C_{kl} + C_{lk}) - \mu \end{bmatrix} \quad (2.45)$$

are semipositive definite. The conditions (2.39)–(2.41) are equivalent to $\Delta_{kl} \geq 0$.

Corollary 3 The map

$$\varphi(a) = \sum_{i,j=1}^n c_{ij} e_{ij} a e_{ij}^* + \mu a \quad (2.46)$$

is completely positive and completely copositive, i.e. $\varphi \in P_n \cap P^n$ iff the following conditions

$$c_{ij} > 0, \quad (2.47)$$

$$c_{ij}c_{ji} \geq \mu^2, \quad i \neq j, \quad (2.48)$$

$$c_{ii} + \mu \geq 0, \quad (2.49)$$

$$[c_{ii}\delta_{ij} + \mu] \geq 0 \quad (2.50)$$

are satisfied.

The φ is not normalized, but since $\varphi(I_n) > 0$, the map

$$\psi(a) = \varphi(I_n)^{-1/2} \varphi(a) \varphi(I_n)^{-1/2} \quad (2.51)$$

is normalized, and the state

$$\sum_{i,j=1}^n \rho^{1/2} \psi(e_{ij}) \rho^{1/2} \otimes e_{ij} \quad (2.52)$$

is a PPT state.

Corollary 4 It follows from Theorems 1 and 2 that choosing $c_{ij} = k$, $1 \leq k < n$, and $\mu = -1$ the map

$$\varphi_k(a) = (k-1) \left(\sum_{i=1}^n e_{ii} a e_{ii} + \sum_{i<j} f_{ij} a f_{ij} \right) + (k+1) \sum_{i<j} g_{ij} a g_{ij} \quad (2.53)$$

is completely positive, while the map

$$(\tau \circ \varphi_k)(a) = kI_n(\text{tr } a) - a \quad (2.54)$$

is known to be k -positive but not $(k+1)$ -positive; i.e., φ_k is completely positive and k -copositive, and consequently the state

$$\frac{1}{kn-1} \sum_{i,j=1}^n \rho^{1/2} \varphi_k(e_{ij}) \rho^{1/2} \otimes e_{ij} \quad (2.55)$$

is NPT (by definition) provided $\text{rank } \rho = n$.

Using Theorems 1 and 2 specialized for the case $c_{ii} = c$ and $\mu = -1$.

Corollary 5 The map

$$\begin{aligned} \varphi(a) &= \sum_{i=1}^n (c-1) e_{ii} a e_{ii} + \sum_{i<j} \left\{ \left(\frac{1}{2}(c_{ij} + c_{ji}) - 1 \right) f_{ij} a f_{ij} \right. \\ &\quad \left. + \left(\frac{1}{2}(c_{ij} + c_{ji}) + 1 \right) g_{ij} a g_{ij} - \frac{i}{2}(c_{ij} - c_{ji}) f_{ij} a g_{ij} + \frac{i}{2}(c_{ij} - c_{ji}) g_{ij} a f_{ij} \right\} \end{aligned} \quad (2.56)$$

is completely positive iff $c \geq 1$ and $c_{ij}c_{ji} \geq 1 \quad i \neq j$.

On the other hand

$$(\tau \circ \varphi)(a) = c \sum_{i=1}^{n-1} h_i a h_i + \sum_{i \neq j} c_{ij} e_{ij} a e_{ij}^* - \frac{n-c}{n} a, \quad (2.57)$$

where $h_1, h_2, \dots, h_{n-1} \in M_n$, $h_i = h_i^*$, $\text{tr } h_i = 0$, $\text{tr } h_i h_j = \delta_{ij}$ and $\text{tr } h_i f_{kl} = \text{tr } h_i g_{kl} = 0$. Using the result of [9] one finds that $(\tau \circ \varphi)$ is k -positive but not $(k+1)$ -positive provided the following conditions are satisfied [10]

$$1 \leq k \leq c < k+1 < n, \quad (2.58)$$

$$\frac{k}{n-k} \leq \frac{c_{ij}}{n-c} < \frac{k+1}{n-k-1}. \quad (2.59)$$

In this case the map φ generates NPT states.

Acknowledgments

The work of M. A. and G.M. has been partially supported by a cooperation grant INFN-CICYT. The work of M. A. has also been partially supported by the Spanish MCyT grant FPA2000-1252. A. K. has been supported by the Grant PBZ-MIN-008/P03/2003.

References

- [1] M. Ohya, Lett. Nuovo Cimento **38**, 402 (1983).
- [2] M. Ohya, IEEE Trans. Information Theory **20**, 770 (1983).
- [3] N. J. Cerf, C. Adami, arXiv: quant-ph/9610005.
- [4] K. Życzkowski, I. Bengtson, Open Sys. Information Dyn. **11**, 3 (2004).
- [5] K. C. Ha, Publ. RIMS Univ. **34**, 59 (1998).
- [6] M. H. Eom, S. H. Kye, Math. Scan. **86**, 130 (2000).
- [7] K. C. Ha, S. H. Kye, V. S. Park, Phys. Lett. **A 313**, 163 (2003).
- [8] M. Horodecki, P. Horodecki, R. Horodecki, Phys. Lett. **A 223**, xxx (1996).
- [9] K. C. Ha, Publ. RIMS, Kyoto Univ. **34**, xxx (1998).
- [10] T. Takasaki, J. Tomiyama, Math. Z. **184**, 101 (1983).
- [11] P. Arrighi and C. Patricot, *On quantum operations as quantum states*, Ann. Phys. **311**, 26 (2004).